



FOR THE PEOPLE
FOR EDUCATION
FOR SCIENCE

LIBRARY
OF
THE AMERICAN MUSEUM
OF
NATURAL HISTORY

1918







A. M. ...
...
...
...

506 (97.3) 10
7

PROCEEDINGS

OF THE

Indiana Academy
of Science

1914

Collected 2/2/16
68

Y. L. L. L. L.
Y. L. L. L. L.
Y. L. L. L. L.

PROCEEDINGS

OF THE

Indiana Academy of Science

1914

H. E. BARNARD, Editor

INDIANAPOLIS :
WM. B. BURFORD, CONTRACTOR FOR STATE PRINTING AND BINDING
1915

18-75534-may 1

TABLE OF CONTENTS.

	PAGE
Constitution.....	5
By-Laws.....	7
Appropriation for 1913-1914.....	9
An Act for the Protection of Birds, Their Nests and Eggs.....	9
Public Offenses—Hunting Birds—Penalty.....	10
Officers, 1913-1914.....	11
Executive Committee.....	11
Curators.....	11
Committees Academy of Science, 1915.....	12
Officers of the Academy of Science (A Table of).....	13
Members.....	14
Fellows.....	14
Active Members.....	22
Minutes of the Spring Meeting.....	39
Minutes of the Thirtieth Annual Meeting.....	45
Program of the Thirtieth Annual Meeting.....	50
Science in Its Relation to the Conservation of Human Life. Severance Burrage.....	55
A Family in One Neighborhood. Amos W. Butler.....	59
The Problem of Feeble-Mindedness. G. S. Bliss.....	61
The Feeble-Minded and Delinquent Boy. Franklin C. Paschal.....	63
The Feeble-Minded and Delinquent Girl. E. E. Jones.....	71
Feeble-Mindedness in the Public Schools. Katrina Myers.....	79
The Alcohol Problem in the Light of Coniosis. Robert Hessler.....	85
Cold Storage Is Practical Conservation. H. E. Barnard.....	101
Changing Conditions in the Kentucky Mountains. B. H. Schockel.....	109
Conservation and Civilization. Arthur L. Foley.....	133
Why Do Our Birds Migrate? D. W. Demis.....	145
Flood Protection in Indiana. W. K. Hatt.....	149
An Apparatus for Aerating Culture Solutions. Paul Weatherwax.....	157
Antagonism on <i>B. Fluorescens</i> and <i>B. Typhosus</i> in Culture. P. A. Tetrault.....	161
Notes Upon the Distribution of Forest Trees in Indiana. Stanley Coulter.....	167
Corrections to the Lists of Mosses of Monroe County, Indiana, I and II. Mildred Nothnagel and F. L. Pickett.....	179
The Mosses of Monroe County, Indiana, III. F. L. Pickett and Mildred Nothnagel.....	181
A New Enemy of the Black Locust. Glenn Culbertson.....	185
A New Leaf Spot of <i>Viola Cucullata</i> . H. W. Anderson.....	187
Oat Smut in Indiana. F. J. Pipel.....	191
Plants New or Rare to Indiana. No. V. Charles C. Deam.....	197

	PAGE
Some Peculiarities in <i>Spirogyra Dubia</i> . Paul Weatherwax.....	203
Report on Corn Pollination IV. (Final.) M. L. Fisher.....	207
Stomata of <i>Trillium Nivale</i> . F. M. Andrews.....	209
The Primrose-Leaved Violet in White County. Louis M. Heimlich.....	213
Continuous Rust Propagation Without Sexual Reproduction. C. A. Ludwig.....	219
Correlation of Certain Long-Cyeled and Short-Cyeled Rusts. H. C. Travelbee.....	231
Some Species of <i>Nummularia</i> Common in Indiana. Claude E. O'Neal....	235
The Genus <i>Rosellinia</i> in Indiana. Glen B. Ramsey.....	251
Some Large Botanical Problems. G. C. Arthur.....	267
The Alba B. Ghere Collection of Birds' Eggs Presented to the Museum of Purdue University. Howard E. Enders.....	273
A Note on a Peculiar Nesting Site of the Chimney Swift. Glenn Culbertson	279
Notes on Indiana Earthworms. H. V. Heimburger.....	281
Notes on Orthoptera and Orthopteran Habitats in the Vicinity of Lafayette, Indiana. Henry Fox.....	287
Some Insects of the Between Tides Zone. Charles H. Arndt.....	323
The Snakes of the Lake Maxinkuckee Region. Barton Warren Evermann and Howard Walton Clark.....	337
Stirring as a Time Saver in Gravimetric Analysis. W. M. Blanchard....	349
The Alundum Crucible as a Substitute for the Gooch Crucible. George L. Clark.....	351
The Correlation of High School and College Chemistry. James Brown....	355
The Chemical Composition of Virgin and Cropped Indiana Soils. S. D. Conner.....	359
Sewage Disposal. Charles Brossman.....	365
Tar Forming Temperatures of American Coals. Otto Carter Berry.....	373
Shawnee Mound, Tippecanoe County, as a Glacial Alluvial Cone. Wm. A. McBeth.....	385
Correlation of the Outcrop at Spades, Indiana. H. N. Coryell.....	389
The Paleobotany of the Bloomington, Indiana, Quadrangle. T. F. Jackson.	395
The Flatwoods Region of Owen and Monroe Counties, Indiana. Clyde A. Malott.....	399
Mechanical Device for Testing Mersenne Numbers for Primes. Thos. E. Mason.....	429
Some Properties of Binomial Coefficients. A. M. Kenyon.....	433
Radioactivity of Spring Water. R. R. Ramsey.....	453
A Tornado at Watertown, South Dakota, June 23, 1914. J. Gladden Hutton.....	473
A Simple Form of the Carey Foster Bridge. J. P. Naylor.....	485
Variation of the Emanation Content of Certain Springs. R. R. Ramsey.	489
The Construction of a Rutherford's Electroscope. Edwin Morrison.....	491

CONSTITUTION.

ARTICLE I.

SECTION 1. This association shall be called the Indiana Academy of Science.

SEC. 2. The objects of this Academy shall be scientific research and the diffusion of knowledge concerning the various departments of science; to promote intercourse between men engaged in scientific work, especially in Indiana; to assist by investigation and discussion in developing and making known the material, educational and other resources and riches of the State; to arrange and prepare for publication such reports of investigation and discussions as may further the aims and objects of the Academy as set forth in these articles.

WHEREAS, The State has undertaken the publication of such proceedings, the Academy will, upon request of the Governor, or of one of the several departments of the State, through the Governor, act through its council as an advisory body in the direction and execution of any investigation within its province as stated. The necessary expenses incurred in the prosecution of such investigation are to be borne by the State; no pecuniary gain is to come to the Academy for its advice or direction of such investigation.

The regular proceedings of the Academy as published by the State shall become a public document.

ARTICLE II.

SECTION 1. Members of this Academy shall be honorary fellows, fellows, non-resident members or active members.

SEC. 2. Any person engaged in any department of scientific work, or in original research in any department of science, shall be eligible to active membership. Active members may be annual or life members. Annual members may be elected at any meeting of the Academy; they shall sign the constitution, pay an admission fee of two dollars and thereafter an annual fee of one dollar. Any person who shall at one time contribute fifty dollars to the funds of this Academy may be elected a life member of the Academy, free of assessment. Non-resident members may be elected

from those who have been active members but who have removed from the State. In any case, a three-fourths vote of the members present shall elect to membership. Application for membership in any of the foregoing classes shall be referred to a committee on application for membership, who shall consider such application and report to the Academy before the election.

Sec. 3. The members who are actively engaged in scientific work, who have recognized standing as scientific men, and who have been members of the Academy at least one year, may be recommended for nomination for election as fellows by three fellows or members personally acquainted with their work and character. Of members so nominated a number not exceeding five in one year may, on recommendation of the Executive Committee, be elected as fellows. At the meeting at which this is adopted, the members of the Executive Committee for 1894 and fifteen others shall be elected fellows, and those now honorary members shall become honorary fellows. Honorary fellows may be elected on account of special prominence in science, on the written recommendation of two members of the Academy. In any case a three-fourths vote of the members present shall elect.

ARTICLE III.

SECTION 1. The officers of this Academy shall be chosen by ballot at the annual meeting, and shall hold office one year. They shall consist of a President, Vice-President, Secretary, Assistant Secretary, Press Secretary and Treasurer, who shall perform the duties usually pertaining to their respective offices and in addition, with the ex-presidents of the Academy, shall constitute an Executive Committee. The President shall, at each annual meeting, appoint two members to be a committee, which shall prepare the programs and have charge of the arrangements for all meetings for one year.

Sec. 2. The annual meeting of this Academy shall be held in the city of Indianapolis within the week following Christmas of each year, unless otherwise ordered by the Executive Committee. There shall also be a summer meeting at such time and place as may be decided upon by the Executive Committee. Other meetings may be called at the discretion of the Executive Committee. The past Presidents, together with the officers and Executive Committee, shall constitute the council of the Academy, and represent it in the transaction of any necessary business not especially provided for in this constitution, in the interim between general meetings.

SEC. 3. This constitution may be altered or amended at any annual meeting by a three-fourths majority of the attending members of at least one year's standing. No question of amendment shall be decided on the day of its presentation.

BY-LAWS.

1. On motion, any special department of science shall be assigned to a curator, whose duty it shall be, with the assistance of the other members interested in the same department, to endeavor to advance knowledge in that particular department. Each curator shall report at such time and place as the Academy shall direct. These reports shall include a brief summary of the progress of the department during the year preceding the presentation of the report.

2. The President shall deliver a public address on the morning of one of the days of the meeting at the expiration of his term of office.

3. The Press Secretary shall attend to the securing of proper newspaper reports of the meetings and assist the Secretary.

4. No special meeting of the Academy shall be held without a notice of the same having been sent to the address of each member at least fifteen days before such meeting.

5. No bill against the Academy shall be paid without an order signed by the President and countersigned by the Secretary.

6. Members who shall allow their dues to remain unpaid for two years, having been annually notified of their arrearage by the Treasurer, shall have their names stricken from the roll.

7. Ten members shall constitute a quorum for the transaction of business.

AN ACT TO PROVIDE FOR THE PUBLICATION OF THE REPORTS AND PAPERS OF THE INDIANA ACADEMY OF SCIENCE.

[Approved March 11, 1895.]

WHEREAS, The Indiana Academy of Science, a chartered scientific association, has embodied in its constitution a provision that it will, upon the request of the Governor, or of the several departments of the State

government, through the Governor, and through its council as an advisory board, assist in the direction and execution of any investigation within its province, without pecuniary gain to the Academy, provided only that the necessary expenses of such investigation are borne by the State; and,

WHEREAS, The reports of the meetings of said Academy, with the several papers read before it, have very great educational, industrial and economic value, and should be preserved in permanent form; and

WHEREAS, The Constitution of the State makes it the duty of the General Assembly to encourage by all suitable means intellectual, scientific and agricultural improvement; therefore,

SECTION 1. *Be it enacted by the General Assembly of the State of Indiana,* That hereafter the annual reports of the meetings of the Indiana Academy of Science, beginning with the report for the year 1894, including all papers of scientific or economic value, presented at such meetings, after they shall have been edited and prepared for publication as hereinafter provided, shall be published by and under the direction of the Commissioners of Public Printing and Binding.

SEC. 2. Said reports shall be edited and prepared for publication without expense to the State, by a corps of editors to be selected and appointed by the Indiana Academy of Science, who shall not, by reason of such service, have any claim against the State for compensation. The form, style of binding, paper, typography and manner and extent of illustration of such reports shall be determined by the editors, subject to the approval of the Commissioners of Public Printing and Stationery. Not less than 1,500 nor more than 3,000 copies of each of said reports shall be published, the size of the edition within said limits to be determined by the concurrent action of the editors and the Commissioners of Public Printing and Stationery: *Provided,* That not to exceed six hundred dollars (\$600) shall be expended for such publication in any one year, and not to extend beyond 1896: *Provided,* That no sums shall be deemed to be appropriated for the year 1894.

SEC. 3. All except three hundred copies of each volume of said reports shall be placed in the custody of the State Librarian, who shall furnish one copy thereof to each public library in the State, one copy to each university, college or normal school in the State, one copy to each high school in the State having a library, which shall make application therefor, and one copy to such other institutions, societies or persons as may be designated by the

Academy through its editors or its council. The remaining three hundred copies shall be turned over to the Academy to be disposed of as it may determine. In order to provide for the preservation of the same it shall be the duty of the Custodian of the State House to provide and place at the disposal of the Academy one of the unoccupied rooms of the State House, to be designated as the office of the Academy of Science, wherein said copies of said reports belonging to the Academy, together with the original manuscripts, drawings, etc., thereof can be safely kept, and he shall also equip the same with the necessary shelving and furniture.

SEC. 4. An emergency is hereby declared to exist for the immediate taking effect of this act, and it shall therefore take effect and be in force from and after its passage.

APPROPRIATION FOR 1913-1914.

The appropriation for the publication of the proceedings of the Academy during the years 1913 and 1914 was increased by the Legislature in the General Appropriation bill, approved March 9, 1909. That portion of the law fixing the amount of the appropriation for the Academy is herewith given in full:

For the Academy of Science: For the printing of the proceedings of the Indiana Academy of Science twelve hundred dollars: *Provided*, That any unexpended balance in 1913 shall be available in 1914, and that any unexpended balance in 1914 shall be available in 1915.

AN ACT FOR THE PROTECTION OF BIRDS, THEIR NESTS AND EGGS.

SEC. 602. Whoever kills, traps or has in his possession any wild bird, or whoever sells or offers the same for sale, or whoever destroys the nest or eggs of any wild bird, shall be deemed guilty of a misdemeanor and upon conviction thereof shall be fined not less than ten dollars nor more than twenty-five dollars: *Provided*, That the provisions of this section shall not apply to the following named game birds: The Anatidæ, commonly called swans, geese, brant, river and sea duck; the Rallidæ, commonly called rails, coots, mud-hens, gallinules; the Limicolæ, commonly called shore birds, surf birds, plover, snipe, woodcock, sandpipers, tattlers and curlew; the Gallinæ, commonly called wild turkeys, grouse, prairie chickens, quails and pheasants; nor to English or European house sparrows,

crows, hawks or other birds of prey. Nor shall this section apply to persons taking birds, their nests or eggs, for scientific purposes, under permit, as provided in the next session.

SEC. 603. Permits may be granted by the Commissioner of Fisheries and Game to any properly accredited person, permitting the holder thereof to collect birds, their nests or eggs for strictly scientific purposes. In order to obtain such permit the applicant for the same must present to such Commissioner written testimonials from two well-known scientific men certifying to the good character and fitness of such applicant to be entrusted with such privilege, and pay to such Commissioner one dollar therefor and file with him a properly executed bond in the sum of two hundred dollars, payable to the State of Indiana, conditioned that he will obey the terms of such permit, and signed by at least two responsible citizens of the State as sureties. The bond may be forfeited, and the permit revoked upon proof to the satisfaction of such Commissioner that the holder of such permit has killed any bird or taken the nest or eggs of any bird for any other purpose than that named in this section.

PUBLIC OFFENSES—HUNTING WILD BIRDS—PENALTY.

[Approved March 13, 1913.]

SECTION 1. *Be it enacted by the General Assembly of the State of Indiana,* That section six (6) of the above entitled act be amended to read as follows: Section 6. That section six hundred two (602) of the above entitled act be amended to read as follows: Section 602. It shall be unlawful for any person to kill, trap or possess any wild bird, or to purchase or offer the same for sale, or to destroy the nest or eggs of any wild bird, except as otherwise provided in this section. But this section shall not apply to the following named game birds: The Anatidæ, commonly called swans, geese, brant, river and sea duck; the Rallidæ, commonly known as rails, coots, mud-hens and gallinules; the Limicolæ, commonly known as shore birds, plovers, surf birds, snipe, woodcock, sandpipers, tattlers and curlews; the Gallinæ, commonly called wild turkeys, grouse, prairie chickens, quails, and pheasants; nor to English or European house sparrows, blackbirds, crows, hawks or other birds of prey. Nor shall this section apply to any person taking birds or their nests or eggs for scientific purposes under permit as provided in the next section. Any person violating the provisions of this section shall, on conviction, be fined not less than ten dollars (\$10.00) nor more than fifty dollars (\$50.00).

Indiana Academy of Science.

OFFICERS, 1914-1915.

PRESIDENT,
WILBUR A. COGSHALL.
VICE-PRESIDENT,
WILLIAM A. MCBETH.
SECRETARY,
ANDREW J. BIGNEY.
ASSISTANT SECRETARY,
H. E. ENDERS.
PRESS SECRETARY,
FRANK B. WADE.
TREASURER,
WILLIAM M. BLANCHARD.
EDITOR,
H. E. BARNARD.

EXECUTIVE COMMITTEE:

ARTHUR, J. C.,	CULBERTSON, GLENN,	MCBETH, W. A.,
BIGNEY, A. J.,	DRYER, CHAS. R.,	MEES, CARL L.,
BLANCHARD, W. M.,	EIGENMANN, C. H.,	MOTTIER, DAVID M.,
BLATCHLEY, W. S.,	EVANS, P. N.,	MENDENHALL, T. C.,
BODINE, DONALDSON,	DENNIS, D. W.,	NAYLOR, JOSEPH P.,
BRANNER, J. C.,	FOLEY, A. L.,	NOYES, W. A.,
BURRAGE, SEVERANCE,	HAY, O. P.,	WADE, F. B.,
BUTLER, AMOS W.,	HESSLER, ROBERT.	WALDO, C. A.,
COGSHALL, W. A.,	JOHN, J. P. D.,	WILEY, H. W.,
COULTER, JOHN M.,	JORDAN, D. S.,	WRIGHT, JOHN S.
COULTER, STANLEY,		

CURATORS:

BOTANY.....	J. C. ARTHUR.
ENTOMOLOGY.....	W. S. BLATCHLEY.
HERPETOLOGY }.....	
MAMMALOGY }.....	A. W. BUTLER.
ORNITHOLOGY }.....	
ICHTHYOLOGY.....	H. C. EIGENMANN.

COMMITTEES ACADEMY OF SCIENCE, 1915.

Program.

WILL SCOTT, Bloomington
 F. B. WADE, Indianapolis
 P. N. EVANS, LaFayette

Nominations.

SEVERANCE BURRAGE, Indianapolis
 W. J. MOENKHAUS, Bloomington
 A. S. HATHAWAY, Terre Haute

State Library.

W. S. BLATCHLEY, Indianapolis
 H. J. BANKER, Coldspring Harbor,
 N. Y.
 A. W. BUTLER, Indianapolis

Biological Survey.

C. C. DEAM, Bluffton
 H. W. ANDERSON, Crawfordsville
 GEO. N. HOFFER, West LaFayette
 A. O. COX, Terre Haute
 J. A. NIEUWLAND, Notre Dame

Distribution of Proceedings.

A. J. BIGNEY, Moores Hill
 JOHN B. DUTCHER, Bloomington
 A. W. BUTLER, Indianapolis
 W. M. BLANCHARD, Greencastle

Membership.

H. E. ENDERS, West LaFayette
 EDWIN MORRISON, Richmond.
 FRED A. MILLER, Indianapolis

Auditing.

E. B. WILLIAMSON, Bluffton
 GLENN CULBERTSON, Hanover

Restriction of Weeds and Diseases.

ROBERT HESSLER, Logansport
 AMOS BUTLER, Indianapolis
 J. N. HURTY, Indianapolis
 STANLEY COULTER, LaFayette
 D. M. MOTTIER, Bloomington

Academy to State.

R. W. MCBRIDE, Indianapolis
 GLENN CULBERTSON, Hanover
 H. E. BARNARD, Indianapolis
 A. W. BUTLER, Indianapolis
 W. W. WOOLLEN, Indianapolis

Publication of Proceedings.

H. E. BARNARD, Indianapolis
 C. R. DRYER, Ft. Wayne
 M. K. HAGGERTY, Bloomington
 R. R. HYDE, Terre Haute
 J. S. WRIGHT, Indianapolis

OFFICERS OF THE INDIANA ACADEMY OF SCIENCE.

YEARS.	PRESIDENT.	SECRETARY.	ASST. SECRETARY.	PRESS SECRETARY.	TREASURER.
1885-1886	David S. Jordan.	Amos W. Butler.			O. P. Jenkins.
1886-1887	John M. Coulter.	Amos W. Butler.			O. P. Jenkins.
1887-1888	J. P. D. John.	Amos W. Butler.			O. P. Jenkins.
1888-1889	John C. Branner.	Amos W. Butler.			O. P. Jenkins.
1889-1890	T. C. Mendenhall.	Amos W. Butler.			O. P. Jenkins.
1890-1891	O. P. Hay.	Amos W. Butler.			C. A. Waldo.
1891-1892	J. L. Campbell*	Amos W. Butler.			C. A. Waldo.
1892-1893	J. C. Arthur.	Amos W. Butler.	{Stanley Coulter.		C. A. Waldo.
1893-1894	W. A. Noyes.	C. A. Waldo.	W. W. Norman.		W. P. Shannon.
1894-1895	A. W. Butler.	John S. Wright.	A. J. Bigney.		W. P. Shannon.
1895-1896	Stanley Coulter.	John S. Wright.	A. J. Bigney.		W. P. Shannon.
1896-1897	Thomas Gray*.	John S. Wright.	A. J. Bigney.		J. T. Scovell.
1897-1898	C. A. Waldo.	John S. Wright.	A. J. Bigney.	Geo. W. Benton.	J. T. Scovell.
1898-1899	C. H. Eigenmann.	John S. Wright.	E. A. Schultze.	Geo. W. Benton.	J. T. Scovell.
1899-1900	D. W. Dennis.	John S. Wright.	E. A. Schultze.	Geo. W. Benton.	J. T. Scovell.
1900-1901	M. B. Thomas*.	John S. Wright.	E. A. Schultze.	Geo. W. Benton.	J. T. Scovell.
1901-1902	Harvey W. Wiley.	John S. Wright.	Donaldson Bodine.	Geo. W. Benton.	J. T. Scovell.
1902-1903	W. S. Blachley.	John S. Wright.	Donaldson Bodine.	G. A. Abbott.	W. A. McBeth.
1903-1904	C. L. Mees.	John S. Wright.	J. H. Ransom.	G. A. Abbott.	W. A. McBeth.
1904-1905	John S. Wright.	Lynn B. McMullen.	J. H. Ransom.	G. A. Abbott.	W. A. McBeth.
1905-1906	Robert Hessler.	Lynn B. McMullen.	J. H. Ransom.	G. A. Abbott.	W. A. McBeth.
1906-1907	D. M. Mottier.	Lynn B. McMullen.	J. M. Ransom.	Charles R. Clark.	W. A. McBeth.
1907-1908	Glenn Culbertson.	J. H. Ransom.	A. J. Bigney.	G. A. Abbott.	W. A. McBeth.
1908-1909	A. L. Foley.	J. H. Ransom.	A. J. Bigney.	G. A. Abbott.	W. A. McBeth.
1909-1910	P. N. Evans.	Geo. W. Benton.	A. J. Bigney.	John W. Woodhams.	W. J. Moenkhaus.
1910-1911	C. R. Dyer.	A. J. Bigney.	E. B. Williamson.	Milo H. Stuart.	W. J. Moenkhaus.
1911-1912	J. P. Naylor.	A. J. Bigney.	E. B. Williamson.	Milo H. Stuart.	W. J. Moenkhaus.
1912-1913	Donaldson Bodine.	A. J. Bigney.	C. M. Smith.	F. B. Wade.	W. A. Cogshall.
1913-1914	Severance Burrage.	A. J. Bigney.	H. E. Enders.	F. B. Wade.	W. A. Cogshall.
1914-1915	Wilbur A. Cogshall.	A. J. Bigney.	H. E. Enders.	F. B. Wade.	W. M. Blanchard.

*Deceased.

MEMBERS.*

FELLOWS.

- ††Abbott, G. A., Grand Forks, N. Dak. †1908
 Professor of Chemistry, University of North Dakota.
 Chemistry.
- Aley, Robert J., Orono, Me. 1898
 President of University of Maine.
 Mathematics and General Science.
- Anderson, H. W., 1 Mills Place, Crawfordsville, Ind. 1912
 Professor of Botany, Wabash College.
 Botany.
- Andrews, F. M., 744 E. Third St., Bloomington, Ind. 1911
 Assistant Professor of Botany, Indiana University.
 Botany.
- Arthur, Joseph C., 915 Columbia St., Lafayette, Ind. 1894
 Professor of Vegetable Physiology and Pathology, Purdue University.
 Botany.
- Barnard, H. E., Room 20 State House, Indianapolis, Ind. 1910
 Chemist to Indiana State Board of Health.
 Chemistry, Sanitary Science, Pure Foods.
- Beede, Joshua W., cor. Wall and Atwater Sts., Bloomington, Ind. 1896
 Associate Professor of Geology, Indiana University.
 Stratigraphic Geology, Physiography.
- Benton, George W., 100 Washington Square, New York, N. Y. 1896
 With the American Book Company.

*Every effort has been made to obtain the correct address and occupation of each member, and to learn what line of science he is interested in. The first line contains the name and address; the second line the occupation; the third line the branch of science in which he is interested. The omission of an address indicates that mail addressed to the last printed address was returned as uncalled for. Information as to the present address of members so indicated is requested by the secretary. The custom of dividing the list of members has been followed.

†Date of election,

††Non-resident,

- Bigney, Andrew J., Moores Hill, Ind. 1897
 Vice-President and Professor of Biology and Geology, Moores Hill
 College.
 Biology and Geology.
- Bitting, Catharine Golden, Washington, D. C. 1895
 Microscopic Expert, Pure Food, National Cammers Laboratory.
 Botany.
- Blatchley, W. S., 1558 Park Ave., Indianapolis, Ind. 1893
 Naturalist.
 Botany, Entomology and Geology.
- Bodine, Donaldson, Four Mills Place, Crawfordsville, Ind. 1899
 Professor of Geology and Zoology, Wabash College.
 Entomology and Geology.
- Breeze, Fred J., care American Book Company, New York, N. Y. 1910
 With the American Book Company.
 Geography.
- Bruner, Henry Lane, 324 S. Ritter Ave., Indianapolis, Ind. 1899
 Professor of Biology, Butler College.
 Comparative Anatomy, Zoology.
- Burrage, Severance, care Eli Lilly Co., Indianapolis, Ind. 1898
 Charge of Biological Laboratory, Eli Lilly Co.
 Bacteriology, Sanitary Science.
- Butler, Amos W., 52 Downey Ave., Irvington, Ind. 1893
 Secretary, Indiana Board of State Charities.
 Vertebrate Zoology, Anthropology, Sociology.
- Cogshall, Wilbur A., 423 S. Fess Ave., Bloomington, Ind. 1906
 Associate Professor of Astronomy, Indiana University.
 Astronomy and Physics.
- Cook, Mel T., New Brunswick, N. J. 1902
 Professor of Plant Pathology, Rutgers College.
 Botany, Plant Pathology, Entomology.
- Coulter, John M., care University of Chicago, Chicago, Ill. 1893
 Head Department of Botany, Chicago University.
 Botany.
- Coulter, Stanley, 213 S. Ninth St., Lafayette, Ind. 1893
 Dean School of Science, Purdue University.
 Botany, Forestry.

- Cox, Ulysses O., P. O. Box 81, Terre Haute, Ind..... 1908
 Head Department Zoology and Botany, Indiana State Normal.
 Botany, Zoology.
- Culbertson, Glenn, Hanover, Ind..... 1899
 Chair Geology, Physics and Astronomy, Hanover College.
 Geology.
- Cumings, Edgar Roscoe, 327 E. Second St., Bloomington, Ind..... 1906
 Professor of Geology, Indiana University.
 Geology, Paleontology.
- Davisson, Schuyler Colfax, Bloomington, Ind..... 1908
 Professor of Mathematics, Indiana University.
 Mathematics.
- Deam, Charles C., Bluffton, Ind..... 1910
 Druggist.
 Botany.
- Dennis, David Worth, Richmond, Ind..... 1895
 Professor of Biology, Earlham College.
 Biology.
- Dryer, Charles R., Oak Knoll, Fort Wayne, Ind..... 1897
 Geographer.
- Eigenmann, Carl H., 630 Atwater St., Bloomington, Ind..... 1893
 Professor Zoology, Dean of Graduate School, Indiana University.
 Embryology, Degeneration, Heredity, Evolution and Distribution
 of American Fish.
- Enders, Howard Edwin, 105 Quincy St., Lafayette, Ind..... 1912
 Associate Professor of Zoology, Purdue University.
 Zoology.
- Evans, Percy Norton, Lafayette, Ind..... 1901
 Director of Chemical Laboratory, Purdue University.
 Chemistry.
- Foley, Arthur L., Bloomington, Ind..... 1897
 Head of Department of Physics, Indiana University.
 Physics.
- Golden, M. J., Lafayette, Ind..... 1899
 Director of Laboratories of Practical Mechanics, Purdue Uni-
 versity.
 Mechanics.

- ††Goss, William Freeman M., Urbana, Ill..... 1893
Dean of College of Engineering, University of Illinois.
- Haggerty, M. E., Bloomington, Ind..... 1913
- Hathaway, Arthur S., 2206 N. Tenth St., Terre Haute, Ind..... 1895
Professor of Mathematics, Rose Polytechnic Institute.
Mathematics, Physics.
- Hessler, Robert, Logansport, Ind..... 1899
Physician.
Biology.
- Hilliard, C. M., Simmons College, Boston, Mass..... 1913
- Hoffer, Geo. N., West Lafayette, Ind..... 1913
- Hurty, J. N., Indianapolis, Ind..... 1910
Secretary, Indiana State Board of Health.
Sanitary Science, Vital Statistics, Eugenics.
- †Huston, H. A., New York City..... 1893
- Kern, Frank D., State College, Pa..... 1912
Professor of Botany, Pennsylvania State College.
Botany.
- Lyons, Robert E., 630 E. Third St., Bloomington, Ind..... 1896
Head of Department of Chemistry, Indiana University.
Organic and Biological Chemistry.
- McBeth, William A., 1905 N. Eighth St., Terre Haute, Ind..... 1904
Assistant Professor Geography, Indiana State Normal.
Geography, Geology, Scientific Agriculture.
- †Marsters, V. F., Santiago, Chile..... 1893
- Mees, C. L., Terre Haute, Ind..... 1894
President of Rose Polytechnic Institute.
- †Miller, John Anthony, Swarthmore, Pa..... 1904
Professor of Mathematics and Astronomy, Swarthmore College.
Astronomy, Mathematics.
- Moenkhaus, William J., 501 Fess Ave., Bloomington, Ind..... 1901
Professor of Physiology, Indiana University.
Physiology.
- Moore, Richard B., Denver, Colo..... 1893
With U. S. Bureau of Mines.
Chemistry, Radio-activity.

- Mottier, David M., 215 Forest Place, Bloomington, Ind..... 1893
 Professor of Botany, Indiana University.
 Morphology, Cytology.
- Naylor, J. P., Greencastle, Ind..... 1903
 Professor of Physics, Depauw University.
 Physics, Mathematics.
- †Noyes, William Albert, Urbana, Ill..... 1893
 Director of Chemical Laboratory, University of Illinois.
 Chemistry.
- Pohlman, Augustus G., 1100 E. Second St., Bloomington, Ind..... 1911
 Professor of Anatomy, Indiana University.
 Embryology, Comparative Anatomy.
- Ramsey, Rolla R., 615 E. Third St., Bloomington, Ind..... 1906
 Associate Professor of Physics, Indiana University.
 Physics.
- Ransom, James H., 323 University St., West Lafayette, Ind..... 1902
 Professor of General Chemistry, Purdue University.
 General Chemistry, Organic Chemistry, Teaching.
- Rettger, Louis J., 31 Gilbert Ave., Terre Haute, Ind..... 1896
 Professor of Physiology, Indiana State Normal.
 Animal Physiology.
- Rothrock, David A., Bloomington, Ind..... 1906
 Professor of Mathematics, Indiana University.
 Mathematics.
- Scott, Will, 731 Atwater St., Bloomington, Ind..... 1911
 Assistant Professor of Zoology, Indiana University.
 Zoology, Lake Problems.
- Shannon, Charles W., Norman, Okla..... 1912
 With Oklahoma State Geological Survey.
 Soil Survey, Botany.
- Smith, Albert, 1022 Seventh St., West Lafayette..... 1908
 Professor of Structural Engineering.
 Physics, Mechanics.
- ††Smith, Alexander, care Columbia University, New York, N. Y..... 1893
 Head of Department of Chemistry, Columbia University.
 Chemistry.

- Smith, Charles Marquis, 910 S. Ninth St., Lafayette, Ind..... 1912
 Professor of Physics, Purdue University.
 Physics.
- Stone Winthrop E. Lafayette, Ind..... 1893
 President of Purdue University.
 Chemistry.
- ††Swain, Joseph, Swarthmore, Pa..... 1898
 President of Swarthmore College.
 Science of Administration.
- Van Hook, James M., 639 N. College Ave., Bloomington, Ind..... 1911
 Assistant Professor of Botany, Indiana University.
 Botany.
- ††Waldo, Clarence A., care Washington University, St. Louis, Mo.... 1893
 Thayer Professor Mathematics and Applied Mechanics, Wash-
 ington University.
 Mathematics, Mechanics, Geology and Mineralogy.
- ††Webster, F. M., Kensington, Md..... 1894
 Entomologist, U. S. Department of Agriculture, Washington, D. C.
 Entomology.
- Westland, Jacob, 439 Salisbury St., West Lafayette, Ind..... 1904
 Professor of Mathematics, Purdue University.
 Mathematics.
- Wiley, Harvey W., Cosmos Club, Washington, D. C..... 1895
 Professor of Agricultural Chemistry, George Washington Uni-
 versity.
 Biological and Agricultural Chemistry.
- Woollen, William Watson, Indianapolis, Ind..... 1908
 Lawyer.
 Birds and Nature Study.
- Wright, John S., care Eli Lilly Co., Indianapolis, Ind..... 1894
 Manager of Advertising Department, Eli Lilly Co.
 Botany.

NON-RESIDENT MEMBERS.

Ashley, George H., Washington, D. C.

Bain, H. Foster, London, England.

Editor Mining Magazine.

Branner, John Casper, Stanford University, California.

Vice-President of Stanford University, and Professor of Geology.
Geology.

Brannon, Melvin A., President University of Idaho, Boise, Ida.

Professor of Botany.

Plant Breeding.

Campbell, D. H., Stanford University, California.

Professor of Botany, Stanford University.

Botany.

Clark, Howard Walton, U. S. Biological Station, Fairport, Iowa.

Scientific Assistant, U. S. Bureau of Fisheries.

Botany, Zoology.

Dorner, H. B., Urbana, Illinois.

Assistant Professor of Floriculture.

Botany, Floriculture.

Duff, A. Wilmer, 43 Harvard St., Worcester, Mass.

Professor of Physics, Worcester Polytechnic Institute.

Physics.

Evermann, Barton Warren, Director Museum.

California Academy of Science, Golden Gate Park, San Francisco, Cal.

Zoology.

Fiske, W. A., Los Angeles, Cal., Methodist College.

Garrett, Chas. W., Room 718 Pennsylvania Station, Pittsburgh, Pa.

Librarian, Pennsylvania Lines West of Pittsburgh.

Entomology, Sanitary Sciences.

Gilbert, Charles H., Stanford University, California.

Professor of Zoology, Stanford University.

Ichthyology.

Greene, Charles Wilson, 814 Virginia Ave., Columbia, Mo.

Professor of Physiology and Pharmacology, University of Missouri.

Physiology, Zoology.

- Hargitt, Chas. W., 909 Walnut Ave., Syracuse, N. Y.
 Professor of Zoology, Syracuse University.
 Hygiene, Embryology, Eugenics, Animal Behavior.
- Hay, Oliver Perry, U. S. National Museum, Washington, D. C.
 Research Associate, Carnegie Institution of Washington.
 Vertebrate Paleontology, especially that of the Pleistocene Epoch.
- Hughes, Edward, Stockton, California.
- Jenkins, Oliver P., Stanford University, California.
 Professor of Physiology, Stanford University.
 Physiology, Histology.
- Jordan, David Starr, Stanford University, California.
 President Emeritus of Stanford University.
 Fish, Eugenics, Botany, Evolution.
- Kingsley, J. S., University of Illinois, Champaign, Ill.
 Professor of Zoology.
 Zoology.
- Knipp, Charles T., 913 W. Nevada St., Urbana, Illinois.
 Assistant Professor of Physics, University of Illinois.
 Physics, Discharge of Electricity through Gases.
- MacDougal, Daniel Trembly, Tucson, Arizona.
 Director, Department of Botanical Research, Carnegie Institute, Wash-
 ington, D. C.
 Botany.
- McMullen, Lynn Banks, State Normal School, Valley City, North Dakota.
 Head Science Department, State Normal School.
 Physics, Chemistry.
- Mendenhall, Thomas Corwin, Ravenna, Ohio.
 Retired.
 Physics. "Engineering," Mathematics, Astronomy.
- Moore, George T., St. Louis, Mo.
 Director Missouri Botanical Garden.
- Newsom, J. F., Palo Alto, California.
 Mining Engineer.
- Purdue, Albert Homer, State Geological Survey, Nashville, Tenn.
 State Geologist of Tennessee.
 Geology.

Reagan, A. B.

Superintendent Deer Creek Indian School, Ibapah, Utah.
Geology, Paleontology, Ethnology.

Slonaker, James Rollin, 334 Kingsley Ave., Palo Alto, California.

Assistant Professor of Physiology, Stanford University.
Physiology, Zoology.

Springer, Alfred, 312 East 2d St., Cincinnati, Ohio.

Chemist.
Chemistry.

ACTIVE MEMBERS.

Allen, William Ray, Bloomington, Ind.

Allison, Evelyn, Lafayette, Ind.

Care Agricultural Experiment Station.
Botany.

Anderson, Flora Charlotte, Wellesley College, Wellesley, Mass.

Botany.

Arndt, Charles H., Lafayette, Ind.

Biology.

Atkinson, F. C., Indianapolis.

Chemistry.

Badertscher, J. A., Bloomington.

Anatomy.

Baker, George A., South Bend.

Archaeology.

Baker, Walter D., N. Illinois St., Indianapolis, Ind.

Care Waldcraft Co.
Chemistry.

Baker, Walter M., Amboy.

Superintendent of Schools.
Mathematics and Physics.

Baker, William Franklin, Indianapolis.

Medicine.

Balcom, H. C., Indianapolis.

Botany.

- Banker, Howard J., Cold Spring Harbor, N. Y.
Botany.
- Barcus, H. H., Indianapolis.
Instructor, Mathematics, Shortridge High School.
- Barr, Harry L., Waveland.
Student.
Botany and Forestry.
- Barrett, Edward, Indianapolis.
State Geologist.
Geology, Soil Survey.
- Bates, W. H., 306 Russell St., West Lafayette.
Associate Professor, Mathematics.
- Bell, Guido, 431 E. Ohio St., Indianapolis.
Physician.
- Bellamy, Ray, Worcester, Mass.
- Bennett, Lee F., 825 Laporte Ave., Valparaiso.
Professor of Geology and Zoology, Valparaiso University.
Geology, Zoology.
- Berry, O. C., West Lafayette.
Engineering.
- Berteling, John B., South Bend.
Medicine.
- Binford, Harry, Earlham.
Zoology.
- Bishop, Harry Eldridge, 1706 College Ave., Indianapolis.
Food Chemist, Indiana State Board of Health.
- Blanchard, William M., 1608 S. College Ave., Greencastle.
Professor of Chemistry, DePauw University.
Organic Chemistry.
- Blew, Michael James, R. R. 1, Wabash.
Chemistry and Botany.
- Bliss, G. S., Ft. Wayne.
Medicine.
- Bond, Charles S., 112 N. Tenth St., Richmond.
Physician.
Biology, Bacteriology, Physical Diagnosis and Photomicrography.

- Bourke, A. Adolphus, 1103 Cottage Ave., Columbus.
Instructor, Physics, Zoology and Geography.
Botany, Physics.
- Bowles, Adam L., Terre Haute.
Zoology.
- Bowers, Paul E., Michigan City.
Medicine.
- Breckinridge, James M., Crawfordsville.
Chemistry.
- Brossmann, Charles, 1616 Merchants Bank Bldg., Indianapolis.
Consulting Engineer.
Water Supply, Sewage Disposal, Sanitary Engineering, etc.
- Brown, James, 5372 E. Washington St., Indianapolis.
Professor of Chemistry, Butler College.
Chemistry.
- Brown, Paul H., Richmond.
Physics.
- Brown, Hugh E., Bloomington, Ind.
- Bruce, Edwin M., 2401 North Ninth St., Terre Haute.
Assistant Professor of Physics and Chemistry, Indiana State Normal.
Chemistry, Physics.
- Bryan, William Lowe, Bloomington.
President, Indiana University.
Psychology.
- Bybee, Halbert P., Bloomington.
Graduate Student, Indiana University.
Geology.
- Canis, Edward N., 2221 Park Ave., Indianapolis.
Officeman with William B. Burford.
Botany, Psychology.
- Caparo, Jose Angel, Notre Dame.
Physics and Mathematics.
- Carlyle, Paul J., Bloomington.
Chemistry.
- Carmichael, R. D., Bloomington.
Assistant Professor of Mathematics, Indiana University.
Mathematics.

- Carr, Ralph Howard, Lafayette.
Chemistry.
- Caswell, Albert E., Lafayette.
Instructor in Physics, Purdue University.
Physics and Applied Mathematics.
- Chansler, Elias J., Bicknell.
Farmer.
Ornithology and Mammals.
- Clark, George Lindenburg, Greencastle, DePauw University.
Chemistry.
- Clark, Elbert Howard, West Lafayette.
Mathematics.
- Clark, Jediah H., 126 East Fourth St., Connersville.
Physician.
Medicine.
- Clarke, Elton Russell, Indianapolis.
Zoology.
- Collins, Jacob Roland, West Lafayette, Purdue University.
Instructor in Physics.
- Conner, S. D., West Lafayette.
- Coryell, Noble H., Bloomington.
Chemistry.
- Cotton, William J., 5363 University Ave., Indianapolis.
Physics and Chemistry.
- Cox, William Clifford.
- Crowell, Melvin E., 648 E. Monroe St., Franklin.
Dean of Franklin College.
Chemistry and Physics.
- Cutter, George, Broad Branch Road, Washington, D. C.
Retired Manufacturer of Electrical Supplies.
Conchology.
- Daniels, Lorenzo E., Rolling Prairie.
Retired Farmer.
Conchology.
- Davis, D. W., Greencastle.
Biology.

Davis, Ernest A., Notre Dame.

Chemistry.

Davis, Melvin K., Anderson.

Instructor, Anderson High School.

Physiography, Geology, Climatology.

Dean, John C., Indianapolis.

Astronomy.

Deppe, C. R., Franklin.

Dewey, Albert H., West Lafayette.

Department of Pharmacy, Purdue University.

Dietz, Harry F., 408 W. Twenty-eighth St., Indianapolis.

Deputy State Entomologist.

Entomology, Eugenics, Parasitology, Plant Pathology.

Dolan, Jos. P., Syracuse.

Donaghy, Fred, Ossian.

Botany.

Dostal, Bernard F., Bloomington.

Physics.

Drew, David Abbott, 817 East Second St., Bloomington.

Instructor in Mechanics and Astronomy.

Astronomy, Mechanics, Mathematics and Applied Mathematics.

DuBois, Henry, Bloomington, Ind.

Duden, Hans A., 5050 E. Washington St., Indianapolis.

Analytical Chemist.

Chemistry.

Duncan, David Christie, West Lafayette.

Instructor in Physics, Purdue University.

Dutcher, J. B., Bloomington.

Assistant Professor of Physics, Indiana University.

Physics.

Earp, Samuel E., 24½ Kentucky Ave., Indianapolis.

Physician.

Easley, Mary, Bloomington, Ind.

Edmonston, Clarence E., Bloomington.

Graduate Student, Physiology, Indiana University.

Physiology.

- Edwards, Carlton, Earlham College, Earlham.
- Ellis, Max Mapes, Boulder, Colo.
Instructor in Biology, University of Colorado.
Biology, Entomology.
- Emerson, Charles P., Hume Mansur Bldg., Indianapolis.
Dean Indiana University Medical College.
Medicine.
- Evans, Samuel G., 1452 Upper Second St., Evansville.
Merchant.
Botany, Ornithology.
- Ewers, James E., Terre Haute.
Instructor in High School.
Geology.
- Felver, William P., 325½ Market St., Logansport.
Railroad Clerk.
Geology, Chemistry.
- Ferry, Oliver P., West Lafayette.
Physiology.
- Fisher, Homer Glenn, Bloomington.
Zoology.
- Fisher, Martin L., Lafayette.
Professor of Crop Production, Purdue University.
Agriculture, Soils and Crops, Birds, Botany.
- Foresman, George Kedgie, Lafayette, Purdue University.
Chemistry.
- Frier, George M., Lafayette.
Assistant Superintendent, Agricultural Experiment Station, Purdue
University.
Botany, Zoology, Entomology, Ornithology, Geology.
- Fulk, Murl E., Decatur.
Anatomy.
- Fuller, Frederick D., 213 Russell St., West Lafayette.
Chief Deputy State Chemist, Purdue Experiment Station.
Chemistry, Microscopy.
- Funk, Austin, 404 Spring St., Jeffersonville.
Physician.
Diseases of Eye, Ear, Nose and Throat.

- Galloway, Jesse James, Bloomington.
 Instruction, Indiana University.
 Geology, Paleontology.
- Garner, J. B., Mellon Institute, Pittsburgh, Pa.
 Chemistry.
- Gatch, Willis D., Indianapolis, Indiana University Medical School.
 Anatomy.
- Gates, Florence A., 3435 Detroit Ave., Toledo, Ohio.
 Teacher of Botany.
 Botany and Zoology.
- Gidley, William, West Lafayette.
 Department of Pharmacy, Purdue University.
- Gillum, Robert G., Terre Haute, Ind.
- Glenn, E. R., Froebel School, Gary, Ind.
 Physics.
- Goldsmith, William Marion, Oakland City.
 Zoology.
- Gottlieb, Frederic W., Morristown.
 Care Museum of Natural History. Assistant Curator, Moores Hill
 College.
 Archaeology, Ethnology.
- Grantham, Guy E., 437 Vine St., West Lafayette.
 Instructor in Physics, Purdue University.
- Greene, Frank C., Missouri Bureau of Geology and Mines, Rolla, Mo.
 Geologist.
 Geology.
- Grimes, Earl J., Russellville.
 Care U. S. Soil Survey.
 Botany, Soil Survey.
- Hamill, Samuel Hugh, 119 E. Fourth St., Bloomington.
 Chemistry.
- Hammerschmidt, Louis M., South Bend.
 Science of Law.
- Happ, William, South Bend.
 Botany.

- Harding, C. Francis, 111 Fowler Ave., West Lafayette.
 Professor of Electrical Engineering, Purdue University.
 Mathematics, Physics, Chemistry.
- Harman, Mary T., 611 Laramie St., Manhattan, Kansas.
 Instructor in Zoology, Kansas State Agricultural College.
 Zoology.
- Harman, Paul M., Bloomington.
 Geology.
- Harvey, R. B., Indianapolis.
- Heimbürger, Harry V., 701 West Washington St., Urbana, Ill.
 Assistant in Zoology, University of Illinois.
- Heimlich, Louis Frederick, 703 North St., Lafayette.
 Biology.
- Hendricks, Victor K., 855 Benton Ave., Springfield, Mo.
 Assistant Chief Engineer, St. L. & S. F. R. R.
 Civil Engineering and Wood Preservation.
- Henn, Arthur Wilbur, Bloomington.
 Zoology.
- Hennel, Cora, Bloomington, Ind.
- Hennel, Edith A., Bloomington, Ind.
- Hetherington, John P., 418 Fourth St., Logansport.
 Physician.
 Medicine, Surgery, X-Ray, Electro-Therapeutics.
- Hinman, J. J., Jr., University of Iowa, Iowa City, Ia.
 Chemist, Dept. Public Health and Hygiene.
 Chemistry.
- Hole, Allen D., Richmond.
 Professor Earlham College.
 Geology.
- Hostetler W. F., South Bend.
 Geography and Indian History.
- Hubbard, Lucius M., South Bend.
 Lawyer.
- Hufford, Mason E., Bloomington.
 Physics.

- Hutchins, Chas. P., Buffalo, N. Y.
Athletics.
- Hutton, Joseph Gladden, Brookings, South Dakota.
Associate Professor of Agronomy, State College.
Agronomy, Geology.
- Hyde, Carl Clayton, Bloomington.
Geology.
- Hyde, Roscoe Raymond, Terre Haute.
Assistant Professor, Physiology and Zoology, Indiana State Normal.
Zoology, Physiology, Bacteriology.
- Hyslop, George, Bloomington.
Philosophy.
- Ibison, Harry M., Marion.
Instructor in Science, Marion High School.
- Iddings, Arthur, Hanover.
Geology.
- Imel, Herbert, South Bend.
Zoology.
- Inman, Ondess L., Bloomfield.
Botany.
- Irving, Thos. P., Notre Dame.
Physics.
- Jackson, D. E., St. Louis, Mo.
Assistant Professor, Pharmacology, Washington University.
- Jackson, Thomas F., Bloomington.
Geology.
- James, Glenn, West Lafayette.
Mathematics.
- Johnson, A. G., Madison, Wisconsin.
- Jones, Wm. J., Jr., Lafayette.
State Chemist, Professor of Agriculture and Chemistry, Purdue University.
Chemistry, and general subjects relating to agriculture.
- Jordan, Charles Bernard, West Lafayette.
Director School of Pharmacy, Purdue University.

- Koezmarek, Regedius M., Notre Dame.
Biology.
- Kenyon, Alfred Monroe, 315 University St., West Lafayette.
Professor of Mathematics, Purdue University.
Mathematics.
- Keubler, John Ralph, 110 E. Fourth St., Bloomington.
Chemistry.
- von KleinSmid, R. B., Tucson, Ariz.
- Koch, Edward, Bloomington.
Physiology.
President University of Ariz.
- Liebers, Paul J., 1104 Southeastern Ave., Indianapolis.
- Ludwig, C. A., 210 Waldron St., West Lafayette, Ind.
Assistant in Botany, Purdue University.
Botany, Agriculture.
- Ludy, L. V., 229 University St., Lafayette.
Professor, Experimental Engineering, Purdue University.
Experimental Engineering in Steam and Gas.
- Malott, Clyde A., Bloomington.
Physiology.
- Marshall, E. C., Bloomington.
Chemistry.
- Mason, Preston Walter, Lafayette.
Entomology.
- Mason, T. E., 226 S. Grant St., Lafayette.
Instructor Mathematics Purdue University.
Mathematics.
- McBride, Robert W., 1239 State Life Building, Indianapolis.
Lawyer.
- McCartney, Fred J., Bloomington.
Philosophy.
- McClellan, John H., Gary, Ind.
- McCulloch, T. S., Charlestown.
- McEwan, Mrs. Eula Davis, Bloomington, Ind.
- McGuire, Joseph, Notre Dame.
Chemistry.

- Mance, Grover C., Bloomington, Ind.
- Markle, M. S., Richmond.
- Middletown, A. R., West Lafayette.
Professor of Chemistry, Purdue University.
Chemistry.
- Miller, Daniel T., Indiana University, Bloomington.
Anatomy.
- Miller, Fred A., 3641 Kenwood Ave., Indianapolis.
Botanist for Eli Lilly Co.
Botany, Plant Breeding.
- Molby, Fred A., Bloomington.
Physics.
- Montgomery, Ethel, South Bend.
Physics.
- Montgomery, Hugh T., South Bend.
Physician.
Geology.
- Moon, V. H., Indianapolis.
Pathology.
- Moore, George T., St. Louis, Mo.
Director, Missouri Botanical Garden.
Botany.
- Morris, Barclay D., Spiceland Academy, Spiceland.
Science.
- Morrison, Edwin, 80 S. W. Seventh St., Richmond.
Professor of Physics, Earlham College.
Physics and Chemistry.
- Morrison, Harold, Indianapolis, Ind.
- Mowrer, Frank Karlsten, Interlaken, New York.
Cooperative work with Cornell University.
Biology, Plant Breeding.
- Muncie, F. W.
- Murray, Thomas J., West Lafayette.
Bacteriology.
- Myers, B. D., 321 N. Washington St., Bloomington.
Professor of Anatomy, Indiana University.

- Nelson, Ralph Emory, 419 Vine St., West Lafayette.
Chemistry.
- Nieuwland, J. A., The University, Notre Dame, Ind.
Professor, Botany, Editor Midland Naturalist.
Systematic Botany, Plant Histology, Organic Chemistry.
- North, Cecil C., Greencastle.
- Northnagel, Mildred, Gary, Ind.
- Oberholzer, H. C., U. S. Department Agriculture, Washington, D. C.
Biology.
- O'Neal, Claude E., Bloomington.
Graduate Student, Botany, Indiana University.
Botany.
- Orahood, Harold, Kingman.
Geology.
- Orton, Clayton R., State College, Pennsylvania.
Assistant Professor of Botany, Pennsylvania State College.
Phytopathology, Botany, Mycology, Bacteriology.
- Osner, G. A., Ithaca, New York.
Care Agricultural College.
- Owen, D. A., 200 South State St., Franklin.
Professor of Biology. (Retired.)
Biology.
- Owens, Charles E., Corvallis, Oregon.
Instructor in Botany, Oregon Agricultural College.
Botany.
- Payne, Dr. F., Bloomington, Ind.
- Peffer, Harvey Creighton, West Lafayette.
Chemical Engineering.
- Petry, Edward Jacob, 267 Wood St., West Lafayette.
Instructor in Agriculture.
Botany, Plant Breeding, Plant Pathology, Bio-Chemistry.
- Phillips, Cyrus G., Moores Hill.
- Pickett, Fermen L., Bloomington.
Botany Critic, Indiana University Training School.
Botany, Forestry, Agriculture.
- Pipal, F. J., 11 S. Salisbury St., West Lafayette.

Powell, Horace, West Terre Haute.

Zoology.

Price, James A., Fort Wayne.

Ramsey, Earl E., Bloomington.

Principal High School.

Ramsey, Glenn Blaine, Orono, Me.

Botany.

Rhinehart, D. A., Bloomington.

Anatomy.

Rice, Thurman Brooks, Winona Lake.

Botany.

Schultze, E. A., Laurel.

Fruit Grower.

Bacteriology, Fungi.

Schnell, Charles M., South Bend.

Earth Science.

Schierling, Roy H., Bloomington.

Shimer, Dr. Will, Indianapolis, Ind.

Director, State Laboratory of Hygiene.

Shockel, Barnard, Professor State Normal, Terre Haute, Ind.

Sigler, Richard, Terre Haute.

Physiology.

Silvey, Oscar W., 437 Vine St., West Lafayette.

Instructor in Physics.

Physics.

Smith, Chas. Piper, College Park, Md.

Associate Professor, Botany, Maryland Agricultural College.

Botany.

Smith, Essie Alma, R. F. D. 6, Bloomington.

Smith, E. R., Indianapolis.

Horticulturist.

Smith, William W., West Lafayette, Genetics.

Biology.

Snodgrass, Robert, Crawfordsville, Ind.

Southgate, Helen A., Michigan City.

Physiography and Botany.

- Spitzer, George, Lafayette.
Dairy Chemist, Purdue University.
Chemistry.
- Stech, Charles, Bloomington.
Geology.
- Steele, B. L., Pullman, Washington.
Associate Professor of Physics, State College, Washington.
- Steimley, Leonard, Bloomington.
Mathematics.
- Stickles, A. E., Indianapolis.
Chemistry.
- Stoltz, Charles, 530 N. Lafayette St., South Bend.
Physician.
- Stoddard, J. M.
- Stone, Ralph Bushnell, West Lafayette.
Mathematics.
- Stork, Harvey Elmer, Huntingburg.
Botany.
- Stuart, M. H., 3223 N. New Jersey St., Indianapolis.
Principal, Manual Training High School.
Physical and Biological Science.
- Sturmer, J. W., 119 E. Madison Ave., Collingswood, N. J.
Dean, Department of Pharmacy, Medico-Chirurgical College of Philadelphia.
Chemistry, Botany.
- Taylor, Joseph C., Logansport.
Wholesale merchant.
- Tetrault, Philip Armand, West Lafayette.
Biology.
- Thompson, Albert W., Owensville.
Merchant.
Geology.
- Thompson, Clem O., Salem.
Principal High School.
- Thornburn, A. D., Indianapolis.
Care Pitman-Moore Co.
Chemistry.

- Travelbee, Harry C., 304 Oak St., West Lafayette.
Botany.
- Troop, James, Lafayette.
Entomology.
- Trueblood, Iro C. (Miss), 205 Spring Ave., Greencastle.
Teacher of Botany, Zoology, High School.
Botany, Zoology, Physiography, Agriculture.
- Tucker, Forest Glen, Bloomington.
Geology.
- Tucker, W. M., 841 Third St., Chico, California.
Principal High School.
Geology.
- Turner, William P., Lafayette.
Professor of Practical Mechanics, Purdue University.
- Vallance, Chas. A., Indianapolis.
Instructor, Manual Training High School.
Chemistry.
- Van Dorau, Dr., Earlham College, Richmond.
Chemistry.
- Van Nuys, W. C., Newcastle.
- Voorhees, Herbert S., 2814 Hoagland Ave., Fort Wayne.
Instructor in Chemistry and Botany, Fort Wayne High School.
Chemistry and Botany.
- Wade, Frank Bertram, 1039 W. Twenty-seventh St., Indianapolis.
Head of Chemistry Department, Shortridge High School.
Chemistry, Physics, Geology and Mineralogy.
- Walters, Arthur L., Indianapolis.
- Warren, Don Cameron, Bloomington, Ind.
- Waterman, Luther D., Claypool Hotel, Indianapolis.
Physician.
- Webster, L. B., Terre Haute, Ind.
- Weatherwax, Paul, Bloomington, Ind.
- Wemmer, M. L., 102 Garfield Ave., Valparaiso.
Professor of Botany.
Botany and Human Physiology.

- Weir, Daniel T., Indianapolis.
Supervising Principal, care School office.
School Work.
- Weyant, James E., Indianapolis.
Teacher of Physics, Shortridge High School.
Physics.
- Wheeler, Virges, Montmorenci.
- Wiancko, Alfred T., Lafayette.
Chief in Soils and Crops, Purdue University.
Agronomy.
- Wicks, Frank Scott Corey, Indianapolis.
Sociology.
- Wiley, Ralph Benjamin, West Lafayette.
Hydraulic Engineering, Purdue University.
- Williams, Kenneth P., Bloomington.
Instructor in Mathematics, Indiana University.
Mathematics, Astronomy.
- Williamson, E. B. Bluffton.
Cashier, The Wells County Bank.
Dragonflies.
- Wilson, Charles E., Bloomington.
Graduate Student, Zoology, Indiana University.
Zoology.
- Wilson, Guy West, Assistant Professor Mycology and Plant Pathology, State
University, Iowa City, Ia.
- Wissler, W. A., Bloomington.
Chemistry.
- Wood, Harry W., 84 North Ritter Ave., Indianapolis.
Teacher, Manual Training High School.
- Woodburn, Wm. L., 902 Asbury Ave., Evanston, Ill.
Instructor in Botany, Northwestern University.
Botany and Bacteriology.
- Woodhams, John H., care Houghton Mifflin Co., Chicago, Ill.
Traveling Salesman.
Mathematics.

Wootery, Ruth, Bloomington, Ind.

Yocum, H. B., Crawfordsville.

Young, Gilbert A., 725 Highland Ave., Lafayette.

Head of Department of Mechanical Engineering, Purdue University.

Zehring, William Arthur, 303 Russell St., West Lafayette.

Assistant Professor of Mathematics, Purdue University.

Mathematics.

Zeleny, Charles, University of Illinois, Urbana, Ill.

Associate Professor of Zoology.

Zoology.

Zufall, C. J., Indianapolis, Ind.

Fellows	70
Members, Active	260
Members, Non-resident	29
	<hr/>
Total	359

MINUTES OF THE SPRING MEETING
OF THE
INDIANA ACADEMY OF SCIENCE.

SOUTH BEND, INDIANA, THURSDAY, MAY 28, 1914.

Public meeting of the Academy in the auditorium of the South Bend High School. The meeting was addressed by Dr. John M. Coulter, of the University of Chicago, upon the subject "Plant Breeding and How it Will Help Solve the Problem of Our Food Supply." Attendance: About 250.

The speaker was introduced by Severance Burrage, of the Eli Lilly Drug Company of Indianapolis, President of the Academy.

BUSINESS SESSION MAY 28, 1914.

At the close of the public meeting addressed by Dr. John M. Coulter the Academy was called into business session by President Burrage.

The following committees reported:

Resolutions:

Upon motion of John S. Wright, seconded by W. S. Blatchley, and passed, "That the Secretary be instructed to wire Dr. C. H. Eigenmann our regrets at his inability to attend this meeting."

In accordance with the motion the following telegram was sent:

"Dr. Eigenmann—The members of the Indiana Academy of Science in session at the spring meeting, South Bend, May 28th, wish to express their regrets that the condition of your health prevented your attendance, and join in wishing you speedy and complete recovery."

"HOWARD E. ENDERS, Secy."

Membership:

The Membership Committee, F. M. Andrews, chairman, read the names of forty-four applicants for membership. Upon motion duly seconded and passed, the secretary cast the ballots for membership of the persons whose names were proposed.

Letters were read in acknowledgment of sympathy from Chas. W. Fairbanks; and of regrets from Dr. Harvey W. Wiley.

With the consent of the Academy, Mr. J. S. Wright was authorized to extend a special invitation to Dr. Wiley to attend the next fall and spring meetings of the Academy.

The choice of a time for the fall meeting was requested.

It was moved (Blatchley) and seconded (Cogshall) that the meeting be set for the Thanksgiving period, as heretofore for several years. The motion was amended, and passed, "to leave the time for the meeting in the hands of the Program Committee, with power to act."

(In the discussion prior to the passage of the amendment, the sentiment was against the Thanksgiving period.)

Announcements:

President Burrage announced that Professor Edward Lee Green, botanist, of the Smithsonian Institution, will be present tomorrow and will accompany the party on its excursion.

Mr. Charles Stoltz, M. D., chairman of the local committee, announced the program for tomorrow (Friday, May 29, 1914). Automobiles for the whole party leave 7:00 a. m. from the front of the Hotel Oliver; return to the same place at 4:00 o'clock in the afternoon. Banquet, 6:30 p. m., at the Hotel Oliver.

Attention was called to the fact that J. B. Garner of Crawfordsville had not received the publications and literature of the Academy. He desires his name on the mailing list.

BUSINESS SESSION MAY 29, 1914.

After the noon luncheon on the field trip—at the Adventist College, north of Berrien Springs, Michigan—the meeting was called to order by the President, Severance Burrage, for the purpose of hearing the report of the Membership Committee.

F. M. Andrews, chairman, presented the names of six persons for membership.

Upon motion, duly seconded and passed, the secretary cast the ballot for their election to membership.

Announcements concerning the further movements of the party were made by local chairman, Dr. Stoltz.

BANQUET AND BUSINESS SESSION.

HOTEL OLIVER, May 29, 1914.

An ample banquet was spread in one of the private dining-rooms of the Hotel Oliver for forty-eight persons. President Severance Burrage, as master of ceremonies, expressed pleasure with the entertainment provided by the local committee. He made a strong plea for a large attendance and an increased membership for the fall meeting.

President Burrage then called upon the following persons for toasts and for expression of their sentiments.

Rev. John Cavanagh, of Notre Dame University, extended the greetings of the city and of Notre Dame to the members of the Academy.

Professor Dennis, of Earlham College: "How the Other Man Looks at It."

Professor Edward Lee Green, of the Smithsonian Institution, Washington, D. C., "The Unchangeableness of Nature, or the Stability of Science and the future of the Academy of Science."

Amos Butler, chairman of the State Board of Charities: "The Problem of Dealing with Mental Defectives."

Judge Hubbard, of South Bend: "Geology of the Regions About South Bend."

Dr. W. S. Blatchley, former State Geologist: "Geological Rambles."

Father Nieuwland, of Notre Dame University: Invitation to members of the Academy to visit him in his laboratories.

Professor Mottier of Indiana University: "Conservation of Our Young Members."

Professor Bodine, Wabash College: "A Personal Appreciation of the Day."

Dr. J. C. Arthur. "The Influence of the Academy of Science."

Eugene Manning, City Comptroller of South Bend, in the absence of the Mayor, extended the good-will of the city.

Dr. Montgomery of South Bend, "Welcome to our Homes, and to our Town."

Dr. Stoltz, South Bend, expressed his appreciation in the name of the local committee.

The *Business Session* was called to order by President Burrage.

The Membership Committee, F. M. Andrews, chairman, reported the

names of five applicants for membership. Upon motion, duly seconded, and passed, the Secretary cast the ballot for their election to membership.

Attention was called by the President to the fact that nearly sixty members were added to the Academy during the spring meeting.

Amos Butler read several letters relative to the passage of the congressional bills on the protection of migratory birds. Attention was also called to the treaty with Canada, aimed to protect migratory birds.

The following motion was passed: "That the President of the Academy of Science be appointed a committee of one to express to Senators Shively and Kern the earnest desire that they support all efforts to make the migratory bird law effective, and support the international treaty with Canada for bird protection in every way possible."

In accordance with this motion the following telegram was sent:

"MAY 29, 1914.

"Hon. Benj. F. Shively, U. S. Senate, Washington, D. C.:

"The Indiana Academy of Science in session here heartily endorses your support of the appropriation to make the bird law effective, and relies upon you to support legislation for the protection of the birds, including the treaty with Canada. We regard these as important to all the people.

"SEVERANCE BURRAGE, President."

Mr. Butler requested all the members of the Academy to write Senators Kern and Shively relative to the bird legislation. Many members pledged to do so.

A resolution, introduced by Professor Bodine, was passed: "That a rising vote of thanks be extended to Dr. Stoltz, Dr. Montgomery, and the other members of the local committee for the fine entertainment today."

Adjournment.

HOWARD E. ENDERS,

Assistant Secretary.

The following were elected to membership May 29, 1914:

Ethel Montgomery, South Bend, Ind. Physics.

Helen A. Southgate, Michigan City, Ind. Physiography and Botany.

Jose Angel Caparo, Notre Dame, Ind. Physics and Mathematics.

Dr. D. A. Rhinehart, 301 N. Walnut Street, Bloomington, Ind. Anatomy.

Thurman Brooks Rice, Winona Lake, Ind. Botany.

- Charles M. Schnell, South Bend, Ind. Earth Science.
- Roy H. Schierling, P. O. Box 172, Bloomington, Ind.
- Philip Armand Tetrault, West Lafayette, Ind. Biology.
- James Troop, Lafayette, Ind. Entomology.
- Charles Stech, Bloomington, Ind. Geology.
- Leonard L. Steimley, Bloomington, Ind., Indiana Club. Mathematics.
- Forest Glenn Tucker, 430 E. Fourth Street, Bloomington, Ind. Geology.
- W. A. Wissler, 415 S. Grant Street, Bloomington, Ind. Chemistry.
- Glenn Blaine Ramsey, University of Maine, Orono, Maine. Botany.
- H. C. Oberholzer, U. S. Department of Agriculture, Washington, D. C. Biology.
- Thomas J. Murray, West Lafayette, Ind. Bacteriology.
- Fred A. Molby, 525 S. Park Avenue, Bloomington, Ind. Physics.
- Preston Walter Mason, Purdue University, Lafayette, Ind. Entomology.
- E. C. Marshall, 409 E. Fourth Street, Bloomington, Ind. Chemistry.
- Clyde A. Malott, 209 S. Dunn Street, Bloomington, Ind. Geology.
- Joseph McGuire, Notre Dame, Ind. Chemistry.
- Fred J. McCartney, Bloomington, Ind. Philosophy.
- Edward Koch, 314 N. Washington Street, Bloomington, Ind. Physiology.
- Charles Bernard Jordan, West Lafayette, Ind. Director of School of Pharmacy, Purdue University.
- Regedius M. Kaezmarek, Notre Dame, Ind. Biology.
- Glenn James, West Lafayette, Ind. Mathematics.
- Thomas F. Jackson, 325 S. Grant Street, Bloomington, Ind. Geology.
- Thos. P. Irving, Notre Dame, Ind. Physics.
- Herbert Imel, South Bend, Ind., care Studebaker School. Zoölogy.
- George Hyslop, Bloomington, Ind., Kappa Sigma House. Philosophy.
- Chas. P. Hutchins, Buffalo, New York. Athletics.
- W. F. Hostetler, South Bend, Ind. Geography and Indian History.
- Paul M. Harman, 111 W. Dunn Street, Bloomington, Ind. Geology.
- Louis M. Hammerschmidt, South Bend, Ind. Science of Law.
- William Happ, South Bend, Ind. Botany.
- Elbert Howard Clark, West Lafayette, Ind. Mathematics.
- John B. Berteling, South Bend, Ind. Medicine.
- Homer Glenn Fisher, 727 Atwater Street, Bloomington, Ind. Zoölogy.
- David Christie Duncan, West Lafayette, Ind. Instructor in Physics Purdue University.

- Albert H. Dewey, West Lafayette, Ind. Dept. of Pharmacy, Purdue University.
- Ernest A. Davis, Notre Dame, Ind. Chemistry.
- Noble H. Coryell, 330 S. Henderson Street, Bloomington, Ind. Chemistry.
- Jacob Roland Collins, W. Lafayette, Ind. Instructor in Physics, Purdue University.
- Flora Charlotte Anderson, Wellesley College, Wellesley, Mass. Botany.
- Paul J. Carlyle, 315 N. Washington Street, Bloomington, Ind. Chemistry.
- William Franklin Baker, Indianapolis, Ind., care Eli Lilly & Co. Medicine.
- George A. Baker, South Bend, Ind. Archaeology.
- J. A. Badertscher, 509 N. Washington Street, Bloomington, Ind. Anatomy.
- William Gidley, W. Lafayette, Ind. Dept. of Pharmacy, Purdue University.
- Willis D. Gatch, Indianapolis, Ind., Indiana University Medical School. Anatomy.
- George Kedzie Foresman, 110 S. Ninth Street, Lafayette, Ind. Chemistry. Purdue University.
- Carl Clayton Hyde, Bloomington, Ind., Kappa Sigma House. Geology.
- Arthur Wilbur Henn, 821 Atwater Avenue, Bloomington, Ind. Zoölogy.
- Ralph Benjamin Wiley, 1012 Seventh Street, West Lafayette, Ind., Purdue University. Hydraulic Engineering.

MINUTES OF THE THIRTIETH ANNUAL MEETING.
INDIANA ACADEMY OF SCIENCE.

CLAYPOOL HOTEL, INDIANAPOLIS, IND., DEC. 4, 1914.

The Executive Committee of the Indiana Academy of Science met in the Assembly Room, and was called to order by the President, Severance Burrage of Indianapolis, Ind.

The following members were present: Severance Burrage, Andrew J. Bigney, Howard E. Enders, W. A. Cogshall, Donaldson Bodine, J. P. Naylor, Glenn Culbertson, John S. Wright, Stanley Coulter, A. W. Butler, and Robert Hessler.

The minutes of the Executive Committee of 1913 were read and approved.

The reports of the standing committees were then taken up.

The Program Committee, John S. Wright, chairman, reported the work completed as indicated by the printed program, with three additional papers.

Amos W. Butler, member of the State Library Committee, reported that progress is being made in the way of housing books of the Indiana Academy of Science. Lack of available funds hampers the State Librarian in binding of exchanges. The State Librarian requests a ruling upon the matter of sending out proceedings to individuals upon request.

On motion, duly passed, the matter of distribution of proceedings is left to the discretion of the Committee on Distribution of Proceedings and the State Librarian.

On motion, duly passed, John S. Wright is authorized to prepare a revised membership blank and have it in order for use next year.

REPORT OF MEMBERSHIP COMMITTEE, F. M. ANDREWS, CHAIRMAN.

The following named persons are proposed for membership in the Academy:

Harry C. Travelbee, 304 Oak Street, West Lafayette, Ind. Botany.

- William J. Cotton, 5363 University Avenue, Indianapolis, Ind. Physics and Chemistry.
- Charles H. Arndt, Lafayette, Ind. Biology.
- William W. Smith, West Lafayette, Ind. Genetics.
- Oliver P. Ferry, West Lafayette, Ind. Physiology.
- Louis Frederick Heimlich, 703 North Street, Lafayette, Ind. Biology.
- Harry Creighton Pepper, West Lafayette, Ind. Chemical Engineering.
- Paul E. Bowers, Michigan City, Ind. Medicine.
- Charles P. Emerson, Hume-Mansur Building, Indianapolis, Ind. Dean Indiana University Medical College. Medicine.
- Ralph Emory Nelson, 419 Vine Street, West Lafayette, Ind. Chemistry.
- Ralph Bushnell Stone, West Lafayette, Ind., Purdue. Mathematics.
- W. H. Hanna, 828 Atwater Avenue, Bloomington, Ind. Mathematics.
- Fred Earl Robbins, 215 Waldron Street, West Lafayette, Ind. Agriculture.
- Ralph B. Stone, 307 Russell Street, West Lafayette, Ind. Mathematics.
- Frank Scott Corey Wicks, Indianapolis, Ind. Sociology.
- F. C. Atkinson, Indianapolis, Ind., American Hominy Co. Chemistry.
- Arthur Iddings, Hanover, Ind. Geology.
- Harry Binford, Earlham, Ind. Zoölogy.
- Samuel Hugh Hamill, 110 E. Fourth Street, Bloomington, Ind. Chemistry.
- Ralph Howard Carr, Lafayette, Ind. Chemistry.
- George Lindenburg Clark, Greencastle, Ind., DePauw University. Chemistry.
- Dr. D. W. Davis, Greencastle, Ind., DePauw University. Biology.
- Ondess L. Inman, Bloomfield, Ind. Botany.
- Richard Sigler, Terre Haute, Ind. Physiology.
- Adam L. Bowles, Terre Haute, Ind. Zoölogy.
- John Ralph Keubler, 110 E. Fourth Street, Bloomington, Ind. Chemistry.
- Dr. Daniel T. Miller, Indiana University, Bloomington, Ind. Anatomy.
- Fred Donaghy, Ossian, Ind. Botany.
- Horace Powell, West Terre Haute, Ind. Zoölogy.
- William Marion Goldsmith, Oakland City, Ind. Zoölogy.
- Muri E. Fulk, Decatur, Ind. Anatomy.
- Michael James Blew, R. R. 1, Wabash, Ind. Chemistry and Botany.
- Harvey Elmer Stork, Huntingburg, Ind. Botany.
- Harold Orabood, Kingman, Ind. Geology.
- Dr. VanDoran, Earlham College, Richmond, Ind. Prof. of Chemistry.

- Carlton Edwards, Earlham College, Earlham, Ind.
 W. C. Van Nuys, Newcastle, Ind. Medicine.
 Barclay D. Morris, Spiceland, Ind., Spiceland Academy. Science.
 Paul H. Brown, Richmond, Ind. Physics and Manual Training.
 Bernard F. Dostal, Bloomington, Ind. Physics.
 O. C. Berry, Waldron Street, West Lafayette, Ind. Engineering.
 James M. Breckenridge, 514 S. Walnut Street, Crawfordsville, Ind. Chemistry.
 Dr. V. H. Moon, Indianapolis, Ind. Pathology.
 H. C. Balcom, 1023 Park Avenue, Indianapolis, Ind. Botany.
 Elton R. Clark, Indianapolis, Ind. Zoölogy.
 A. E. Stickels, 768 Massachusetts Avenue, Indianapolis, Ind. Chemistry.
 John C. Dean, Indianapolis, Ind. Astronomy.
 Dr. G. S. Bliss, Fort Wayne, Ind., State School for Feeble-Minded. Medicine.
 On motion they were recommended.

The Treasurer, W. A. Cogshall, reported as follows:

Balance from 1913	\$254 78
Collected to December 1st	133 00
	<hr/>
Total	\$387 78
Expenses December 1st	174 79
	<hr/>
	\$212 99

Upon motion, duly passed, the report was received and turned over to the Auditing Committee.

On publication of Proceedings, H. E. Barnard, editor, reported the publication and distribution of Proceedings for 1913.

On motion the following members were recommended as Fellows:

Fellows, Wm. L. Bryan, Indiana University, Psychology; E. B. Williamson, Bluffton, Biology; A. W. Kenyon, Purdue, Mathematics; J. A. Nieuwland, Botany; Wm. M. Blanchard, DePauw University, Chemistry.

The matter of avoiding duplication of reports was discussed. It is deemed desirable to have committees report at the executive session and that only the Secretary's summarized report be read at the general meetings.

The Secretary reported the binding of the minutes of previous sessions of the Academy, and the purchase of a new Secretary's book. The Secretary reported that the State Librarian has consented to deposit the bound copies of the minutes in the safe of the State Library.

President Burrage reported that he had appointed the following persons as delegates to the State Tax Association: Judge R. W. McBride, William Lowe Bryan. Alternates, E. B. Williamson, Dr. Cecil C. North. Adjournment.

GENERAL SESSION. ASSEMBLY ROOM, 2:00 P. M.

The meeting was called to order by President Severance Burrage.

The minutes of the Executive Committee were read and approved.

On motion, duly passed, the persons who were recommended for membership were elected members of the Indiana Academy of Science.

On motion, duly passed, the five persons who were recommended by the Executive Committee were elected fellows in the Academy of Science.

The regular program was then taken up. The papers numbered 1-11 were read in general session; after which the Academy separated into two sections as follows: Section A—Bacteriology, Botany and Zoology, presided over by President Burrage and H. E. Enders being Secretary. Section B—Chemistry, Physics, Engineering, Geography, Geology, Mathematics, and Meteorology; W. A. Cogshall presiding and A. J. Bigney being Secretary.

EVENING SESSION, 8:00 O'CLOCK.

Business:

The report of the Nominating Committee was as follows: President—W. A. Gogshall, Bloomington; Vice-President—W. A. McBeth, Terre Haute; Secretary—A. J. Bigney, Moores Hill; Assistant Secretary—Howard E. Enders, Lafayette; Press Secretary—Frank B. Wade, Indianapolis; Treasurer—William M. Blanchard, Greencastle; Editor—H. E. Barnard, Indianapolis.

On motion the report was accepted, and the persons named were elected for the ensuing year.

SATURDAY, DECEMBER 5TH, 9:00 A. M.

Business:

The following resolutions were presented by the chairman of the Resolution Committee, Stanley Coulter of Purdue University:

Resolution No. 1.

Resolved, That the Indiana Academy of Science, recognizing the importance to the State and Society of the Preservation of the Public Health, hereby endorses the proposal for a thoroughly trained county health commissioner, who will give his entire time to this service, and urges the Legislature to pass a law to that end.

Resolution No. 2.

Resolved, That we extend a vote of thanks to Mr. Lawrence, manager of the Claypool Hotel, for the use of rooms for the meetings of the Academy, and for his continued kindness in aiding us in every way possible.

Resolution No. 3.

(On suggestion of R. R. Ramsey.)

Be it Resolved, That as certain periodicals, such as Science Abstracts A and B, *Baiblaita zu der Annalen der Physik* (in Physics), and in all probability other magazines in other lines are not on the list of exchanges, that such exchanges be sought and obtained if possible. If not successful, periodicals whose nature is primarily abstracting, should be put on the mailing list.

On suggestion of Will Scott that our journals should be open to foreign scientists, Stanley Coulter moved, and it was passed, that a committee be appointed to take up the matter with the Smithsonian Institution regarding the opening of our scientific journals to foreign scientists for publication of their papers.

On motion the Academy endorsed the plan of the State Geologist to cooperate with the United States Geological Survey, with a view to the prevention of floods in Indiana (see resolution of last year), and ordered that a copy of that resolution be sent to the Governor, Lieutenant-Governor, and Speaker of the House.

Adjourned to sections.

AFTERNOON SESSION, 1:00 O'CLOCK.

Section B met to complete the reading of its papers.

Adjournment.

A. J. BIGNEY, Secretary.

SEVERANCE BURRAGE, President.

PROGRAM
OF THE THIRTIETH ANNUAL MEETING OF THE
INDIANA ACADEMY OF SCIENCE,
CLAYPOOL HOTEL, INDIANAPOLIS,
FRIDAY AND SATURDAY, DECEMBER 4 AND 5, 1914.

GENERAL PROGRAM.

FRIDAY, DECEMBER 4.

Meeting of Executive Committee	10:30 a. m.
General Session	2:00 p. m.
Section Meetings.....	4:00 p. m.
Informal Dinner, Tickets \$1.00. For reservation, apply to W. M.	
Blanchard, DePauw University, Greencastle.....	6:00 p. m.
Business Session.....	7:45 p. m.
General Session.....	8:00 p. m.

Address by the Retiring President, Severance Burrage.

Symposium on Feeble-mindedness.

SATURDAY, DECEMBER 5.

General Session.....	9:00 a. m.
Section Meetings.....	9:45 a. m.
General Session, followed by Section Meetings (provided the time is required to complete the program).....	2:00 p. m.

LIST OF PAPERS TO BE READ.

Address by the Retiring President, Mr. Severance Burrage.

GENERAL SESSION AT EIGHT O'CLOCK, FRIDAY EVENING.

Subject: "Science in Its Relation to Conservation of Human Life."

Symposium: Some Scientific and Practical Aspects of the Problem of
Feeble-mindedness.

1. The Feeble-minded Family..... Amos W. Butler, Indianapolis
2. The Problem of Feeble-mindedness..... Dr. G. S. Bliss, Ft. Wayne
3. The Feeble-minded and Delinquent Boy. Dr. F. E. Paschal, Jeffersonville
4. The Feeble-minded and Delinquent Girl. Dr. E. E. Jones, Evanston, Ill.
5. Feeble-mindedness in the Public School. Miss Katrina Myers, Indianapolis

Discussion opened by Mr. F. S. C. Wicks.

GENERAL—TWO O'CLOCK FRIDAY.

6. The Alcohol Problem in the Light of Coniosis,
20 minutes.....Robert Hessler
7. Cold Storage, Practical Conservation, 20 minutes.....H. E. Barnard
8. Changing Conditions among the Cumberland Plateau Mountain
People, lantern, 20 minutes.....Bernard H. Schockel
9. The Conservation of Energy, 30 minutes.....Arthur L. Foley
10. Agriculture in Southern Indiana, 15 minutes.....C. G. Phillips
11. The Chief Reason for the Migration of Our Birds, 15 minutes.....D. W. Dennis

BACTERIOLOGY.

12. An Aeration Apparatus for Culture Solutions, with charts,
10 minutes.....Paul Weatherwax
13. Autagonism of *B. fluorescens* and *B. typhosus* in Culture,
10 minutes.....P. A. Tetrault

BOTANY.

14. Notes on the Distribution of the Forest Trees of Indiana,
15 minutes.....Stanley Coulter
15. A New Enemy of the Black Locust, 5 minutes.....Glenn Culbertson
16. The Parasitic Fungi Attacking Forest Trees in Indiana,
10 minutes.....Geo. N. Hoffer
17. A New Disease of *Viola cucullata*, lantern, 5 minutes.....H. W. Andersons
18. Oatsmut in Indiana, 15 minutes.....F. J. Pipal
- 18a. Weed Seeds in soil, 10 minutes.....F. J. Pipal
19. Additions to Indiana Flora, 3 minutes.....Chas. C. Deam
20. Some Peculiarities in *Spirogyra dubia*, 5 minutes.....Paul Weatherwax
21. Stomata of *Trillium nivale*, 10 minutes.....F. M. Andrews
22. Final Report on Cross Pollination of Corn, 3 minutes.....M. L. Fisher
23. The Primrose-Leaved Violet in White County, charts
and specimens, 10 minutes.....Louis F. Heimlich
24. Continuous Rust Propagation without Sexual Reproduction,
10 minutes.....C. A. Ludwig
25. Correlation of Certain Long-cycled and Short-cycled Rusts,
10 minutes.....H. C. Travelbee
26. Some Species of *Nummularia* Common in Indiana, 10 minutes
.....C. E. O'Neal.
27. The Genus *Rosellinia* in Indiana, 2 minutes.....Glenn B. Ramsey

28. Cultivating and Breeding Medicinal Plants, lantern,
20 minutes.....Fred A. Miller
29. Some Large Botanical Problems, 10 minutes.....J. C. Arthur

ZOOLOGY.

30. The Alba Gehre Collection of Birds' Eggs in the Museum
of Purdue University, 10 minutes.....Howard E. Enders
31. A Study of the Maturation Period in the Mole-Cricket, blackboard,
10 minutes.....F. Payne
32. Note on a Peculiar Nesting Site of Chimney Swift,
3 minutes.....Glenn Culbertson
33. Mosaics in *Drosophila Ampelophila*, chart, 5 minutes..Horace M. Powell
34. New Mutations in the Genus *Drosophila* and their Behavior
in Heredity, chart, 10 minutes.....Roscoe R. Hyde
35. Notes on Indiana Earthworms, 10 minutes.....H. V. Heimburger
36. Insects of the Between-Tides Zone. 30 minutes.....Chas. H. Arndt
37. Regeneration in *Sagartia*, 5 minutes.....D. W. Davis
38. The Relation of Birds to Aquatic Life as Exemplified by
Observation and Studies made at Lake Maxinkuckee,
15 minutes.....Barton W. Evermann
39. The Reptiles and Batrachians of the Lake Maxinkuckee
Region, 20 minutes.....Barton W. Evermann.
40. A Physical and Biological Survey of Lake Maxinkuckee,
20 minutes.....Parton W. Evermann

CHEMISTRY.

41. The Quantitative Determination of Copper, 5 minutes..W. M. Blanchard
42. The Alundum Crucible as a Substitute for the Gooch
Crucible, 5 minutes.....George L. Clark
43. Some Recent Work in Dairy Chemistry, 20 minutes.....George Spitzer
44. Analysis of Zirconium Minerals, 10 minutes.....James Brown
45. Correlation of High School and College Chemistry,
20 minutes.....James Brown
46. Chemical Composition of Virgin and Cropped Indiana Soils,
10 minutes.....S. D. Conner

ENGINEERING.

47. Sewage Disposal, lantern. 15 minutes.....Charles Brossmann
48. Extension of Empirical Curve by the Addition of Estimated Values
to a Series of Observations, charts, 20 minutes.....Albert Smith

49. Tar-Forming Temperatures of American Coals, charts, 20 minutes
 O. C. Berry.

GEOGRAPHY.

50. Shawnee Mound, Tippecanoe County, A Glacial Alluvial Cone,
 charts and photographs, 10 minutes..... William A. McBeth

GEOLOGY.

51. Stratigraphic Correlation of the Outcrop at Spades, Indiana,
 15 minutes..... H. N. Coryell
52. Pennsylvania Fossil Plants of the Bloomington Quadrangle,
 5 minutes..... J. F. Jackson
53. Preliminary Geological History of Dearborn County, 10 minutes
 A. J. Bigney
54. Notes on the Cause of Asterism in "Starolite" (Asteriated Quartz),
 charts and specimens, 10 minutes..... Frank B. Wade
55. The Mississippian Section of Monroe County, charts, 15 minutes
 J. W. Beede
56. The Flatwoods Region of Owen and Monroe Counties, Indiana,
 25 minutes..... Clyde A. Malott

MATHEMATICS.

57. Mechanical Device for Testing Merseane Numbers for Primes,
 5 minutes..... Thomas E. Mason
58. Some Properties of Binominal Coefficients, 20 minutes..... A. M. Kenyon

METEOROLOGY.

59. The Watertown, S. D., Tornado of June 23, 1914, 10 minutes
 J. Gladden Hutton

PHYSICS.

60. A New Lantern and Projector, lantern, 10 minutes..... Arthur L. Foley
61. Some Text Book Inconsistencies, 5 minutes..... Arthur L. Foley
62. The Mechanism of Light and Heat Radiations, 10 minutes
 James E. Weyant
63. A Simple Form for the Carey Foster Bridge, lantern, 5 minutes
 J. P. Naylor
64. The Change of the Radio Activity of Certain Springs, lantern,
 10 minutes..... R. R. Ramsey
65. Radio Activity of Spring Water, lantern, 10 minutes..... R. R. Ramsey
66. A Radio-Active Electroscope, lantern, 5 minutes..... Edwin Morrison

SCIENCE IN ITS RELATION TO THE CONSERVATION OF HUMAN LIFE.

SEVERANCE BURRAGE.

Nearly every branch of science has direct or indirect relation toward the conservation of human life. Unfortunately the appreciation of scientific work has been from the industrial rather than from the humanitarian viewpoint, and such researches as have resulted in discoveries that have commercial importance have been the ones to receive the plaudits of the public.

Chemistry, physics, geology and biology in all of their subdivisions have undoubtedly contributed in the work of saving human life. For example, we have in *chemistry* the studies of the impurities in the air, water, food and drugs, practical applications on the purification of water and sewage, and so on. In *physics* and its various branches we have the practical application of safety devices of all kinds, rescue apparatus for mines, the developments in rapid communication, climaxed by the invention of the wireless telegraph and telephone, most useful in the prevention of accidents, the various inventions which protect employes from dangerous machinery, the development of fire-fighting apparatus, and special protective devices against floods, earthquakes, cyclones and other disasters. In *geology*, the selection of proper building stone, the dangers from the corrosion of building stone in different climates, the selection of proper building sites, and, indirectly the discoveries of coal, oil, and other things essential in many phases of human existence. In *biology*, particularly in the subdivisions, bacteriology, medicine and sanitary science, we find some of the most important discoveries resulting in the prevention of disease and death. Even in entomology—the life histories of various mosquitoes and flies have important bearing on the prevention of disease. In bacteriology the discovery of the causes of transmissible diseases, through the research of Pasteur, Koch and others, methods in the rapid diagnosis of disease, protective inoculation against disease, resulting from the work of Jenner Von Behring and others, are familiar to you all. In medicine the application of asepsis and cleanliness to surgical methods has revolution-

ized this important branch of medicine and made it a most important factor in the conservation of human life; the discovery of selective chemical substances for the treatment of specific diseases, such as quinine for malaria, salvarsan for syphilis and most recently ipecac for amebic dysentery and pyorrhea.

In sanitary science the development has been most remarkable. This includes its practical applications in sewage and garbage disposal, street cleaning and the sanitary construction of pavements, the sanitation of heating and ventilation of factories, workshops and schools, the medical inspection of schools, the sanitation of railway cars, and stations, and such special sanitary devices as individual drinking cups, dental lavatories in railroad cars, and the various applications of sanitation to the farmhouse and rural dwelling.

The above outline, which is obviously incomplete, suggests some of the things which science in its various branches has done that have been and can be applied in the conservation of human life. Granting that science has done all of these things and many more, I would raise this important question: Is the public at large getting the full benefit of all of this scientific work? Is the public taking advantage of these discoveries of science? In my opinion it is not.

Notwithstanding intensive efforts on the part of state boards of health, extension departments in our universities, instruction given before farmers' institutes, educational activities of anti-tuberculosis societies and insurance companies, we find the death rate from preventable diseases decreasing very little if at all. In some communities the deaths from preventable diseases are on the increase. In our own State we find very little change in the last ten years in the deaths from preventable diseases.

This Academy is of course particularly interested in Indiana. Can not this Academy suggest or recommend ways and means to apply throughout this State the various developments of science relating to health and disease prevention in such a way as to create a healthier and longer-lived citizenship? A commission appointed recently in Massachusetts to investigate the high cost of living stated—

“The increased vital efficiency of the citizens of this State (Massachusetts) which would result from a conservation of the present waste of health would, if expended in labor, increase the earnings of those whose health is impaired and also lessen the burdens of


those who are at present unnecessarily ill. This increase in earnings would thus tend to reduce the cost of living, increase the total earnings of the citizens and make the average income larger."

Thus conservation of health means higher wages, which enables the worker to keep ahead of the increasing price of the commodities of life. Surely this is worth striving for, and in my opinion Indiana can decrease the death rate and lengthen life and thus bring about this condition. To accomplish it I would urge the adoption of a law which would provide for a full-time health commissioner in every county in the State. This commissioner must be especially trained in sanitary science and the various applications of the other sciences in so far as they affect the prevention of unnecessary deaths. If such a law were passed, and backed up by an intelligent public, we would have the healthiest State in the Union in a very short time. I state my belief because where such sanitary applications have been thoroughly carried out, we have as a result healthful conditions. I would cite as an instance of this the Panama Canal district. While the names of Goethals and Gorgas will go down to posterity as constructors of the Panama Canal, they should receive more credit for the sanitary organization and administration which made the construction of the canal possible. They converted one of the most unhealthful localities in the world, where the death rate was over 70 per 1,000, to the most healthful spot with the death rate of less than 6 per 1,000, a death rate lower than that of any other civilized community in the world.

Another instance of the practical applications of sanitary science has been in our military camps. Reports show that out of 12,000 men in these camps there was not a single case of smallpox or typhoid fever for a period covering two years. Typhoid fever has long been one of the pests of camp life, but through improved sanitation and typhoid vaccination, this disease has been absolutely eliminated. Other diseases in these camps over which there was not such perfect control showed a great reduction.

With such fine examples of successful sanitary administration it seems to me justifiable to make application to our own communities, with of course necessary modifications. I therefore would suggest that this Academy at this session pass resolutions favoring the passage of a law at the next Legislature which would provide for a competent, full-time health commissioner in each county in this State. I know of no way in which the Indiana Academy of Science can better further the best interests of the State with reference to the conservation of human life.






THE 'C' FAMILY

FIVE GENERATIONS	MALE	FEMALE	SEX UNK.	TOTAL
INDIVIDUALS RECORDED	24	28	5	57
MENTAL CONDITION				
PEOPLE MINDED	17	19	..	36
INSANE	7	1	..	8
NORMAL	10	6	..	16
NORMALITY IN QUESTION	4	2	5	11
SEX OFFENDERS	3	4	..	7
ILLEGITIMATE	1	3	5	9
INTENTION INMATES	6	12	..	18

THE 'C' FAMILY



THE 'C' FAMILY

	MALE	FEMALE	SEX UNKNOWN	TOTAL
FIVE GENERATIONS				
INDIVIDUALS RECORDED	24	28	5	57
MENTAL CONDITION:				
PEOPLE MIMED	17	19		36
INSANE	1	1		2
NORMAL	6	6		12
NORMALITY IN QUESTION	4	2	5	11
SEX OFFENDERS	3	4		7
ILLEGITIMATE	1	3		4
INSTITUTION INMATES	6	12		18

A FAMILY IN ONE NEIGHBORHOOD.*

AMOS W. BUTLER,
Secretary Indiana Board of State Charities.
Indianapolis.

Not long ago I had handed to me a little card containing these words:

“When shall we apply the same intelligence to breeding human beings that we apply to breeding cattle?”

It came at a time when we were investigating the family histories of some of the state's wards whose names are recorded in the registration of the Board of State Charities. The application was made more striking by having before me charts of some of these families, some of them running back five or six generations. These tell a story of degeneracy that is appalling. I have thought you would be interested in one of them whose visible beginning was in a pair of feeble-minded ancestors about a hundred years ago. It includes five generations, represented by fifty-seven individuals.

There would be some changes in the chart as a result of subsequent investigations but in the main the facts are as given.

<i>The “C” Family.</i>		Sex		
	Male.	Female.	Unknown.	Total.
Individuals recorded	24	28	5	57
Mental condition:				
Feeble-minded	17	19	.	36
Insane	1	.	1
Normal	3	6	.	9
Normality in question.....	4	2	5	11
Sex offenders	3	4	.	7
Illegitimate	1	3	5	9

*Read at Indiana Academy of Science, Indianapolis 1914.

Our investigations brought to light the fact that eighteen members of this family have been at some time of their lives inmates of public institutions in Indiana. Our information as to the length of time five of these were on public support is incomplete; it was before our present registration of institution inmates was begun. Concerning the other thirteen, however, we have accurate data. To date they have spent a total of 203 years, 5 months in public institutions.

Eleven have been in county poor asylums 44 years, 3 months; 7 have been in orphans' homes 40 years, 7 months; 8 have been in the School for Feeble-Minded Youth 110 years, 11 months; one has been in the Indiana Girls' School 7 years, 8 months. If their maintenance has averaged but \$125 a year, the total cost has already amounted to more than \$25,000. This is not all. There are now five young women of this family in the School for Feeble-Minded Youth at Fort Wayne. Their ages are 22, 23, 24, 28 and 32 years, an average of 25.8 years. At the annual average per capita cost of maintenance in that institution last year (\$140.68) these five young women cost the State \$700.00 a year.

I ask this question: What are we going to do about it? It is one of our problems.

A legislative commission in New Jersey has recently made a report after a thorough investigation of the problem of mental defectiveness. Other commissions in New York, Massachusetts, and elsewhere have been, or are, engaged in this work. Is it not time that Indiana should wake up and have a commission to study the problem of mental defectiveness? What is the condition? What are the needs? How shall they be met? What is the wisest plan to follow? What are we going to do to save ourselves this continually increasing population of mental defectiveness, that is shown by this chart and could be shown by a hundred others, to be growing up in the State of Indiana without our knowledge, without our thought, without our effort to prevent it? The question comes to us: What will we do about it?

THE PROBLEM OF FEEBLE-MINDEDNESS.

G. S. BLISS.

The first recorded attempt to do something for feeble-mindedness occurred in the year 1800, when Dr. Itard, a French physician, tried to educate a so-called "wild boy" found in the woods. The attempt failed because the boy was feeble-minded, and was followed in France by several abortive attempts to educate feeble-minded persons.

The first successful attempt in this direction was made by the School for the Deaf and Dumb at Hartford, Conn., in 1836. They took several feeble-minded children and succeeded in training them a little in school work and in forming better habits of life.

In 1846 Dr. Seguin, a pupil of Dr. Itard, opened a successful school for mental defectives in France. This attempt succeeded so well that other schools were soon founded for this most unfortunate class. In 1848 Massachusetts started the first state school in the United States. This was followed by other States, and in 1879 Indiana established her present school.

All these schools were started with the idea that mental defect was curable, and that the idiot or imbecile could be educated to become a self-supporting and dependable citizen. This we now know to be an impossibility, and the fact is coming to be more generally recognized that there is no cure for mental defect. It is a condition, not a disease.

Insanity is a disease attacking a developed brain and is often cured; feeble-mindedness is never cured, but may be greatly relieved by proper training and care.

There are between 5,000 and 6,000 feeble-minded persons in Indiana needing institutional care today, and only about one-fourth of these are receiving it. These people are at large, reproducing defectives in an ever-increasing amount, like the waves from a pebble thrown into a lake. If we are to protect the coming generations of our sons and daughters, grandsons and granddaughters from this growing burden, we must wake up to the condition and *do* something about it.

What to do, and how to do it, constitutes one of the most serious problems for our State.

I presume if I were to ask any member of this body before me tonight, what in his estimation was the best measure, he would say sterilization or asexualization. And to that measure I would give a hearty amen, if it were as practical a proceeding as I wish it were. Theoretically it is good, but in practice, owing to ignorance, false sentimentality, honest disbelief in the measure by honest people, it is not as easily carried out as most of the persons here would think. If applied, it should be used on persons at large, not on those already segregated and cared for by public institutions; but the problem here is first to catch them, and then to decide where to draw the line.

I believe that for many years to come, segregation will be our best method of dealing with this problem. Mr. Butler has recommended for this State a real practical step in the establishment by this Legislature of a commission to investigate this condition throughout the State and report to our next Legislature.

I have also recommended a large farm, 2,000 acres or more, somewhere in the south central part of the State where the adult boys and men can live as useful and happy as may be; and also another smaller farm where the older women could care for chickens, turkeys, and small fruits, living their lives apart from the world, where they are such complete failures.

I believe that better marriage laws, permitting no one to marry without a clean bill of health, would be a help.

Whenever alcohol and vice are abolished in this world the feeble-mindedness from those causes will cease, and the public registration of venereal disease would prove a potent weapon against mental defect. I believe, that every case of syphilis and gonorrhoea should be registered with the health officer as well as smallpox or typhoid fever.

I hope Indiana will realize in the near future the momentousness of this problem, and by meeting it and better preventing the reproduction of defect, place herself where she belongs, at the pinnacle of those States who prevent, as well as provide for this burden of feeble-mindedness on her community.

THE FEEBLE-MINDED AND DELINQUENT BOY.

FRANKLIN C. PASCHAL.

In an examination of the relation of feeble-mindedness to delinquency, we find ourselves in the realm of the higher degrees of mental defect, the moron and the borderline cases. These, not the imbeciles, are the ones who present the difficult problem to the student of delinquency, for when those of the lower grade come into contact with the law, their antisocial behavior is recognized as but a manifestation of the deficiency. But the delinquent whom we classify as a high grade defective is not so easily disposed of, and it is this class with which this paper deals.

Only within the past few years have the courts begun to recognize that each case is an individual case and that an understanding of the violator is fully as important as an understanding of the law violated. This has come about largely through the appreciation that a great many of these persons have grave mental defects which were not of a sufficient degree to be recognized by the community. As a result, the juvenile courts of the larger cities and many penal institutions are depending upon the findings of the clinical laboratories to guide them in the disposition of the cases which appear before them. These institutions are finding extremely difficult, almost hopeless, the task of readjusting in society those who from congenital or early developmental causes are equipped with inadequate mental machinery.

A delinquency is an abnormal reaction to stimuli furnished by the environment. There are many conditions which operate to produce abnormal reactions, such as mental depressions, a craving for excitement, instability and, very frequently, a mind not completely unfolded. If judgment, foresight, and moral appreciation are undeveloped, then inhibitions are deficient and the resulting anomalies of behavior will quite likely become criminal acts or delinquencies. The feeble-minded boy is the tool of his environment. He can not see his way forward in the situations that arise, nor can he control his environment. The more complex the situation in which he is placed, the less liable he is to solve his own problems and the greater the probability that his reactions will be construed as antisocial.

The types of delinquencies of boys of the feeble-minded class, then, are many and depend upon the peculiar combinations of circumstances which chance may throw about them in their environments.

The feeble-minded boy is usually a member of a family of degenerate type. This degeneracy may be due to feeble-mindedness itself, to intemperance, or other causes may be to blame, but at least the family has fallen into the lowest strata of society. What mental defect the boy may be given is then of a lower instead of a higher order. The instruments for the implanting of the higher ideas and ideals, the church and the school, are either absent or ineffectual. If the family in its descent has become criminal, the boy is trained in the criminal paths, into which he falls quite readily. If not actually trained, he is encouraged by a family attitude which countenances this sort of thing.

Even though the family is not directly responsible, it has thrown him among associates of the lowest kind and surrounded him by the atmosphere of the slums or, in the small town, of the saloon and the gang. These people will train him, will assist him and will encourage him in starting upon a career of antagonism to law and order. A high level of intelligence is required to rise of itself above surroundings such as this, but our feeble-minded boy stays where he is put. He makes an excellent tool; he does not reason, nor appreciate the full nature of his acts, but he can be induced to perform quite difficult feats for those whose only labor is his instruction and who receive the lion's share of the profits. And, of course, having performed the act, he is the one who pays the penalty if caught, while the real criminal remains hidden.

The public schools, the churches and the charitable organizations have long fought the various instruments of suggestion, the dime novel, the yellow periodical, the moving picture film of similar nature, *et cetera*, most potent factors in the production of juvenile delinquency; but how much more influence do these things have when the mentality is so low that the counterbalancing elements are not present. When the environment is of the slum type, how much additional material for suggestion is presented that can not be controlled other than by the removal of the boy from these surroundings. He does not possess the wider and deeper mental interests, so seeks the activities, which even the feeble-minded demand, in some lower form.

Only recently have we recognized that weak-mindedness is present in a great portion of those boys who fail to progress in school and soon

begin to practice truancy, itself a delinquency. From this it is but a step to the gang with its series of petty thieving. The feeble-minded boy is often the oldest and largest boy in his room and consequently a leader. Under his guidance the younger normal boys learn to play truant, to smoke cigarets, to steal and to practice sexual vices. It is through the moral deterioration of these unfortunate followers that this defective does the greatest damage.

Not being capable of being aroused by the interests that are presented to the normal boy, our defective is looking about for excitement, and this he finds in stealing or other delinquencies. We often find in a good home a boy of this kind, who, although his parents do all that seems possible, is continually getting into trouble and eventually finds himself in some serious situation. Such a case is particularly hard for our juvenile courts, as it is not easy to take the boy from his home, yet the parents are seldom able to deal with him. A mother will overlook the faults of this boy because she feels that he is not to blame, yet she will not acknowledge even to herself that his is an incurable deficiency. She makes allowances and covers up his faults instead of enforcing the discipline that she would give a normal son. The feeble-minded boy is even more in need of strong and rigid discipline, for even this child may learn the effects of fire if it is hot enough.

Space will not permit a discussion of what is called Moral Imbecility, but a few words are necessary at this point. By moral imbecility is meant, to quote Tredgold, "that class of persons who are so constituted that they are utterly devoid of any real moral sense, and of the consciousness that any obligation is morally due from them to their fellows. Such defect is inherent and it may rightly be called moral deficiency. . . . Their moral defect is in fact, latent But although latent, moral defectives of this kind are not of necessity actual criminals; they may well be described as potential criminals."

Many psychologists declare that moral imbecility does not appear independent of grave mental deficiency, some say that it may appear with a deficiency which is slight, while others say that it may be present without accompanying defect of mentality. That portion who have unquestionable mental defects fall within the range of this paper. The moral sense, sympathy, benevolence or social instincts may be lacking, and if the boy is without the intelligence to simulate these instincts or to de-

termine what society considers to be the correct attitude, delinquencies will quite likely result. These are the ones who commit the acts of violence, malicious destruction, assault, rape, sodomy, and other sexual abnormalities and perversions. No hope can be held out in these cases.

Many there are, even of feeble-minded families, who living in the less exacting environment of the thinly populated districts, exist unaided by tilling their small plot or by performance of the tasks of drudgery and attract not even the notice of the communities in which they live. But when the unusual occurs, hardships or difficulties arise, they lack the judgment and foresight necessary to carry them through. Then the easiest way out is the only way, and they come into conflict with the law. This is not the class of the delinquent boy, but is the class into which the delinquent boy often later falls, and this element of his future must be considered in handling his case. This is the class found in large numbers in our reformatories and prisons.

A few cases from our own laboratory will serve to make concrete some of these types:

A colored fellow of very low mentality is serving a sentence for petit larceny. He had once been committed to an Illinois prison for cutting a white foreman who had attempted to take advantage of him by withholding a portion of his pay. While traveling to a point in Indiana at which he expected to find work, he fell in with a white man who took some railroad brass and told this fellow to carry it into the next town where they would sell it and divide the money. Not appreciating the nature of his act, he walked into town with the brass under his arm and when arrested, his director, who had followed at a distance, disappeared. Under a controlled environment and cared for, this subject is a very hard worker at the menial tasks and says that the only mar to perfect happiness is that he can not have tobacco.

One boy of sixteen years is but eight years of age mentally and comes from a feeble-minded family and a vicious environment. His brother has been in our institution and other relations have been incarcerated. When he was eight years old, his mother refused longer to own him and at that age he was thrown upon his own resources. But he soon provided for his future care by performing certain acts which obtained admission for him into the reform school, and he has been a ward of the State almost continually since that time. Besides his mental incapacity, he is physically subnormal and distinctly unstable. He is also a pervert and is a contam-

inating influence even in a reformatory. With such a background as this, a favorable prognosis is impossible.

One boy of sixteen years comes from a family that we have reason to believe is mentally subnormal. The mental examination shows him to be a moron, and our information is to the effect that his environment has not been particularly bad. Though employed, he stole a wheel which he traded for a billy-goat he had long desired.

Another boy is now with us for the second time. His congenital defect is of syphilitic origin and he presents many physical malformations, among which is a nose quite small and deformed. He had previously been in the reform school and was sent to us first for continuing his petty thievery after returning to his home, a small town in western Indiana. He was not long absent upon parole before he returned with a new sentence for horse-stealing, which act was without purpose but probably was due to the suggestion of associates. He recently appeared at our office with a smile upon his face and informed us that he had made a discovery. Upon being supplied with a nail, he passed it in one side of his nose and out the other. It afforded him great pleasure to exhibit this accomplishment to a clinic before a medical association.

One boy of about eighteen, whom the psychological examination showed to be of low level, was convicted upon a charge of petit larceny. In the institution he was a hard worker both in the shop and in the school, but he could not accomplish a great deal. He explained in the most pleading terms to me at the time of his entrance, that he had been without work and while sleeping in a barn he found a fur overcoat which he took because he was so cold. Rather a delinquency than a crime is it not? And the cause? Mental incapacity, the inability to compete on equal terms with his fellows.

From a scientific point of view, one of the most interesting cases we have had was that of a young man of the borderline class, having sufficient intellect to appear normal and about whose home life and environment we have not been able to obtain sufficient data. He stabbed and killed a fellow inmate, almost a total stranger to him, for the sole reason that he wished to obtain a transfer to the State prison, where he would be given tobacco. Even a prison is not a complete protection against a mentality such as this.

Many more instances could be given from this one institution which

are of great interest because of the light they throw upon the essential relations of feeble-mindedness to delinquency, and many more yet must be recorded before the subject will be fully understood.

What shall we do with these feeble-minded and delinquent boys? It is the duty of society to develop their scant capacities and prepare them for things they will be able to do, then to surround them with an environment in which they will be able to do their part and thus get the greatest happiness out of their narrow lives, while society is freed from the menace.

There has been for some time a movement on foot in several States looking toward the establishment of state institutions for defective delinquents where they may be given permanent custodial care. Massachusetts now has such a law, and in New York the Governor last year vetoed a similar bill. Our reformatories are not feeble-minded institutions and can not hold these boys indefinitely. Our feeble-minded institutions have more than they can do with a lower class and are not suited to the requirements of these. In an institution such as suggested, provision could be made for the effective development of the abilities of each one. Each could be given duties that could be made to appeal to his interests and which are within his capabilities. Removed from the competition to which he is not equal, his planning done for him, the cares and troubles to which he is subjected in the world eliminated, his life could be guided so as to give him the maximum of happiness. Furthermore, he would be beyond the power of those who seek such as he to further their own ends. And, again, he would no longer be able to bring into the world others of his kind, to endure a difficult life and to furnish more cases with which society would have to deal.

But since we do not have such an institution at the present time, it is our duty now to do all that we can to assist them with our present machinery. Many of these delinquents have special abilities which can be developed, if we will find them. Already there have been many instances in which the finding of an adaptability has furnished an outlet for the hitherto recalcitrant individuality. We must develop them mentally as far as possible, teach them to read and write, if they prove able to learn, for here some mental interests may be aroused. Closely related to the mental defect in many of the cases, is a physical defect due often to malnutrition and improper care during infancy or early childhood. So far as is possible, these physical handicaps must be removed. The training in

any trade can only be a matter of learning a certain number of movements, yet this may be sufficient to enable them to earn a living on the outside when we are forced to release them. Agricultural pursuits are especially advantageous, as here the demands of the community are not so complex. At such a time as we release them, it is our duty to see that they are placed in environments to which they are best adapted and where the effects of our care and guidance may be such as to insure for them peaceful lives which for society is the greatest protection.

The investigation of delinquents is now being carried on through departments of research in a number of penal institutions throughout the country. In the Indiana Reformatory we are gathering much valuable material of all phases of this subject, but there is one thing which interferes with our efficiency. Every psychologist and psychiatrist recognizes that feeble-mindedness, as well as insanity, is evidenced in other ways than by intelligence alone, and while a psychological analysis will bring forth the defect in the majority of cases, there are a great many of the borderline type that can be rightly understood only by careful investigation of the heredity, family history, developmental history and environmental conditions of the subject. This work presupposes trained field agents upon whom a great amount of work would necessarily fall. In most cases our laboratory now has no information as to those particulars other than that which we have been able to obtain from the inmate, and this is unreliable often because of false representation, but more often from a lack of knowledge of the things desired. Our men know pitifully little about themselves or their families, especially in those cases in which we are the most anxious to obtain accurate information. Many a man has considered it strange that we should expect him to know the year of his mother's death, the number of years he spent in school, the number of times he failed there or whether he was six or twelve at the time of an illness. This developmental period, which is very important to us, exists for him only as a hazy portion of his existence. It is only as legislatures will provide for such needs of our state clinical laboratories that we will be able to contribute to the fullest extent to the thorough understanding of the relation of feeble-mindedness to delinquency.

THE FEEBLE-MINDED AND DELINQUENT GIRL.

E. E. JONES.

It is a signal victory for the cause of education and social service in Indiana that the Indiana Academy of Science has seen fit to place on its program a symposium on "Some of the Scientific and Practical Aspects of the Problem of Feeble-Mindedness." My part in this symposium is to discuss the "Feeble-Minded and Delinquent Girl." She presents a separate and distinct problem in the social and educational development of a State, simply because she is a girl. To her sex belongs the important function of bearing the offspring of the race, and through maternal functionings, mental and physical qualities of the race are propagated. Thus, the feeble minded girl occupies a most strategic position in the problem of race improvement and development, a matter in which scientists have been profoundly interested since 1869, the date when Fances Galton published his wonderful work—Hereditary Genius.

It is fitting that the science of this age devote itself assiduously to the problems of racial improvement, for by no other means than that of science can such a program of civilization be accomplished.

My first problem this evening is to define the feeble-minded girl. What is she? What qualifications, limitations, and possibilities does she possess? What is her type of mind? What is her physical endowment? How much does she know? How far can she be educated? In what sense can she ever become self-sustaining? We must have a comprehension of these facts before we can fully define the feeble-minded girl. At the present time we have numerous scales such as the Binet-Simon Scale for measuring quantitatively and qualitatively general intelligence; and in defining the feeble-minded girl, we must bring into play the use of all such instruments of measurements as will determine her mental and physical endowments. Such scales have been employed in the feeble-minded institutions and with feeble-minded children in the public schools with marked success, and we are looking to a time in the near future when these and other scales will be perfected and refined until it will be possible for us, early in the

life of the individual, to determine the mental and physical defects that produce delinquency and feeble-mindedness.

Feeble-mindedness is a term used to describe individuals who have not attained a normal mental status when compared with individuals of the community. If their mental status is of such a character that they are not able to make the necessary adjustments in a complex social life they are deemed feeble-minded. This does not mean that there are no adjustments that they can make, but that they are able to react only to the simpler situations in life. Some feeble-minded children display remarkable alertness and acute sensitivity, and superficially one would not expect that there is any mental defect; but upon closer examination and study such an individual is found to be deficient in all matters that require complex associations and comparisons. By applying any one of the numerous scales for determining intelligence to such an individual, parts of the scales are answered with very great ease—namely, those parts that pertain to fineness of discrimination in sense impression, either visual or auditory. But when any part of the scale that requires reasoning, associations, comparisons, or complex mental processes is applied to her, she fails miserably. If a child is six years of age and only measures three years in intelligence there is some reason to expect feeble-mindedness. If a child is seven years of age and only measures three years, it becomes quite evident that development is so far retarded that the individual may be very well classified as feeble-minded. Between the ages of seven and sixteen, if the mental age is found to be four years or more below the biological age, there is reason to expect that you are dealing with a feeble-minded person. This is an arbitrary standard that is fairly well adopted by psycho-clinicians in the United States. It is possible, however, to find individuals sixteen years of age who only measure twelve years of age psychologically whose mental retardation is due to disease or other causes than native weakness. Such individuals would form exceptions and should not be regarded as feeble-minded, for there would be a possibility of their very rapid development at a later period; but on the whole we are pretty safe in defining a feeble-minded individual as one whose mental development is as much as four or more years below the normal of a child of his age.

Since having defined the feeble-minded girl, I shall endeavor to treat my topic from three different standpoints. First, how to discover her: second, what are her symptoms? and, third, what shall we do with her?

As to the first question of how we may discover the feeble-minded girl in the elementary schools, and indeed in the homes, I have this to offer. It is my belief that the public schools will have to provide themselves with psycho-educational clinics for the determination early in the life of the child of any sort of mental deficiency. There should be available to every school in the State such a clinic, for no school of a hundred or more children is so fortunate as to be without those whose mental deficiency may be low enough to be designated as feeble-minded. Recent statistics from the New York City schools, Chicago, New Orleans, Omaha and elsewhere, show that about two per cent. of all the children in the public schools are feeble-minded. It is possible that this is too high a percentage, but even if only one child in a hundred is feeble-minded it is extremely important that that fact be determined very early in its life. The psycho-educational clinic will perform the important function in a community of determining, not only all stages of mental deficiency, but also all the stages of mental acceleration; and it is extremely important that those individuals whose rank in intelligence is considerably above the average should also be known and the educational needs adapted to them in a suitable manner. Without such scientific aid in the diagnosis of the child early in her life much energy is wasted in trying to train and educate the child who may be uneducable. Psycho-educational clinics would also serve as a means for determining all grades of mental development in all children and would thus serve as a corrective agency in the proper development of all children. Teachers are generally unskilled in the matter of mental diagnosis of their pupils. They teach upon the assumption that all children have mental capacity about equal. Upon this assumption many a feeble-minded child has suffered punishment and humiliation for laziness, indifference, lack of zeal, inattention, etc., when as a matter of fact the child did not possess more than a third or a half of the mental capacity to do the task assigned. It is my belief that the State should support enough psycho-educational clinics in different parts of the commonwealth as to be available for the use of all teachers and parents. By this agency the feeble-minded girl would be detected early in her life and would be under close observation for a number of years and could finally be disposed of to the best advantage of the public schools, the parents, and the social interests of the State.

My second point is, "What are the symptoms of feeble-mindedness?"

It is safe to say that there are no two cases precisely alike, and it is rare that we find mental conditions that are strikingly the same in different individuals; however, it is possible to name some of the more general characteristics. One of the most prominent characteristics of the feeble-minded girl is mental stupor. She apparently dreams, sits in the presence of certain powerful stimuli unmoved. She is lethargic, inactive and apparently mentally depressed. My own experiments show that the sense organs of feeble-minded girls are about normal, their eyes may be deficient and they may have defective hearing, but this does not seem to be any more a characteristic of the feeble-minded girl than the average high school girl. My data show that the percentage of such defects are about the same for the two classes of girls. It should be said, too, that many feeble-minded girls are supersensitive; their vision is very sharp, their hearing is extremely acute and other senses seem to be abnormally developed. It is only in the organization of this sense material in the higher brain centers that their mental weakness is discovered. The feeble-minded girl normally does not like to play; complex games are difficult for her to comprehend and she can only be taught them with very great patience and much repetition. By the use of the Bergstrom kronoscope, I have secured the reactions of several hundred feeble-minded children. These reactions are both slow and irregular. There is no reliability in the response to a stimulus. Reactions may not be slow in some instances, but they are invariably inconsistent and show great mean variations. I also have the records of several hundred girls as to their vital index which is found by dividing the weight by the vital capacity. The median of the vital index of these feeble-minded girls is several points lower than the median for high school girls. Feeble-minded girls are usually below normal in height and weight. The feeble-minded girl is irresponsible morally, she is not mentally capable of knowing the nature of crime or its ultimate results. She sees in her own acts, which may be immoral, no social significance whatever. She is in no sense responsible for her acts of immorality. Feeble-minded girls are subject to fits of anger and lack of control. This seems to be merely a phase of retardation in her development and is the line of least resistance through which she reacts upon an unfavorable environment. Her acts are nearly all upon the low level of response to sense stimuli. The feeble-minded girl is only educable to a very small degree; she may learn to read, but she can not comprehend

very well what is read; she simply pronounces the words and they are for her the names of peculiar visual stimuli closely akin to the names of persons whom she knows; but these words as groups of words have really no meaning for her.

It frequently happens in feeble-minded girls that there is some special line of action or work in which they can excel; this frequently offers possibilities for education which may be fruitful. These possibilities are easily discovered by the psycho-clinician who may have the girl under observation for a considerable length of time. Frequently feeble-minded girls can do simple sewing, cooking, cleaning, occasionally manifest talent in art or industrial work to a certain extent. Feeble-minded girls are usually strongly sexed. For this reason they are easily brought under the influence of lewd men and are led into immorality. It should be said, however, that in the cases of this kind that have come under my own observation the girl has not comprehended at all the nature of her crime. For her the immorality has been a mere species of play, and she is not at all responsible for her act. Juvenile courts, however, rarely take this into consideration in disposing of the feeble-minded girl. Such girls are usually spoken of in the juvenile courts as sexual perverts. This characterization, however, is a sample of the looseness with which many courts exhibit scientific knowledge in their ministration of justice. It is my belief that many of the girls of this character who are sent to corrective institutions as sexual criminals possess only the normal sex development of the race, and are in no sense abnormal. They have been led into their immorality by men of low character who are ever ready to take advantage of mental weakness, and such girls are so constituted that they cannot possibly comprehend the ulterior results of the sexual act. It is considered no more seriously by them than the gratification of any other sensual pleasure. It must be borne in mind, too, that mere response to sense stimuli is one of the predominant characteristics of the feeble-minded girl, which fact places her far down in the scale of human intelligence, more nearly in the category of the lower animals than that of human beings, who respond to complex situations with judgment and high discriminative powers. The latter she cannot do, because she has not the cerebral connections for such reactions.

My third question is: "What shall we do with her?" This can be answered only in the light of her diagnosis. We must know her mental

and physical possibilities, her powers to respond intelligently to complex situations, such as we find in our own civilization, and whether or no she is able to be self-conservative in a social mechanism such as our State affords. We have already answered this question negatively. We know positively that the feeble-minded girl cannot survive intelligently in the State of Indiana. She will fall if left to herself. She will end in prostitution and crime unless she is protected. The State cannot afford to turn her loose upon society, for obvious reasons. Then what shall we do for her? My statement of the case will be straightforward and above board. First of all she should not be allowed to attend the public school. As soon as she is discovered by the psycho-clinician, when all the expert evidence is in with reference to her and it is positively determined that she comes well within the class of feeble-minded, she should be taken from the public school and placed in a school which is equipped for the careful treatment of such cases. This school should be centrally located for a large territory in the community, and every means should be employed to protect such children to and from school. Parents should be warned of the dangers to which the feeble-minded girl is subjected on the streets, on the playground, in alleys, outhouses and barns and on vacations; and in cases where there is any possibility that parents will not adequately protect the feeble-minded girl from immorality, she should be taken from them and placed in an institution for such mental delinquents for life. She should never marry, for under no circumstances should she be allowed to propagate her kind. In my judgment, as a perfect safeguard to society, she should be sterilized at the pubescent period.

Her education should proceed in such an institution according to lines of her interests. She should be made happy in the work that her likes demand, and should remain protected throughout her lifetime. The State of Indiana probably has at the present time several hundred feeble-minded girls at large, attending no school, under poor parental supervision, running the streets, responding to sense stimuli, gradually going into prostitution, giving birth to illegitimate children, and placing upon society some of her heaviest burdens. It is the duty of the State to bring them under control and save them from the life of social degeneracy which inevitably awaits them if they are allowed to mingle freely with licentious men and are afforded no protection from their sexual suggestions. Her only salva-

tion is in protection from the State, and that protection should continue throughout her whole lifetime.

The question is frequently asked if feeble-minded girls should ever be allowed to marry. In my judgment no feeble-minded girl should ever marry, even though she has been sterilized at pubescence. For the danger lies in the fact that she has not the intelligence adequately to comprehend the meaning of the nuptial tie. The obligations of this relationship would mean nothing to her, and she could not be held responsible for violations of those obligations. If she were free from state control, and should be permitted to marry, even though she had been sterilized at puberty, there would still be the tendency to fall into prostitution and crime which would be unavoidable. It would afford the means of spreading venereal disease and stimulating prostitution, which I feel no State can afford to permit.

One of the greatest social problems of the day for Indiana and all other States is the proper control and education of the feeble-minded delinquent girl. If she is not brought under control, she will propagate her kind, and it is probable that the percentage of feeble-mindedness will increase. With its increase comes added expenditure for state institutions, juvenile courts, medical aid, and waste in education, etc., which increases with leaps and bounds. But if the State takes under its protection and care all feeble-minded girls and boys, there will soon be a great decline in many of the social wastes which at present are sapping the resources of the State. It is difficult to estimate the whole cost to the State of the offspring from one degenerate woman. Fortunately we have a few statistics on this point. The Germans have studied with care the long line of descendants from a few degenerate women, and have calculated their cost to the state. For example, a Margaret Siler, who is characterized as a weak-minded prostitute, was the mother of six children. After 180 years the history of her progeny is as follows: She had 1,286 descendants; of these there were 200 criminals, 280 adult paupers, 300 died of congenital diseases, there were 50 tramps, and she cost Germany \$150,000,000 in legal proceedings alone. Another instance is that of Ada Joirk, a feeble-minded prostitute and drunkard. Seven hundred nine of her descendants have been accounted for. There were 141 beggars, 64 in the poorhouse, 287 vagabonds, and 76 sexual criminals. She cost Switzerland \$1,250,000 in 120 years, and through the lines of con-

genital heredity the terrible work of this one deficient and diseased woman is still going on. It gives us a new phase of the problem of eternal life, and makes it an educational and social problem, rather than a theological one.

Compare such statistics with that of the descendants of Jonathan Edwards, the great New England theologian. One thousand three hundred ninety-four of his descendants were identified in 1900, of whom 295 were college graduates; 13 presidents of our greatest colleges; 65 professors in colleges, besides many principals of other important educational institutions; 60 physicians, many of whom were eminent; 100 or more clergymen, missionaries, or theological professors; 75 were officers in the army and navy; 60 prominent authors and writers; 100 or more were lawyers of whom one was our most eminent professor of law; 30 were judges; 80 held public office, of whom one was a vice-President of the United States; 3 were United States senators; several were governors, members of congress, framers of state constitutions, mayors of cities, and ministers to foreign courts; one was president of the Pacific Mail Steamship Company; railroads, banks, insurance companies, and large industrial enterprises have been indebted to their management. Almost if not every department of social and intellectual progress has felt the impulse of this healthy and long-lived family. It is not known that any of them ever committed a crime, or died in a poorhouse.

Let us protect for life every feeble-minded girl in the commonwealth, and thus cut off one of the most potent influences for social corruption which now embarrasses the State.

FEEBLE-MINDEDNESS IN THE PUBLIC SCHOOLS.

KATRINA MYERS.

“Education,” says Seguin, “is the right of every child, the duty of every parent, the bond of the community.”

In ideal conditions natural talent is allowed to form the basis for training for social usefulness; to each child is given the bent of his natural genius or trade.

For many years our public schools were organized to fill the needs of the mediocre or average pupils. Educational plans that include children at opposite ends of the intelligence scale belong to very recent history. Elastic grading now permits brilliant pupils to skip grades; while an increasing number of slow-witted and even defective minds have special schools adjusted to their needs.

Imagine one hundred ordinary first or second grade pupils. Here investigators find: one stutterer; two or three who lisp; one seriously anaemic; several badly spoiled children; one immature, a year or two retarded in mental and moral growth; one morally weak; two imbecile or feeble-minded. Then, there is one passive, inactive child; several over-sensitive, nervous children; one superficially precocious child, and several superior—eager, ardent, imaginative, social. Four suffer from defective hearing; twenty-six now, or will very soon, show eye strain or have defective vision; about a dozen have asymmetries or deformities; about thirty have nasal obstruction or diseased throats; and several others possess serious peculiarities of temperament. Only twenty-five of the hundred are physically and mentally without blemish.

All these children represent an actual, positive asset in human society. If they are not saved for constructive activity, many of them will become a destructive force later on.

The question arises: How are the powers of individual pupils to be definitely known, so that schools may measure to their greatest efficiency?

Some of these peculiarities are related to the size and shape, observable defects; some are alterations of internal structure, not apparent to

the untrained observer. Many pupils, who have observable peculiarities, are very capable mentally and give normal response to training; while many, who baffle the most conscientious teaching, present no outward signs of disordered organizations at all.

To remedy all cases of mental and physical deviation, there must be definite localization of the defects, which only thorough medical, psychological and pedagogical examination will reveal.

The earlier suspected abnormalities are discovered, and proper corrective treatment and training are given, the greater chance these children have to become more nearly like normal beings. Brain energies are broken by physical irritations and other strains. This is why certain conditions produce retardation and an arrest or paralysis of the inhibitory or moral sense, and explains why removal of disturbances is often closely related to moral and mental regeneration.

For years psychologists have been endeavoring to formulate some intelligence measuring scale that could be applied to the age and grade of the average pupil. The great benefit that would accrue to both pupils and teachers from an accurate intelligence standardization can hardly be estimated. Then, only, can training be given each child that will insure full individual development.

The Binet-Simon Measuring Scale of Intelligence is a series of questions "arranged in groups according to their difficulty as determined by age difference in performance". The questions relate to general intelligence, to information that the average normal child should absorb from every day associations and not to what he is taught at school. The insufficiency or retardation of backward children is later estimated by comparison of their results with those of normal children. The series is merely a sorting test; but, in the hands of experts, it has been amply demonstrated that it is very valuable, and gives a surprisingly close estimate of a child's mentality.

A child who has for no adequately known reason fallen behind two years in his school work, should be carefully tested and watched. He may be ill; his mind may be "slowing down". When children are found three or four years behind children of their age, the intelligence tests will undoubtedly disclose more serious conditions. It is well recognized that minds of most educated persons reach the limit of intellectual development between the ages of twenty and forty. Minds of the great mass of

mankind reach the limit of development between fourteen and twenty-one. A mind that never gets beyond thirteen is just able to make a living. Above eight and below thirteen comes the moron, a person between normal and imbecile. He can be taught routine tasks, lay bricks, make parts of shoes, do tailoring, farm work, etc., as well as any one, provided some one else does the directing and planning; he never gets beyond twelve years old. True imbeciles never develop beyond seven. The mind of an idiot closes before three.

When the brain stops developing—and it may occur as early as the third year—the time is practically past when it is possible to give a training that will help the child to earn a living.

Any one who deals with a large number of persons realizes how intelligence varies from those with practically none to the very gifted, and that responsibility varies according to the intelligence. Some of these persons, under simple environment, seem to function normally; but when placed where the environment becomes too complex normal functioning becomes impossible.

Among a number of misfit pupils observed in grade classes and recently tested, were two, thirteen years old, who had made no real progress for five years; two, eleven years old who had lost four years; a girl of thirteen who had made no real progress for six years; but all had been promoted, though the work accomplished had been, at best, mere rote work, with no more real intelligence than that of a parrot taught to say a rhyme. Two girls of seven years were mentally less than three.

To attend a class where normal children are receiving instruction does not help the undeveloped child. "Learning can not penetrate like a cold storage chill." "Mind building is like house building; there must be a foundation on which it rests."

Undoubtedly all intelligent persons agree that, no matter what a child's lack of mentality, the hopelessly idiotic and imbecile types should be trained in the schools the States maintain for their segregation and care. These most deficient children comprise only a very small per cent. of the school membership ($\frac{1}{2}$ of 1%). Often the most troublesome public school cases are pupils of the borderland types; those just between normal and subnormal.

Since the compulsory education law exempts children, mentally and physically disabled from its operation, these pupils, and those of lower

mentality, having become annoying, in the past had been habitually excluded from school. These excluded, subnormal children, untrained and undisciplined, have often become the men and women scourges of society. Many teachers, principals and physicians consider such pupils as merely slow, and earlier in their school life, fondly expect them to brighten up; but, the incapacity once recognized, the problem is an altogether different one; for, when we measure the intelligence, we have measured the degree of the irresponsibility. Their right to a training is the same as is that of their better endowed brother, and their need is greater. To attempt to lead a feeble-minded child along the school way of the normal child can result only in failure.

How would you feel if you were fourteen years old, with a mental capacity of a child of seven, obliged to sit in a 5A grade class, absolutely incapable of understanding the work? Can you blame a child for rebelling against such an overburdening "education"?

Misfit children are naturally affectionate, kind, and tractable. Maliciousness and carelessness are not their inherent traits but are acquired from lack of capacity to understand required conditions. Continued misunderstanding and rejections make them hopeless and rebellious.

A desire for greater school efficiency is demanding that classification must be made on a basis of native ability. For hundreds of our public school pupils, the hope of escape from a life of utter inefficiency lies in the ungraded schools of our public schools. For here only can they *now* be given training and treatment adapted to their subnormal, individual capacities. Destroyed brain tissue can never be renewed. We know our limitations. It is not possible to make good out of a poor thing; but wonderful things can be done.

The widespread need of such schools can be estimated when you consider that reliable investigators compute the number of feeble-minded pupils to be from two to four per cent. of all children of public school age. All cities, in our country, of 100,000 population and over, and many small towns, now maintain separate schools for feeble-minded and defective children, in care of more or less specially trained teachers. Indianapolis now has two such schools, in which thirty-one pupils are enrolled. According to the lowest expert estimate, this is $4\frac{1}{2}$ per cent. of the feeble-minded and defective children now in the grade classes that need to be given special opportunities.

Perhaps it is well nigh impossible for most persons to become as little children who live under the crushing conviction that they have no brains. Failures and rejections burn deep gloom into sensitive child minds. Their errors should be corrected by putting in their place more practical, simple truths.

Since books spell discouragement to the dull, inapt mind, it is generally wise to excite an interest in things unrelated to failure; for expected failure creates a hopeless habit of thought. For these pupils there should be an abundance of stimulating devices to excite an interest. These minds have lain dormant for years, or have *never* been awakened to natural child curiosity. This varied stimulation should be thought provoking, not nervously exciting.

Frequent change in manner of presenting old forms, and gradual introduction of new material arouses confidence. Successes create assurance; and, even if the progress be slow, it surely follows. Success is a relative term; but it carries the highest valuation in mental and character development.

Manual work offers great variety and simplicity in subjects and employs and trains more faculties than any other school work—observation, attention, concentration, comparison, coördination, decision, judgment, all are involved in its simplest problems. So, it is admirably adapted to the needs of the undeveloped child.

Almost every child's best is good in something; and it is only by our honest trying that we shall be able to draw a finer and better efficiency from the unused and often ill-directed capacities of children who possess limited possibilities.

Having discovered an underlying trait, something for which the child has liking and ability, the worst struggle is over. Then, thru the newly discovered aptitude, it is comparatively easy to bring the pupil into natural association with other school duties. Tasks and lessons coördinate. He reads to learn, and even numbers have a new meaning. He enters with zest into games, songs and all school exercises. In the natural life of the schoolroom the child is socialized.

The training requires the hand of iron in the glove of velvet—thoroughness, patience, resourcefulness, open-mindedness, sympathy, hope. The wisdom of Solomon, perhaps, could not always solve the problems these children present. That is why failure now so frequently attends our ef-

forts. But we know that everything that makes these children—who never can become men and women in intelligence—more capable with their hands, more reasonable, better self-controlled, more helpful to themselves and to others, will assure them a more certain degree of success as individuals and in their relation to society.

I summarize under six headings:

1. The State should demand and provide careful medical and psychological examination of all children in the grade schools, who are two or more years retarded in school work.
2. The State should provide for correction of all physical defects in children diagnosed as having *remediable* defects.
3. School systems should be obliged to organize and maintain separate schools for all feeble-minded and mentally defective children now in grade classes.
4. Schools for feeble-minded and mentally defective children should be in charge of specially trained teachers and supervisors.
5. The Compulsory Education Law should be amended so as to include in its operation children *not* in good mental condition.
6. The State Institution for Feeble-minded should organize and maintain a department where teachers for feeble-minded and mentally defective pupils can receive practical and theoretical training.

NOTE. Acknowledgment is made, for some valuable statistics used, to Dr. H. H. Goddard, Dr. W. J. E. Wallin and Dr. M. Groszmann.

THE ALCOHOL PROBLEM IN THE LIGHT OF CONIOSIS.¹

ROBERT HESSLER.

The alcohol problem is a very important one, so important that it has entered politics. Men vote wet or dry, and women are seeking suffrage. If women vote and vote "dry", will we have prohibition? Will the passing of a prohibition law free the State from the alcohol problem?

There are all sorts of factors that enter into the question, and there are all sorts of problems connected with the alcohol problem. Here I desire to call attention to one that is usually neglected, the dust problem or the bad air factor. (How to develop the subject in a 20-minute paper is in itself a problem. Technical details must be omitted.)

The study at first was one of individuals and then one of the family, and finally of the entire relationship. I shall show a number of charts. Some represent several hundred individuals and are so full of details that it would require considerable time to go over them. Most of these charts will be referred to very briefly in order to give as much time as possible to the deductions or conclusions.

A word regarding the charts that were shown: On account of the difficulty of reproducing them properly in colors they are omitted here; only one is given to illustrate the general idea, how the data were diagrammed. Perhaps the few references to charts will be understood in the light of the diagrams. In the original charts, or genealogies, males were indicated by blue lines; females by red lines. Black lines represented members whose sex was unknown. Short lines indicated individuals who died young. The children, sibs, are given in the order of age, beginning at the left.


Intermarriages are shown by short lines, drawn upwards.

No offspring (that is, race suicide,) is indicated by a circle.

The accompanying diagram represents three generations.²

¹Coniosis. For an explanation of the term and of the theory of Coniosis see Proc. Ind. Acad. Science, 1911.

²The original blue lines (males) are here represented by light lines, and the red (females) by heavy black lines.



1. A father and a mother.
2. The three children: a son married (with one son living and one son dead); a daughter, unmarried; a daughter married but childless.
3. The young generation.

A number of charts were shown, some of five or six generations and representing several hundred individuals. As a rule those living under simple life conditions, as in the Northwest, still have old time fertility, while those living in towns and cities show a gradual decline and even total extinction.

Among charts shown may be mentioned that of a family where for several generations, after coming to America, the sibs of each generation numbered about twelve. The parents being two, a family of twelve children of course means a six-fold increase. The moment the more recent descendants began to live under town and city conditions, the death rate increased, many did not reach maturity at all, and some of the living have no offspring, race suicide appeared with a vengeance; also narcomania, i. e., the desire for strong drink and narcotics.

Why do men drink? There are all sorts of reasons. The three chief classes of drinkers are:

Social drinkers, keeping up old-time customs.

Habit drinkers, some think they require an appetizer; others, something to settle their dinner, to aid digestion, etc.

Relief drinking, as where a man feels bad but feels better, or thinks he does, after a drink or after several in succession. It is with this class that this paper is mostly concerned. Such men as a rule "pour it down," they are not particular about flavor or taste, they drink for the effect; they do not take strong drink as a "stimulant," just the opposite, as a sedative. Writers usually speak of this sort of drinking as "Misery drinking." Relief drinking is a better name in most instances.

We must constantly ask of the man who drinks or who drinks to excess, Where, when, why?

The man living in isolation, the moonshiner, or the farmer, who can drink alcohol with impunity, may find it a powerful poison when he removes to the city. Alcohol and infective matter in the air (and of course also in water and food) make a bad combination. The body may be unable to overcome their combined influences.

Reliable data concerning people who use narcotics are difficult to ob-

tain. If one asks directly many become offended. Some hesitate to speak of the "black sheep" in the family; and the black sheep themselves may magnify the failings of others. Often black sheep are not as black as painted. Without knowing the facts in the case, we should not be too severe on the man who drinks or the man who requires the use of tobacco for his well-being, as he believes.

A large chart was shown of six generations since the original English immigrants, the relationship now consisting of several hundred, but only a few branches are more or less known.

Here we are at once in the midst of things. There are all sorts of deteriorating factors, including narcomania. The alcohol problem crops out all over one-half of the chart.

Some remember the days of "free whiskey" when everybody drank and when it was no disgrace to be drunk. Today there is a different sentiment and one may well ask, Why does an apparently sensible man drink and drink to excess? Why does he drink to excess occasionally or not quite to excess at all times? To say a man drinks because he wants to is not a satisfactory reply.

Many patent medicines are full of alcohol and their effect on the body is that of alcohol. Some men who are ashamed to go into a saloon, or to call for plain whiskey, use nostrums that are compelled to pay the whiskey tax. Such preparations are used because they give ease; the man or woman using them "feels better". As in the case of plain alcohol, the dose must gradually be increased, or more frequently repeated, to get the desired effect, that of a sedative.

Many of the subjective complaints of the men and women who use alcohol are similar to those of neurasthenics and hysterics, so called. Many complain of "fear, moods, depression, sleeplessness, restlessness, tremors, general weakness, tearing pains, anorexia, palpitation, etc." According to my observations many individuals who are regarded as neurotics or neuropathics are dust victims, and because narcotics give relief, they use them. A study of the chart will show this.

Red squares in the chart indicate a victim of narcomania. At times the habit of using alcohol or morphia or even aspirin was acquired through a doctor's prescription. Physicians nowadays are careful for whom they prescribe narcotics; they fear habit-forming drugs. Many physicians themselves are said to be victims; long and irregular hours, loss of sleep,

etc., including attendance on patients living in unsanitary homes, even their own ill-ventilated offices, may lead a doctor to seek relief.

A green square indicates country life, or good air life, in contrast to the yellow, or polluted city air life. The half and half squares at the bottom mean that some members are living in the country, others in villages, town and cities.

Black lines indicate no definite data regarding individuals.

There are eleven children in the first generation in this country. Today the descendants of one seem to outnumber those of all the others; this was a man who married a strong-minded German catholic. The chart showed that these lead mainly a simple country life, with large families and a practical absence of narcomania.

TIME SPENT IN GOOD AND IN BAD AIR.

Primitive man	0/24	0/7	0/365	
Hunter and trapper	0/24	0/7	x/365	
Squatter	0/24	0/7	x/365	
Farmer	0/24	x/7	x/365	
Villager	x/24	x/7		} 8 hours of labor. } 8 hours of recreation. } 8 hours of sleep. } Under what air conditions?
Townsmen				
Cityman				
Slum dweller	24/24	7/7	365/365	

This chart (table) is an attempt to express in a general way time spent in good and in bad air. At one extreme is the savage, living say in the mild tempered South Sea Islands, with good air at all times; at the other extreme is the slum dweller with 24 hours of bad air, 7 days a week, 365 days a year.

Air conditions under which people live can not be considered collectively; each individual must be studied separately. Ordinarily we assume that a farmer is leading the good air life, but when the farmer comes to town every Saturday, or perhaps daily in spring and fall, and loafs for hours on street corners, with clouds of infected dust blowing about, he may carry home enough infection to last him for days.

The ideal of the union labor man is the eight-hour day. But we must question where? Under what air conditions—good, bad, indifferent—

are the eight-hour periods spent? Is the recreation time spent at home in the suburbs making garden or in outdoor air; or is the time spent in the heart of a dirty city, in a saloon, pool rooms, or loafing on the dusty street corners? The five-cent theater habit is only too often accompanied by the bad habit of "going out to see a man" and chewing a clove after it. In short, the time spent in recreation under bad air may be the real factor for inefficiency and premature breakdown and not the work in a clean and well-ventilated shop.

Eight hours of sleep: In what sort of bedroom? With good air, free ventilation? Or closely housed and with perhaps no daylight coming into the room at any time?

The slum dweller of course represents the worst conditions in all respects. Naturally the weeding out process is actively at work. To see the end results at their best one must turn to the overcrowded cities of China and India—with bodily adaptation at the expense of mentality. Our unsanitary cities in time produce a class of people little different from John Chinaman, with all that that implies, including the use of narcotics and sedatives, if not opium then tobacco and alcohol, or cheaper coal tar preparations.

THE WHERE, WHEN, WHY OF DRINKING.

A question, or rather several questions, a student must constantly ask is, Why do men drink? Why does an apparently sensible, decent sort of man have a craving for strong drink? Why do some demand it more or less constantly, some periodically? Under what conditions is the craving most marked? In short, Where, When, and Why do men drink?

Here are a few representative cases. The figures are of course only relative; one can not express the complex life of a man mathematically, there are too many exceptions.

GOOD AND BAD AIR AND THE DESIRE FOR DRINK.

Mr. W., farmer	0/24	0/7	4/365	(4 times a year to town = sick)
Mr. H., farmer	0/24	1/7	75/365	(visits to church and town)
Mr. X., professional	8/24	6/7	290/365	(15 days in wildwoods)
Mr. X., businessman	8/24	6/7	300/365	($300 \times 8 = 2400$ out of 8760 hours)
Mrs. X., the wife	x/24	1/7	75/365	($75 \times 2 = 150$ out of 8760 hours)
Mr. B., mechanic	24/24	7/7	365/365	("Always thirsty")

Explanations of the chart (table) must be brief.

Mr. W. leads a quiet isolated life on the farm; has retired from active work; is well-to-do. He lives in good air twenty-four hours a day. In the summer when doors and windows are open he attends a country church; in winter when doors and windows are closed he remains at home; says he can not bear close air. In recent years he comes to town about four times a year; formerly he got sick on or after every trip, "would get faint and dizzy and feel like falling," and friends only too often offered him a drink of whiskey, to which he was opposed. When it was pointed out to him that he was a marked dust victim he in time learned how to guard himself; he no longer comes to town when dust clouds blow about and spends little time in dusty buildings.

Mr. H., an active farmer, lives in good air twenty-four hours a day, that is as far as infection, found in city dust, is concerned. Sundays he attends a village church. Occasionally he comes to town, perhaps fifteen times a year; his absences from home he thinks amount to about seventy-five a year, varying from an hour or less to several hours. Formerly when he came to town he had an almost irresistible craving for strong drink, i. e., on dusty days, as I soon found. He felt bad, was miserable; he learned that one or more drinks made him feel good, and only too often the one or two drinks increased to a sufficient number to make him drunk; and then there was great remorse and he vowed never to drink again. But until he learned *why* he had such a craving, how it depended on inhaling infected dust, he only too often could not resist the use of alcohol. In the country when he was tired and would "spit cotton," water would quench his thirst, but the phlegm due to spit dust required something stronger "to cut it."

Mr. X, a professional man, works in the heart of a city eight hours a day for six days in the week. He wants to smoke all the time and frequently wants a drink; at times he is on the verge of being drunk. The desire for strong drink is almost irresistible "when there is strong mental strain", as he supposed, but as a matter of fact it was at times of close confinement to ill-ventilated offices and public buildings. Now the most interesting phase in this man's history, the where, when and why, is this: For two weeks or so every year he goes off to the northern wildwoods; then he has no desire for strong drink, scarcely touches it, and there is little desire for smoking, that is, there is no excessive smoking.

Mr. X, a business man, working, as he believed, "under high pressure."

Certainly he was always "steaming up," constantly smoking and many times a day taking a drink. He did not give me an opportunity to study him fully but I found he had a blood pressure of over 200 mm., nearly twice the pressure an open air or pure air man carries. To such men alcohol gives relief; their drinking must be looked upon as "relief drinking." When he came to understand the danger from a high blood pressure he wanted to know what to do. The proper thing is to reduce hours indoors, offset bad air influences as much as possible by good air influences. But he was the sort of man who took something rather than do something. Is there any drug that will keep down the pressure that can be taken in place of alcohol? Why yes, a host of them. The question here arises: To what extent shall a physician recommend the substitution of one drug for another? Is any drug at all harmless? To depress the bodily activity, to depress the blood pressure artificially is not good medical treatment.

In connection with this business man should be mentioned his wife. She had chronic ill health when she first came to me. She was "low pressure," "could not keep up steam" in fighting off infection. She now lives up to *good* air advice and has reduced her ills, her reactions, her symptoms, to a minimum. There are perhaps seventy-five annual "exposures," as by going to church, shopping, theater in summer, etc. If the average length is two hours, there would be about 150 hours of more or less polluted air inhalation in a year.

The husband stays at home Sundays in good air, but his eight hours of weekday exposure amount to at least 2,400 hours a year, out of a total of 8,760 hours.

Mr. B. is a mechanic, almost a "mere laborer," who works with his hands; little mentality is required. He lives in the heart of the city over a store where rent is cheap. He works in a dirty shop. He gets bad air, air contaminated by infection, twenty-four hours a day for seven days a week and for nearly the whole year, the only time he gets good air is when four times a year he and his family visit "the old folks on the farm." This man is "always thirsty;" he "always needs something to cut the phlegm." Many patent medicines full of alcohol serve the same purpose as alcohol in the form of whiskey, brandy, gin, etc. His only objection to beer is "It takes too much to get relief." Such a man will scarcely listen to a physician who tries conscientiously to help him. The

evolutionist will say nature still weeds out those who can not resist using alcohol excessively.

There are all sorts of reasons why men drink. I am here only calling attention to a neglected factor. Alter the environment or remove a man from the abnormal environment and, unless he has deteriorated too far, he will likely cease to drink or to drink to excess.

Many of you are teachers. Apply the formula to yourselves: Under what air conditions do you spend your time? How good or how bad is the air under which you work, and must work, and the air at home and on your vacations? The teacher has a limited number of hours a day at school, with two days of rest a week, with holidays now and then and an annual vacation of several months—time for lymphatics to empty themselves, ready for the next season's work.

At what time do you feel "fagged out?" At what time do you feel the need or the desire for a sedative (tobacco, alcohol, opiate, coal tar preparations, etc.)? When is there a call for a stimulant (coffee and tea especially)?

Today we hear much of efficiency. Under what conditions is a man most efficient? To what extent is the air factor considered in estimating efficiency? The above formula might perhaps be of service in working out some cases; and similarly in the matter of longevity. How many reach a possible hundred out of the many born in country, village and city? Is a relatively short "high pressure life" of forty-five or fifty more desirable than a longer life under less strenuous conditions? Is fifty years (or often much less) better than "a cycle of Cathay?" Is the attempt to become adapted to dirty cities worth while—when it is at the expense of mentality?

In a study of the alcohol problem and its evils one is constantly reminded of the fact that man is an outdoor animal and that he thrives poorly in houses and cities. Geologically speaking it was only day before yesterday that he attempted to live in houses; yesterday in cities; and today he is undergoing the weeding out process as never before. Man of course will not die out, there will not be complete race suicide.

Man afflicted by the ills of domestication and urbanization seeks relief. There are two methods:

Take something, usually some drug that gives temporary relief.

Do something, alter an unfavorable environment, or if that is not feasible, remove to a better one.

Only the physician knows how prone people are to take something and how they hesitate to do something.

The history of medicine itself is one long search for panaceas, for "cures." We are only beginning to realize that the search is futile. We more and more see that the proper method of securing survival of the best is to annihilate the pests and parasites that prey on us. Just as weeds and pests do not thrive under clean culture, in the same way disease germs do not thrive in a clean city among clean people; quack remedies are as useless in the city as they are useless on the farm.

There are always some who will survive most adverse conditions; this can be seen in the crowded unsanitary cities of China¹ and India, and in our own slums. But such a type is not what the believer in a better humanity has in mind. People who constantly battle with infection have so little mentality left that they are but little above unreasoning brutes. Fittest to live under an unsanitary environment, abnormal in the light of man's past history, does not mean best mentally.

Man in his search for cures in time has found means for relief. He has discovered a series of substances that give ease; they may be grouped under the name of narcotics.

Many plants have such properties, from the poppy with its opium and codeine and morphine, down to mild ones like hops. In time man has learned to prepare alcohol, varying in strength from mild, as in fermented milk and fermented honey and fruit juice, up to distilled alcohol in its pure form.

Alcohol is an old remedy. It constantly appears in new disguises, under the fanciful name of some nostrum or patent medicine. Recently a number of so-called synthetics have come into use, made from coal tar. Their number is constantly increasing. An individual in search of ease soon learns which suit best; unfortunately many are of the habit-forming kind.

NARCOTICS OF THE METHANE SERIES.²

Hydrocarbons: Pentane, Octane.

Hydrocarbons, unsaturated: Amylene, Pental.

Alcohols: Methyl, Ethyl, Propyl, Butyl, Amyl. Amylene hydrate.

¹Even the supposed adapted or immune Chinese have their ills and their means of transient relief. Now that the opium trade is imperilled they will likely resort to the white man's remedies, alcohol and tobacco and synthetics.

²This list was compiled from Cushney.

Ethers: Ethyl, Ethyl acetate, Ethyl nitrite. Amyl nitrite.

Aldehydes: Paraldehyde, Methylal, Acetal, Sulphonal, Trional, Tetronal.

Ketones: Hypnone.

Esthers: Urethane.

Acids: Butyric, Bromacetic, Chloracetic, Formates, Propionates, etc.

Halogen Substitutions: Chloroform, Ethylene chloride, Ethylidine chloride, Chloral, Buthyl or Croton Chloral, Chloretone (Trichlorpseudo-butylalcohol), Chloralamide, Chloralose, Bromoform, Ethyl bromide, etc.

Ethyl alcohol is the alcohol used for drink in the various alcoholic preparations. Chemically and pharmacologically alcohol is in suspicious company, among chloroform, chloral, ethers, and so forth. Some preparations are solids, as chloretone, whose chemical name ends in alcohol.

There is an almost endless variety of synthetics and new ones are constantly appearing "Made in Germany"—and we are the best customers.¹

One can scarcely speak of any as real "cures," but they are palliatives; they give ease from the ills of civilization. The more people are massed under unsanitary conditions the greater the demand for them, indeed the sanitary condition of any community can be estimated by the demand for such remedies.

In connection with this Methane series should be mentioned the Aromatic series; among the chief are: Terepene, Menthol, Guaiacol, Resorcin, Phenacetin, Lactophenin, Sedatin, Phenocoll, Salicylic acid, Salicylates, Salol, Aspirin, Salophen, Analgen, Antipyrine, Salipyrine, Hypnal, Cocaine, Eucaine, Novacaine, Holocaine, etc.

These have a reputation for relieving pains and aches variously referred to as rheumatic or neuralgic, including many forms of headache. Although some are obtained from plants, the great majority are synthetically prepared from coal tar; new ones are constantly appearing.

As already mentioned, individuals differ greatly. Those who do not

¹Twenty years or so ago, when engaged in work among the insane, I tried many of these preparations. At times sheriffs who brought in new patients preceded me in experimenting, as where alcohol was given to make obstreperous lunatics tractable; more than once patients were "dead drunk" and of course offered no resistance.

Chloral, paraldehyde and sulphonal, etc., were used when patients were restless and sleepless; some readily acquired habits and demanded a "nightcap." Chlor form and ether were of course used in surgical cases.

One of the most interesting of the series is Amyl nitrite a vasodilator much more effective than alcohol, acting almost instantaneously. Among the insane it was used in cases of continued epileptic convulsions, often with doubtful results.

get the desired relief or ease from alcohol, or who have enough pride not to resort to its use, or who fail to get relief from opiates, or who fear taking such on account of habit-producing effects, may find a suitable remedy among this Aromatic series.

That some acquire the aspirin habit has already been mentioned. There are all kinds of dope fiends and new ones appear every now and then.

I have not looked up the statistical aspects of this matter. In the case of the most commonly used narcotics or sedatives we all know that the annual bill for alcohol and tobacco in our country is enormous.

It appears that in 1912 the United States used 135,826,000 gallons of distilled spirits, and 1,925,367,000 gallons of beer.

There may be some justification for the use of beer, a mild alcoholic drink, but there is little to be said in favor of strong alcohol, of which only a few ounces can be consumed by the body in 24 hours. It seems some European countries have diminished the consumption of strong alcohol, while in our own country the amount consumed seems to be increasing. With increasing sanitation in old world cities there is less demand for strong drink, less need or demand for relief or misery drinking. Although there is a greater consumption of alcohol in our own cities, there is less actual "drunkenness," using the term in its old significance, i. e., being completely overpowered.

A chart with hundreds of individuals was shown where race suicide was strongly operative, and, although there was narcomania, there were practically no individuals who drank to the point where they would fall into the hands of the police.

Although at times a family may be reduced to one individual, that individual may be worth more to the community than half a dozen in the slums. The fable of the fox and lion applies.

In contrast to the last chart was shown a large one of a family which is in the main still rural, and yet there is gradual disappearance of the old colonial stock. There are all sorts of factors for race suicide. The bad air factor is usually overlooked. Domestication and urbanization and consiosis are usually not considered.

The botanist constantly speaks of "naturalized plants." In the case of annuals it does not require many years of observation to determine whether a plant is really naturalized, that is, whether it can continue itself successfully year after year.

From a study of many family histories, one may question whether the white man, the European, is truly naturalized in this country.

How the use of alcohol and opiates crops out in biographies, also the matter of race suicide, is an interesting subject. In the light of my study of people who drink it is not difficult for me to understand why a man like Poe drank.

The family chart of Herbert Spencer is interesting (chart shown). He was a dust victim, and so were his parents; he was the single survivor out of nine children. He experimented more or less with narcotics and sedatives in his search for relief from symptoms of ill health.

THE EVOLUTION OF DRINK AND NARCOMANIA.

Hunting and Fishing Stage:

- "Home life" very simple.
- Absence of fermented drink.
- Use of vegetable narcotics at ceremonials and festivals.

Pastoral Stage:

- Nomadic tent life.
- Use of leather bottles and fermented milk, a weak alcoholic drink.
- No special desire for "stimulants" under simple life conditions or by outdoor people.

Agricultural Stage:

A fixed home implies domestication, the ability to live under indoor conditions for successive generations.

Domestication means:

- Re-breathed air
 - Soil pollution
 - Water pollution
 - Stored food
- } Attended by "ills of domestication."

Many "incurable ills" or "diseases" are preventable reactions.

Invention of pottery and fermented drinks from fruit juices and grains, wine, cider, beer.

Use of alcohol for relief from disagreeable symptoms. ("Symptomatic treatment" survives today).

Additional use of various narcotics: Opium, henbane, hasheesh, etc.

Handicraft Stage:

Town and city life means urbanization.

Great increase in house and town ills.

The "ills of civilization" are reactions, largely preventable.

International commerce and introduction of cosmopolitan diseases, that is, specific diseases of definite etiology.

The search for remedies or "cures" for ills and diseases incident to house and town life.

Discovery of distilled spirits.

Alcohol regarded as a panacea, first by physicians, then by the people.

Names used: Aqua vitae, Au de vie.

The role of religions, favorable or unfavorable to alcohol.

Severe weeding out and adaptation of humanity to unsanitary city conditions.

Tobacco (introduced into Europe about 1586) a factor of increasing importance in prevalence of ills and diseases and race suicide.

Industrial Stage:

Cities and manufacturing towns the graveyard of man.

Constant need of fresh country blood.

Overcrowding and lack of sanitation = prevalence of specific diseases

Daily and constant exposure to dusty air = prevalence of coniosis.

Excessive use of strong drink and "dope" = narcomania.

The Alcohol Problem a serious one in dirty cities and complicated by the tobacco habit = the "Tobacco Problem."

The tobacco habit produces a "spitting habit" and sets a low standard of cleanliness.

"Spit Dust" is responsible for the prevalence of so-called "American Diseases."

Catarrh, dyspepsia and nervous prostration are mainly reactions to an abnormal environment to "bad air."

Speeding automobiles an important factor; make clouds of dust.

Blood pressure an important factor in the ultimate fate of the individual.

Alcohol, sedatives and narcotics give relief and lead to "relief drinking," to inebriety and drug habits.

Such a chart could be extended indefinitely; here are some additional observations:

Pure air people are temperate people.

People living in good air do not crave alcohol or dope, not even a cathartic pill.

The number of saloons, tobacco shops, drug stores, of patent medicine advertisements in newspapers and of doctors, shows an unhealthy state of affairs. Another sign of national decay and race suicide is the reversion to primitive beliefs, faith and mind cures—a mode of treatment resorted to by people when in despair at the medical profession failing to cure or benefit. Many ills are incurable, they should be looked upon as reactions, not diseases.

Narcomania is exceedingly prevalent in our cities and towns, but men who are reeling drunk are less and less in evidence. The vicious are rapidly eliminated.

"A sound mind in a sound body" fails to consider the influence of environment, how an abnormal environment weeds out the best mentally. The robust teacher fails to understand the delicate child that reacts to an abnormal environment. The best barometer or thermometer for a school-room is a teacher who is not too robust. The best physician for prevalent ills is the one who himself is not too robust, not too immune.

Survival of the fittest does not mean survival of the best when applied to unsanitary city conditions. If this were true the people of crowded Chinese or Indian cities would head civilization.

It still holds true that "the good die young" on account of unsanitary life conditions. Many are killed off by alcohol and narcotics. Many a young man of promise finds his death in the cup.

Several charts were shown to illustrate how the tobacco habit and the spitting habit are related to the alcohol problem, and in turn to the race suicide problem, how unsanitary air conditions lead to prevalent ill health and to terminal infections that kill. Charts were shown based on the census reports, from which it appears that in our State today the rate of decennial increase is constantly diminishing, and likely at the next census there will be a deficiency, in other words, the loss will be greater than the gain. In Northern Indiana only a few counties have gained in population. those with industrial cities.

Industrial cities, like unsanitary cities, have been compared to huge parasites that drain the country of its best blood. Such cities have little use for a man over 45 or 50. Yet such cities may point with pride to their low death rates. The explanation is of course simple: Worn out men go away to their old homes, to die.

Some cities, really overgrown villages, have a bad water supply, and the brewers advertise their clean or pure beer; yet the Prohibitionists are making little or no effort to get good water. Is it any wonder that many cities vote "Wet?" The first effort of the Prohibition advocate should be to give the people clean water and clean air. Fresh water does not necessarily mean pure water, nor does fresh air mean clean air. Saloons flourish in proportion to their unsanitary surroundings and the patronage of low grade laboring men.

The solution of the Alcohol Problem depends upon education and cleanliness—clean people, clean homes, clean cities, clean streets, clean water, clean air. In the light of Coniosis the greatest of these is clean air.

COLD STORAGE IS PRACTICAL CONSERVATION.

H. E. BARNARD.

Cold storage is essentially the application of scientific temperature control in the solution of an economic problem, a practice that regulates prices without increasing them and prevents deterioration while eliminating waste.

But to the average consumer there is no hint of conservation in cold storage and little reason for the practice except that born of greed. June butter in January, spring chicken at Christmas, fresh eggs months after they were taken from the nest, summer fruit in winter weather—do these reversals of the season's horn of plenty, this carrying the products of flush markets over the time of scant production, increase the cost of food to the consumer, reduce its value to the producer or in any way injure the masses of the people who by its consumption and in its production find health and wealth?

The world's development has been along the lines of easy and abundant food production and the most progressive nations have been the best fed. No people living in a hand-to-mouth fashion have lifted themselves above the poverty of their surroundings; no man can be an efficient member of society whose life is an alternate feast and famine. That is why the savage, ignorant of methods for conserving his food supply, is still a savage.

The food supply is perishable. Fruits and vegetables are seasonable; that is, for the most part suitable for use only during the months in which they reach maturity. Meats cannot be kept after slaughter except by special treatment; even the cereals deteriorate with age and the store is depleted by vermin. And so we have a season of plenty when food fresh from the fields and orchards gluts the markets and later seasons of scarcity when natural causes have destroyed the surplus of earlier months. These seasonal variations in the food supply are also subject to yearly fluctuations, for the abundant crop of one year may be succeeded by the scant crop of another. The fact that foods are perishable makes it necessary, if they

are not to be wasted, to supply some adequate means for holding in check the processes of decay which, if allowed to operate, would make them unfit for use.

Cold storage is the modern way for arresting food spoilage. It is the latest and most successful method of storing the surplus of one season against the want of the next, and of preventing the fluctuation of prices from below the cost of production at harvest to a point beyond the resources of the purse the rest of the year.

It is of especial interest to the health officer, both because of this phase, which, in so far as it affects the available food supply, touches the great problem of nutrition, and because of the general impression that goods held beyond what may be termed a natural period of usefulness are not suitable for food. Whether food deteriorates and to what extent should be understood by him in order that he properly may draft and enforce cold storage laws. During the last few years extensive investigations have been made to determine the deterioration of food in cold storage. The results of these investigations are the more interesting because, in some instances, at least, they upset generally accepted theories. Ever since cold storage has been practiced, cold storage chicken has been viewed with askance by the public, and cold storage food has been held accountable for every unexplained illness.

The flood of ill-designed, crudely drawn bills presented to the lawmakers of the various States during the last few years was without doubt a well-intentioned attempt to meet the demand for a careful regulation of the business of cold storage, both with the idea of protecting the health of the consumer, and, in some little-understood way, reducing the cost of living. In effect, however, the passing of many of the bills suggested would have meant the destruction of a most important industry. Yet the value of cold storage is clear to everyone who has given intelligent study to the subject. Even where cold storage facilities are not available, the necessity for them is recognized, and in Canada, at least, the government, appreciating the need of cold storage plants, has adopted the policy of subsidizing the construction of refrigerating warehouses. A committee appointed by the French government to study the recent increase in the prices of food stuffs has pointed out that this is in no small measure due to the fact that France has as yet practically no system of holding food stuffs in cold storage. Unfortunately, men who should be thoroughly famil-

iar with the practice of cold storage still appreciate neither its purpose nor its effect. The Commission on Cold Storage appointed by the Governor of Massachusetts to investigate the subject, addressed a circular of inquiry to secretaries of Boards of Health in the different States, asking for opinions or suggestions as to the need of regulation of the industry and the form which it should take. Twenty-one state boards answered the inquiry and in every instance recorded a belief that legislation for the regulation of cold storage of food and food products is necessary. That the information, however, was not on the whole of great value was shown by the fact that one official recommended restricting the time limit of storage to ten days, another to ninety days, several did not think storage for more than three months desirable. When the information of those we count as sanitary experts is so limited, need we wonder at the fear of cold storage products so long held by the average consumer? Before satisfactory legislation is enacted we must know why we need regulation and what, if any, bounds of restriction are necessary. The business should be regulated by practical laws which do not have for their purpose its destruction and which are intended rather to put a stop to the practice of the storing of food unsuitable for refrigeration, and which has, even before its entry into storage, deteriorated and become unfit for food, and to insure the withdrawal of all goods before they have been held sufficiently long to undergo such physical change as may render them undesirable for human consumption.

The report of the Massachusetts Commission referred to recognizes in cold storage a fundamental necessity in the distribution of the food supply of the nation. It finds that cold storage enables perishable food products to be brought to market with the least possible deterioration, and that it enables the surplus of one season to be carried over to meet the demand during the season of natural scarcity. In this way, by distributing the seasonal output of perishable food stuffs evenly through the market year, it helps to equalize supply and demand. The price of the food supply to the consumer is not materially influenced by cold storage. It has been argued that the possibility of storing food products against a rising market may lead to speculation on the part of the middleman, and no doubt the facilities offered by cold storage may be abused to manipulate prices. This possibility, however, is more theoretical than actual, because of the enormous practical difficulties in the way of artificially controlling the supply of food. It is impossible to determine in advance, for instance, whether

January and February will be relatively warm months as during the winter of 1911, or bitter cold months as during the winter of 1912. In 1911 the warm months brought about a very large production of eggs, and consequently eggs in storage were taken out at a loss to the owners. The possibility of such conditions obtaining acts as a deterrent to the speculator, and all data at hand shows that the manipulation of food prices is not materially increased by the practice of cold storage.

As above suggested, there have been desultory attempts to regulate the cold storage industry by legislation. The government of the United States, although it has discussed the enactment of such legislation for several years, has as yet taken no action. Several States, however, have enacted cold storage laws of varying character. The first cold storage law of record, in the United States at least, was enacted by the State of Indiana in 1911, similar legislation following in the States of New York and New Jersey in the same year. In 1912 the National Association of Food Officials gave to a committee the task of drafting a model cold storage bill. After many months of careful work and investigation and after the revision of several tentative drafts, the committee recommended as a model bill for enactment in the several States a draft which during the legislative sessions of 1913 was enacted in approximately its original form as a law in the states of California, Iowa, Nebraska and North Dakota, and by authority conferred upon it adopted by the Louisiana State Board of Health as the law for that State. In 1912 the Massachusetts legislature enacted a cold storage law, drafted after a most comprehensive investigation of the subject by a committee of the legislature appointed for that purpose. The latest law at the time of writing is that enacted in the state of Pennsylvania. The Pennsylvania law differs in several points from the model bill and indeed from the early legislation upon the subject, which will be referred to in detail later.

At the present time eleven States regulate the cold storage industry by law. The State of Kansas regulates the storing of certain food products, but has no general law. The Canadian government, appreciating the necessity for developing a cold storage industry, in 1907, passed a cold storage act entitled, "An act to encourage the establishment of cold storage warehouses for the preservation of food products." This act, while primarily not intended as a regulative measure, but rather drafted for the purpose of subsidizing the construction of warehouses, is in effect regulative in that

the application for subsidy is limited to warehouses which are constructed under plans subject to the approval of the Department of Agriculture.

Cold storage is defined as the holding of food products at or below the temperature of 40° F. in warehouses refrigerated for that purpose. A cold storage or refrigerating warehouse is held to be an establishment employing refrigerating machinery or ice for the purpose of refrigeration in which articles of food are stored for thirty days or more at a temperature of 40° F. or below. This provision varies somewhat in the several States. The State of Nebraska, for instance, requires that goods must be held in storage for sixty days before being legally cold-stored, while a bill pending in the State of Connecticut holds that eggs must be labeled "Cold Storage" if held for more than fifteen days. The time limit imposed by most of the laws is the natural limit of twelve months, that is, from one productive season to the next. The time limit, however, is not uniform in the several States. The State of Pennsylvania fixes a different limit for different articles of food. It limits the storage of whole carcasses of beef or parts thereof to four months, whole carcasses of pork or parts thereof, of sheep or parts thereof and of lamb or parts thereof to six months, the whole carcasses of veal or parts thereof to three months, of dressed fowl drawn to five months, of dressed fowl undrawn, ten months, eggs eight months, butter nine months and fish nine months. As a rule the law requires that goods which have been in cold storage shall be sold under a label advising the purchaser of their character. The Pennsylvania law even goes so far as to require that food sold from labeled containers must be wrapped in a package stamped on the outside with the words, "Wholesome Cold Storage Food." The Massachusetts, Iowa, Louisiana, Nebraska and North Dakota laws require the display of a sign marked "Cold Storage Goods Sold Here." The Indiana law requires only that eggs taken from cold storage be sold from a receptacle bearing the words, "Cold Storage." All laws are uniform in requiring that goods be marked with the date of entry into storage and the date of withdrawal therefrom, except that the laws of New Jersey and Delaware require only the marking with date of entry and the Nebraska law does not require the date of withdrawal on goods to be shipped outside the State. In nearly every case the warehouseman is required to report the quantity of goods in storage to the proper officials at the end of each three months' period. The Massachusetts law, however, requires the report but three times a year. This provision, while not in any way affecting the character of the goods in storage, is undoubtedly an attempt on the part of

the legislatures to minimize the possibility of the cornering of the food supply by giving all information concerning stocks on hand to the public. As an additional protection under certain conditions the officials of several of the states are authorized to call for more frequent reports than are specifically authorized in the statutes.

In six of the eleven States enforcing a cold storage law, the State Board of Health and its executives and inspectors are charged with the enforcement of the act. In five States the work is done under the supervision of the food commissioner or dairy and food commissioner, as the case may be. In every case, except the State of Delaware, it is made the duty of the official or executive board to issue licenses for the operation of cold storage plants. These licenses are issued after an inspection has shown them to be sanitary and properly equipped and operated, and the board or officials charged with the enforcement of the act have power to withdraw the license if the plant becomes unsanitary or is operated in violation of the law. An important provision of practically every law is that authorizing the officials to extend the time of storage if inspection at the end of the storage period shows the goods still to be in satisfactory condition and suitable for use as food.

Unquestionably the public has the impression that prices are artificially and arbitrarily raised by reason of withholding goods from market in storage warehouses. The special committee of the Chicago Association of Commerce, which made a thorough study of cold storage in its many phases, says of this argument against storage:

“Exhaustive examination of the statistics compiled under the directions of your committee, and a comparison of these statistics with the facts obtained by the department of agriculture, after an exhaustive research demonstrates clearly that the prices of butter, eggs, poultry and fish have been more uniform during the year since cold storage has become a factor in the care of food products than before that period. These statistics also show that taking an average for a period of years, prices on the whole have been lower than during the years when cold storage was unknown.”

This statement is in substantial agreement with the conclusions reached by the Massachusetts committee and undoubtedly is an accurate gauge of the effect of cold storage upon the price of food. Nevertheless, in view of the persistent criticism of the new industry and of the too general impres-

sion that high prices are the result of manipulation somewhere between the farm and the consumer rather than a decreasing supply for an increasing demand, the legislation enacted may be assumed to have definite value both to the warehouseman and to the consumer in that on the one hand the consumer knows where and how much goods are being held for future use and the warehouseman is protected from a criticism which, if persistently indulged in, must prove a serious injury to his business. This statement may, indeed, be applied to all the phases of cold storage legislation, and where the laws have been in force the longest, I believe that without question the industry receives most credit from the consumer, and cold storage food properly handled in storage and sold under an open label out of storage is not only viewed without suspicion, but indeed purchased and consumed with greater satisfaction.

The cold storage industry is not a local business, but is very largely a feature of interstate commerce. Public warehouses could not be maintained for the convenience of local trade. They must depend upon the large shipments collected in one part of the country to be distributed at centers of population. For this reason legislation affecting the industry should properly originate at Washington instead of as at present in the several States. It is perhaps unfortunate that the Federal government did not point the way to uniform and reasonable state legislation by itself enacting a fair and equitable law. The bills proposed for enactment by Congress have, however, been framed without a proper understanding of the subject, and for that reason have not met the favor of those engaged in the industry, the states' officials charged with the regulation of the food supply, the retail trade dependent so largely for a supply upon the warehouse, or the consumer, who wishes only to be protected against unfit food, manipulated prices and deception.

The regulations drafted by officials charged with the enforcement of the laws have been generous and pertinent. The laws have been construed liberally and with regard for the warehouseman. In general, goods held at low temperature in process of manufacture, such as beer and meats in cure, have not been held to be in storage. The technical features of the stamping and tagging have been made as simple as possible and in practice the dating of the time of entry and withdrawal is easily and economically done. There is still some dispute as to whether the small dealer, as for instance the butcher, who may carry small stocks of meats longer than the usual thirty-day period, and the hotel and restaurant, should be held to be

operating cold storage or refrigerating warehouses. In so far as storage may affect the quality of food stuffs there is no difference between the large public warehouse and the private ice box, except that in all probability goods cannot be handled as successfully at the smaller plant. However, the stock of goods held at the hotel or butcher shop for local consumption is never so great as to influence the market, and for that reason the generally recognized necessity for the publication of storage holdings does not obtain. Moreover, unless legislation presumes to label cold storage goods all the way from the warehouse to the consumer's table, there is no necessity in the case of the individual plant for the system of marking followed by the warehouseman. Goods taken from storage are sent to the hotel kitchen or to the home of the consumer without delay, and deterioration is avoided, as might not be the case with the careless handling of goods drawn from cold storage for distribution over a larger area.

Recognizing a strong sentiment for cold storage regulation and the fact that such legislation is already in force, not only in Western States where no warehouses are in operation, but in the populous Eastern States of Massachusetts, New York, New Jersey and Pennsylvania, it behooves the industry to demand adequate protection by federal legislation, protection against unwise state legislation, protection against the loudly expressed yet admittedly erroneous statement that the cold storage industry is employed to manipulate prices to the detriment of the consumer, protection against the firmly established impression that goods deteriorate markedly in storage, protection against the oft-repeated tale that food-poisoning follows the ingestion of cold stored goods. Legislation that accomplishes these facts will not operate to curb the development of the industry, but rather to stabilize and encourage the use of refrigeration by the producer and of cold stored foods by every consumer.

With the passage of adequate cold storage legislation and the development of a practice of labelling which declares the character of the goods to the purchaser, the idea now held that cold storage is an artifice used by the speculator to force higher prices and a practice which spoils food instead of preserving it will no longer obtain.

And when cold storage is no longer feared, our markets will be widened and the food supply enlarged by the thousands and hundreds of thousands of tons of edible products which now rot on the ground for want of facilities to preserve them to such a time that they can find a profitable market.

CHANGING CONDITIONS IN THE KENTUCKY MOUNTAINS.

B. H. SCHOCKEL.

(Illustrations by the author.)

This summary of changing conditions in the plateau of eastern Kentucky is based upon a month's field work, supplemented by previous and subsequent studies. To refresh the reader's general conception of the region an introductory review is made of its topography, surroundings, and settlement.


TOPOGRAPHY AND SURROUNDINGS.

Eastern Kentucky is a part of the Cumberland Plateau and consists of thirty-five counties with an area of some 12,943 square miles, that is, about one-third of Kentucky. It is a part of the Southern Appalachian Highlands. To the east of it are the parallel ridges and valleys of the Greater Valley of the Appalachians; to the west is the Blue Grass region. The top of the plateau is a part of the Cretaceous Peneplain, with monadnocks on it, and slopes gently westward in Kentucky from an elevation of about 2,000 feet to a height of 1,200 to 1,500 feet. This peneplain has been dissected by dendritic drainage to a topographic stage of maturity, the valleys being from 500 to 800 feet deep with narrow bottom lands, and the tops of the ridges averaging in many instances from 10 to 50 feet in width. The ridges, locally known as mountains, in general bear on their shoulders and crests hardwood forests sprinkled with conifers. Most of the lower slopes are cleared. From the top of Pine Mountain the Kentucky country appears to be a billowy wilderness. One cannot see any valleys, nor any sign of life; but beneath those forested waves are sylvan slopes to enchant one, and a sinister labyrinth of gashed valleys to enthrall one in mountain poverty.

Owing to the topography the roads are serpentine; since the bed rock is of shale and friable sandstone chiefly, good road material is scarce; furthermore, the people are poor, and what we term shiftless and ignorant; therefore, their highways are in a most wretched condition.

SETTLEMENT.

In the sixteenth century James I introduced Scotch settlers into northern Ireland, who became the Scotch-Irish. Some of them emigrated to America; and their descendants, augmented by English, native Irish, Pennsylvania Dutch, and others, formed the van of the 300,000 frontiersmen who passed through Cumberland Gap, from 1775-1800, to settle in Kentucky.




1. Creek-road, "upright farm," and forested ridge, near Pine Mountain Postoffice, Ky.

Some of these found a home in the plateau region, which offered clear springs, magnificent forests, abundant game, and good valley land sufficient for that first generation of hunter-farmers. No one could have foretold then the coming of canal and railroad.

The first permanent settlements in the Kentucky mountains were made in the decade 1790 to 1800. Inlay's map of Kentucky (1793), shows "settlements" on Rockcastle River, the upper Louisa Fork, and a fork of Red River. By 1800 the population was 7,964, which was about four per cent. of the population of the State; it is now about 600,000, which is about twenty-five per cent. of Kentucky. Genealogical records of this people are utterly lacking. Their names and survivals in customs and language point to English and Scotch-Irish ancestry in general, although a few German and Huguenot names are found.

Between 1800 and 1840 the mountain region was an integral part of the State, for various reasons. Four interstate, transmontaine routes trav-



2. An example of the poorest highways in the mountains, near Pine Mountain Postoffice, Ky.

ersed the plateau in leading from the Ohio and the Blue Grass countries on the west to the Big Sandy and Kanawha region on the east, and thus on to the tide water settlements. The plainsmen bought lean cattle in the Blue Grass and sent them in droves of from 200 to 300 through the mountains to the Potomac, where they were fattened and sold in Baltimore and Philadelphia. Large droves of hogs followed the same routes. The hog and cattle drivers bought corn at the homes of the mountain people and brought news from the outside world. The slender state appropriations for roads were impartial, the mountain counties being favored equally with the lowland.

But between 1830 and 1850 the four interstate roads declined gradually to a wretched condition and state of non-use; for the Blue Grass and Ohio regions were finding other routes to market, by use of steamboats, etc. Therefore, the mountain counties lost their market and received little outside help for roads. As a result the people have lived isolated by topography and social antipathy.

During the Civil War thousands of the mountaineers, whose ancestors had fought in the Revolution and the War of 1812, joined the Union army and received a practical education. Some received similar training as soldiers of the South. After the war many returned home. But the growth of formal education and broader outlook thus stimulated has been slow.

In 1878, Shaler, of the Kentucky Geological Survey, saw in the eastern, and then most inaccessible portion of the region, men hunting squirrels and rabbits with old English "short-bows" and wrote: "These were not the contrivances of boys of today but were made and strung, and the arrows hefted, in the ancient manner. The men, some of them old, were admirably skilled in their use; they assured me that, like their fathers before them, they had ever used the bow and arrow for small game, reserving the costly ammunition of the rifle for the deer and bear."

Recently outside capital has begun to develop the coal and timber resources of the region, a fact which is bringing about many changes in the mountain country rapidly. As a result, the inhabitants are facing the crisis brought about by the sudden mingling of a primitive people with the exploitative phase of modern civilization.


CHANGING CONDITIONS.

Mineral Resources.

Coal is the chief mineral resource of the region. The seams occur in every county, increasing in number and thickness towards the southeast and reaching their climax in the Black Mountain region. The layers are favorably disposed for mining, except in the Pine and Cumberland mountains, where complex structure renders mining difficult. The coal is bituminous, the most desirable varieties being as follows: Cannel, found in limited basins throughout the field; coking, appearing in large amounts only in the vicinity of Pound Gap; and high class steaming coals, occurring in quantity in the southeastern counties and at a few places along the western margin.

Coal was exported in 1827, probably earlier; but until the railroad came, the output was insignificant. Though production is rather small at the present time, and limited to a few mines scattered along the railroads, the region is beginning to become an important coal center.

The first extensive exploitation began in the region about Middlesboro, in 1892. At present most of this coal is shipped south. Some two years



3. A primitive mountain home, near Buckhorn, Ky.

ago a branch of the L. & N. railroad was pushed up the North Fork Kentucky River to Hazard, and extensive coal mining began. Hazard is now in its Ugly Duckling stage, has a population of about 2,000, boasts one of 3,500, and altogether is a scar upon the beautiful landscape, like a "boom"

town of the west. But the most spectacular development is taking place at Jenkins, on the headquarters of the Elkorn Creek, at the foot of Pound Gap, Pine Mountain, known in literature as "The Trail of the Lonesome Pine." Eighteen months ago a branch of the B. & O. Railroad reached the site, where a few months prior there had been but one mountain cabin. Jenkins now has brick buildings three stories high; a great power plant; palatial residences; a splendid hospital; a concrete dam causing an artificial lake, upon which are pleasure boats; and a town reservoir, into which spring water is filtered from the mountain. Indeed it is growing as fast as Gary, Indiana, in its early days. Most of this coal is shipped across Indiana to Gary.

At a shaft mined by two mountaineers near Booneville, good cannel coal sells for seven cents per bushel. Cause, poor transportation.

Essentially all of the mineral rights have been bought by outside capital, much for \$1.50 per acre, and in some cases for fifty cents. Sometimes the mountain people made the further mistake of giving up the farming rights also.

At an early period iron and salt within the region were the source of considerable traffic, but not now. Oil, gas, and clays, although in progress of exploitation for the last two decades, do not promise to become important.

Forest Resources.

The primitive forests were splendid. But since an early day, lumber has been shipped to an outside market; therefore, the timber area has been reduced, and, although it remains the chief source of wealth, the end is almost in sight. About thirty per cent. of the region was in wood in 1910, not all of which was primitive.

The mountaineer's way of lumbering is to cut a few choice trees and "snake" them down to the creek, where as logs, rafts, or railroad ties, they await the coming of the flood, or "tide", to be floated down stream. Thus a man can produce ten ties per day, for which he received thirty-eight cents apiece, this summer, near Beattyville. But lumbering corporations are beginning to attack the two remote corners of the Southern Appalachian Highlands—the Smoky Mountains and the Kentucky Plateau—and after the onslaught, in which stumps three and one-half feet in diameter are left to rot, the hills are gaunt with slash, or black from resultant forest fires. Consequently increased erosion is resulting on the slopes, with augmented

harmful deposition on the flats below. Furthermore, within and below the plateau, the streams increasingly are characterized by short periods of flood, and long intervals of low stage. Also, the supply of drinking water and water power is becoming less constant.

Most of the timber is owned by outside capital. The United States government is seeking to buy wooded land for forestry having completed the first step last September, when it purchased the 60,000-acre Biltmore estate, near Asheville, N. C.

A passing forest industry is the digging of ginseng and other roots. The normal market price for the first is five dollars to seven dollars per



4. Snaking logs in the Southern Appalachian Highlands.

pound, but now, owing to the war, the price has fallen. One old man and his wife, digging "sang" in the woods, wanted to know why it is that the Chinese cannot live without the root, and what would happen to that people when the supply shortly would give out in America, but then consoled themselves with the thought that probably the Chinese have enormous amounts of it stored away in anticipation.

Animal Resources.

Wild game is becoming surprisingly scarce, due to overhunting, and lax observation of hunting laws. In general the supply of fish is low, some causes being: Dynamiting and seining, lack of restocking, and inconstant

and turbid streams resulting from deforestation. What little meat is eaten, is chiefly swine and chicken. Sheep continue to suffer from exposure and dogs. Beef is walked to market to obtain cash with which to pay taxes. The Ayrshire stock is giving way to a fine short-horn type of cattle, owing to the opening of stock yard cattle markets, as at Mt. Sterling. The mountain mule and pony are being displaced by larger types. Goats suffer from the rough, wet, winter climate.

The development of pasturage for live stock would prove to be a fundamental advantage. Timothy is the chief forage crop; clover is second. A diminutive Japanese clover has filtered into the mountains, and takes possession of deserted fields. It is good for grazing, but it is too small to be cut.

Agriculture.

About 80 per cent. of the land is in farms, of which 45 per cent. is improved, and 23.5 per cent. in woodland. The average size of the farm is 85.7 acres, of which about thirty-nine acres are improved (Kentucky: 85.6; 55.4—Indiana: 98.9; 78.6). The average value of all crops per farm in 1910 was \$310.70. (Kentucky: \$536.20—Indiana: \$947.60). The average value of implements and machinery per farm in 1910 was \$32.3. (Kentucky: \$80—Indiana: \$190). About 6.6 cents worth of fertilizer was used per improved farm acre in 1909. (Kentucky: 8.7—Indiana: 12.8).

The total value of all crops in 1909 was 24.8 million dollars, of which cereals amounted to 12.2 million, vegetables 3.8, hay and forage 1.1, and fruits and nuts 1.1. The total area in cereals was 921,538 acres, of which corn constituted 841,744 acres; oats, 39,341; wheat, 36,403; ryè, 1,579; and barley, 510. Some 21,397 acres were devoted to potatoes, 5,673 to sweet potatoes and yams, and 10,713 to edible beans (a staple food in the mountains). Sorghum was raised on 21,970 acres, and hay and forage on 162,944 acres. There were 1,825,895 apple trees out of a total of 2,425,047 fruit trees. Peaches ranked second to apples.

The average production of corn per acre in 1909 in the region was 18.7 bushels; in Kentucky, 24.2; in Indiana, 40. The corresponding figures for wheat were 9.9; 12.8; and 16.3. Similar data for potatoes were 76.6; 91.8; and 99.4. The respective figures in tons of forage per acre were .8; .9; and 1.2.

The shale soil, which is most common, is fairly fertile, and produces good crops of corn under good cultivation, on gentle slopes. The chief

causes for the low productivity are steep slopes, poor cultivation, and lack of crop rotation. The shale soil washes less than almost any other soil under like circumstances. The wonder is that the soil produces as much as it does.

A few years ago Berea College, with the help of the United States government, employed a special investigator and demonstrator to work with the mountain farmers within reach of Berea. The success was such that a number have been appointed in other localities. About Berea, heavy breaking plows are replacing the one-mule plow, and the disk harrow is pushing back into the mountains. More than twice as many shallow culti-



5. Stumpage and slash which will invite forest fires in the Southern Appalachian Highlands.

vators as single shovel and double shovel ploughs were sold in Berea last spring. The practice of sowing cow peas and rye for forage and turning under is spreading, as is the use of commercial fertilizer. Crop rotation is displacing the fallow system.

Further education in agriculture is being given at the missionary and settlement schools, as at Oneida, Hindman, Buckhorn, and Blackie. But agriculture in the interior of the region is yet primitive, and improvements are slow in penetrating. A common sight is corn growing among girdled trees.

The few gardens which are being introduced about the settlements and mining and lumbering camps are giving favorable results.

Naturally, the region is a splendid fruit country, especially for apples; but spraying is unknown, and the stock has degenerated. Therefore the trees bear abundant crops of gnarled, sour fruit. One mountain woman told us to take as many apples as we wished, since they were of no value except to sharpen the teeth on. Often apples are sold for ten cents per bushel, are given away, or rot; cause, poor transportation.

Manufacture.

Manufacturing within the region always has been meagre, primitive, and for local use, except in the case of salt in the early days.

In 1901 Bell and Boyd counties contained 172 manufacturing establishments, with an aggregate capital of \$5,201,489, an amount which was more than one-half of that invested in manufactures in all the thirty-five counties in 1910. The cause for the emergence of these two counties is the recent growth of Ashland and Catlettsburg on the Ohio River, and Middlesboro near Cumberland Gap, a local supply of coal being the factor in each case. Hazard and Jenkins soon will rank as manufacturing cities.

The status of manufacturing for 1900 is indicated in the following table:

	Establish-ments.	Capital Per Establish-ment.	Men, 16-Years and Over.	Women, 16 Years and Under.	Children Under 16.	Capital.	Value of Products.
Kentucky Mountains..	1,156	\$7,221	4,853	44	85	\$8,347,993	\$11,993,195
Kentucky.....	9,560	10,886	51,101	9,174	2,687	104,070,791	154,166 365

The mills are small and are driven by water, animal, and hand power. Machine made goods from the outside have supplanted the linsey-woolsey cloth, counterpanes, and baskets formerly made in the cabins. But, recently, the missionary and settlement schools have begun to sell such goods outside of the mountains for the people, to supply cash, and therefore the industries are reviving, in part. The W. C. T. U. Settlement School at Hindman, for example, sold \$1,800 worth of such goods last year.

Distilling always has been a widespread industry in the mountains, since thereby corn, the chief crop, is converted into a product which can be marketed with profit, and since the custom has been inherited. Illicit distilling increased greatly after the imposition of the liquor tax of the Civil

War. In 1877 the government began to suppress "moonshining" in the region. By 1882 the supremacy of the law had been established. But in 1894 the liquor tax was increased from ninety cents to one dollar and ten cents, which resulted in increased "moonshining". The counties have been voted "dry", which encourages the illicit traffic. About the coal mining centers, "blockading" is increasing greatly, the whiskey being brought to town under vegetables and in milk cans.

Transportation.

Transportation is the basic problem of the region. Poor communication within it has influenced greatly every phase of life always, and bad connections with the outside have isolated the country since 1850.



6. A primitive mill, near Cornettsville, Ky.

Of a total of 17,432 miles of road, there were within the entire region, in 1904, eighty-three miles surfaced with stone, and four miles with gravel. The present wagon freight is said to be about 44 cents per ton-mile. The average haul for a load of cross-ties is from eight to ten miles, and eight to twelve ties constitute a load.

Logs delivered at the railroad for twenty dollars per load are said to consume sixteen dollars in transportation. From Buckhorn to the railroad is eight miles. A team will make this trip for four dollars in good weather.

The charges in this case are about 88 cents per ton-mile. The average cost of transportation in the United States by wagon is 23 cents per ton-mile.

The old law that every man must work on the roads six days annually is enforced feebly. By a statute passed in 1894, road taxes can be levied by the county and a road commissioner appointed. But this new law is proving a failure in the mountains and is giving way to the old custom because the mountain county is too poor to pay the commissioner's salary, and because the mountain man may pay the tax in work, a fact which introduces again the old problem of road-work enforcement. In 1904 the total expenditures upon the highways in a number of rugged mountain counties amounted to about \$24 per mile. The average expenditure for the State, much less dissected, was \$43.57. The history of the mountain roads emphasizes the inability of the people to provide themselves with efficient highways, and manifests the great need for outside help, state or federal. In general, road material would have to be imported at great expense. The costs of roads steadily increase as the forest retreats towards the headwaters.

In 1907 the United States Department of Public Roads, as an object lesson, built and macadamized in Johnson County, 5,780 feet of road, and constructed through Cumberland Gap, 12,300 feet of macadam pike, and graded 900 feet more, at a total cost of \$7,050 per mile. This work demonstrates again that the construction of good highways in the mountain region, while possible, cannot be done without outside help. Besides the government routes there is a short stretch of macadam road (1 to 20 miles) in five marginal counties, of which, however, Boyd County alone lies strictly within the mountain region. The coal company at Jenkins has surveyed and built six miles of well-graded dirt road connecting Jenkins and McRoberts. Owing to the enforcement of the road laws in Knott County, a fairly good ungraded dirt road extends thirty miles between Hazard and Hindman. Immediately west of Pine Mountain in Leslie County, no wagon roads were attempted till 1890, and few exist now,

Before the advent of railroads, highway improvements were negligible, but the past twenty years have seen progress. Numerous stretches of road, eight to ten miles in length, afford somewhat fair transportation for wagons to the railroads. Where the development of coal and timber has increased the wealth of the community greatly, substantial bridges have been built. Progress has been slowest in the rugged, extreme southeastern

section of the region, even though railroads have begun to penetrate. There the primitive saddle and sleds drawn by oxen are still in use.

Except for lumbering, the streams are used but little. The North, Middle and South forks of the Kentucky River penetrate into the interior. They join at Three Forks, near Beattyville. Thence to Carrollton are 350 miles of good waterway. In 1853 some five locks were completed by Kentucky at a cost of \$4,000,000, which assured good navigation for 300-ton steamers for a distance of over 100 miles. The Federal Government made improvements at the close of the Civil War. Since then the waterways have been declining. In 1887 there were passing Three Forks annually, 50,000,000 feet of lumber, in logs.

Railroad building began in 1856, but made no headway until between 1870-90. The progress has been slow and confined to marginal counties until recently. Within the past five years it has penetrated the North Fork Kentucky River to McRoberts, a few miles west of Pine Mountain, and up the Poor Fork of the Cumberland River, by way of the gap at Pineville. The railroads have been built for the coal and lumber, and not primarily for general traffic. Since the advent of railroads, the conditions which have made possible "the mountaineer" have been passing away.

But in general the region is still landlocked.

Population.

In 1910 the total mountain population was 561,881, representing an increase of 18 per cent. over that of 1900. (Kentucky: 2,289,905; 6.6 per cent.) There was an average of 43.4 people per square mile (Kentucky: 56:9; Indiana: 74:9). The density is greatest along the main river routes and in mining sections. The people continue to be distributed as clans in valleys, which are surprisingly heavily populated. Of necessity the people depend upon the lower slopes of the hills to an extent equal to or greater than on the limited bottom lands, their "shoe-string farms" being found strung along little gullies as well as in broader valleys. A few farms are on the mountain sides, especially on benches or "coves" of somewhat gently sloping land, formed above some massive sandstone ledge. The average size of the mountain family is about 5.2. (Kentucky: 4.6; Indiana: 4.1.) The rural population increased 17.1 per cent. in the last decade. There was no urban population (towns of 2,500 or more) in 1870. In succeeding decades, as Ashland and Middlesboro developed as centers of coal mining, it numbered 3,280; 7,466; 17,428; and 24,004. These two

cities are unique in the region in having a population greater than 5,000; but they soon will be joined by Jenkins and Hazard, about which coal mining is developing rapidly. In 1910 less than one-half of one per cent. of the total population was foreign born. These people were chiefly skilled miners from England, Sweden, Germany, and Switzerland, who drifted in by way of Pennsylvania. In seven counties there were no farmers of foreign birth; and in only one county did the foreign born exceed 21. Recently, Southern Europeans have begun to come, particularly Italians and Hungarians. By 1920 the number of foreign born will have increased greatly. In 1900 about two per cent. of the population were negro, and in 1910 two and one-half per cent. In three counties there were no negroes; and in sixteen, less than twenty.

The problem presented in the region by the rapid increase in population with no corresponding increase in foodstuffs, probably is not greatly overdrawn in the following statements by a mountain graduate of Berea College: "The pioneer of 1850 who sat in his front door watching the deer rove the unbroken forest, today sitting in the same place can see acres of spoiled farm land. A few years ago the people produced enough on their farms to support themselves. Today one-half of the food consumed is brought in by the merchants. Twenty-five years ago our hillsides produced forty bushels of corn per acre. Today the average yield of corn per acre is a little less than twenty-five bushels. (In 1909 it was 18.7.) The independent farmer of yesterday has been transformed in the last few years to a man dependent upon his staves and ties for support. Now, his farm has grown up in bushes, and his timber is almost exhausted. . . . Such is the condition of a vast number of our mountain farmers."

There is an emigration of the mountain families, or of sons and daughters, particularly from the marginal counties, where a fringe of mountain territory has been put in touch with outside progress and humanity, and where mountain peoples are buying adjacent lowlands. Some are moving to Oklahoma and the Far West. This in part accounts for a decrease in population of five counties.

Public health is not as good as might be expected at first thought. The situation has been summarized by Miss Verhoeff (in "The Kentucky Mountains") as follows: "Endurance and muscular strength are common, but a strong constitution is exceptional. Bad housing and sanitation, ill-cooked and insufficient food, exposure to weather, and . . . poverty, have

had their detrimental effects, which have been augmented by a close inter-marriage of families and by an inordinately large use of liquor."

In general the mountain man is quicker than the Indiana plainsman, but not as strong. A month's field work did not bring to note any of the storied giants of the hills, though there probably are some. Not all of the people are lank.

About two generations ago trachoma penetrated into the mountains, and is spreading rapidly, despite the efforts of the state and settlement schools, and the Federal Government. Of over 4,000 people examined in five counties, 12.5 per cent. had this disease. A report from the W. C. T. U. Settlement School at Hindman names twenty-five per cent. for that locality. Adenoid and turbinate cases are common. Several clinics held at Buckhorn revealed that 90 per cent. of those examined were afflicted with hookworm. Splendid work is being done, but the area to be covered is a vast one, and assistance is needed greatly. Superstitions that diseases are visitations of the Lord to be borne with resignation are disappearing slowly.

The people continue to be poor. In 1900 land was worth \$5.00 per acre, and in 1910 \$9.66. (Kentucky: \$13.24 and \$21.83; Indiana: \$31.76 and \$62.36.) The average value of all farm property per farm in 1900 was \$860; and in 1910 it was \$1,359. (Kentucky: \$2,007; and \$2,986; Indiana: \$4,410 and \$8,396.) The average value of farm buildings per farm in 1910 was \$247. (Indiana: \$1,230.)

Institutions.

There is great need of education. In 1900, 24.3 per cent. of the voters were illiterate, and a decade later, 20.7 per cent. (Kentucky: 15.3 and 13; Indiana: 5.6 and 4.1.) In eight counties in 1900, the illiterate voters constituted from 30.5 per cent. to 35.8 per cent. of the total. In 1910, 61.6 per cent. of the children, ages 6 years to 20, were in school. (Kentucky: 60.8; Indiana: 66.) Corresponding figures for children from 6 years to 14 years were 73. (Kentucky: 76.)

However, improvement is being made. In 1900, there were more than 20 counties without a local publication. Now, there are but few counties without a press, and several have more than one.

Formerly, the term of school lasted but three months in the year. The teachers received no training except in the common schools. The buildings were tiny, two or three teachers in some cases teaching in the same room.

But now, the term lasts six months (closing at Christmas owing to bad roads.) Also, many of the teachers receive some training in the normal department of the settlement and missionary schools. Furthermore, there is but one teacher in each room, though in it is no library, few modern desks, and little equipment. In one mountain school visited by the writer in 1914, the pupils were sitting in rough board pews, the boys on one side of the room and the girls on the other. The walls, floors, and seats were dirty. Some of the children wore but one garment. Two of them were suffering from trachoma. The equipment owned by the school consisted of one wall map and three calendars. The only object on the desk was a small switch. The girl-teacher, who was a graduate of the institute at Oneida, had charge of 69 pupils and, besides, without pay, was teaching a "moon-light" school of evenings, to which people of all ages were coming. She did not show any surprise or nervousness when our group of ten men in nailed boots filed in. Nor did the children pay much attention to the visitors. The third grade droned out its reading lesson, and then the second grade carried out its solemn program in spelling. There was a solemnity about it all which the outsider does not understand until he becomes acquainted with the gravity of these people in their gatherings. Progress was being made, though it seemed a pity that the children should have to learn the definition of some words which probably they never will have occasion to use. The day of the "shouting school" (in which the pupils indicate that they are studying by reading aloud) has passed in the mountains. In a second school, a girl, younger than the teacher above, was in charge. She had had no training beyond the common school. There were a few modern desks, but also some rough hewn pews. When I tipped to the door and took a photograph of the interior she showed less surprise than an Indiana school mistress would have exhibited, but she smiled when some of her children awakened to the situation. In some sections a holiday week is declared during the corn harvesting season. Mission and W. C. T. U. settlement schools are coming into the country, as at Buckhorn, Hindman, Pine Mountain Postoffice, and Blackie.

Berea College, on the western margin of the region, serves as a university for the mountains, and is sending its extension department with wagon and camp into the remote sections. The reader is referred to the December, 1912, number of *The American Magazine*, for the story of the heroic foundation of Oneida Baptist Institute, and is reminded of Bulletin Number 530 of the United States Bureau of Education for the story of the

opening of "moon-light" schools in the Kentucky mountains in 1911 for children, parents, and grandparents. When the feud breaks out, mountain mothers from the section in which blood is shed, anxious to get their sons out of danger, are wont to urge them to attend school at Berea College and elsewhere.

Though in some sections enthusiasm for education is becoming great, in others there is great apathy, because of lack of interest on the part of the people, lack of practical teaching, illiterate teachers, poverty, poor roads, and political interference in school affairs. In some districts it is still thought by the school trustee that "the lickinest teacher makes the



7. "The telephone whispers through the silent hills," near Booneville, Ky.

knowinest younguns". Changing conditions are indicated by an incident in which two teachers appeared in the same schoolroom, each determined to become the sole teacher. The following among the pupils was about equally divided at first, but presently they moved away from the teacher using the "A. B. C." method and grouped themselves about the more progressive instructor who was following the sentence method. The broad effort is being made to teach the people how to work and live according to modern ideas, and yet to retain the desirable traits of their own civilization. This is a delicate task, involving much more than merely academic training.

Religion is undergoing transition slowly. Formerly if a speaker did

not shout and gesticulate he might be termed a good *speaker* but not a good *preacher*. The early attitude towards the settlement workers was indicated in a mountain sermon in which the congregation was told to "beware of the fetched on women who come in here wearing gold watches, and their shirt fronts starched so slick that a fly would slip off and bust out his brains". But, a year later, the same mountaineer said that since these women were administering to the needy under conditions so harsh that even the mountain people would not venture out, "I allow as how they are welcome to stay in the mountains as long as I live." One mountain patriarch, who has given his farm and essentially his all in founding a settlement school in the valley of his home, gives some of his reasons as follows: That there was much whiskey and wickedness in the community where his grandchildren must be reared, was a serious thing for him to study about. He heard two of his neighbors say that there is neither heaven nor hell. One of them said that when a man is dead he is just the same as a dumb beast. Another said that he could not rear his large family of children to be as mean as he wished. The founder's idea was that a good school "would help moralize the country." Formerly the Presbyterian religion was most prevalent, but it gave way to the "Hardshell" Baptist creed, since in the mountains the educational qualifications for the latter were less severe than for the former. The disciples of this religion have in turn given way before the "Missionary Baptists." Methodists are also numerous. The most vivid disputes in the mountains were wont to be about religion. But now there is a significant change toward toleration in that preachers frequently exchange pulpits with pastors of other denominations, and that the use of a church is often tendered to another denomination which temporarily is without a place of worship. The following can be interpreted as a groan of growth: "The church in eour holler, hits about dade. Part ov the folks wants an eddicated preacher, an parts wants an old-timer, an so they don't get nary one". The funeral preaching had become the sole opportunity for social gathering until the recent advent of "camp meeting week", and the coming of the extension school on wheels.

Changing conditions have not yet affected greatly the political situation in the mountains. Since the Civil War so many of the inhabitants have been Republicans that party arguments have been one-sided, and the contests have been within the organization. Unity of feeling gives the representatives considerable power in the State Legislature. Political discussions are said to be confined in general to stump speeches con-

cerned with national issues, and hence are of little help concerning local problems. However, since the mountain men are good at politics, some make of the local contests a profitable business. Recently in some sections such men have turned their attention to the school, for the sake of profit in the appointment of teachers. There the trustee runs for office upon a platform statement of which teachers he favors. In some sections the vote runs high in school elections, while it is light on other matters. An increasing number of women vote on school affairs. Another favorite field of the politician is the handling of road taxes.

Deep seated prejudice, due to poverty, exists against taxation of any kind. In 1906 the per capita state and county tax was \$4.62 for Woodward County, in the Blue Grass, while in the mountains it ranged from \$0.40 in Elliott County to \$1.75 in Harlan. Little returns are obtained by taxation of lumber and mineral resources.

The feud was transplanted from Europe into the Blue Grass, the Kentucky mountains, and elsewhere. It survived among the isolated valleys of the mountains, where it was fostered by folk-song, the flaring resentment of the Indian fighter and pioneer, and the habits of thought natural in isolated communities where for a long time there was neither sheriff nor jury and where, even to this day, the government hardly has been able to inspire confidence or dread. The Civil War greatly increased and intensified the feud: Prior to 1860 few weapons had been used in the mountains, and few deaths had resulted. In the region in 1860 there were 10,098 slaves and 1,280 free colored people. The lines grew sharp between the Union and Confederate counties, as well as between opposing families, and between opposing members of a family. Modern arms were introduced into the region. The physiography of the land favored bushwhacking. During the war the Kentucky mountaineers suffered more sharply than the mountain people of any other State, except Tennessee. Also, many of the principals of the post-war feuds were boys during the Civil War, whose imaginations were filled with all of these horrors. It is said by the mountain people that the actual numbers engaged in the feuds has ranged from 10 to 60 on a side; that the duration has been from 1 to 40 years; that perhaps not 10 per cent. of the mountain people have had a personal difficulty sufficient to cause fighting; probably not 40 per cent. of them have gone to a court house to prosecute or defend a case; and that half of the enlisted partisans never have faced the

music in a show down fight, the number actually having figured in the ambuscades being small.

In some parts of the region, as about Oneida, education is causing the decline of the feud; but in other sections it flourishes, as near Pound Gap (Trail of the Lonesome Pine), near where, it is said, some eleven men were killed in three months during the spring of 1914.

The home, also, is changing. One still can see the windowless log cabin with its "dog-trot"; but some roofs are of shingles, and some of tin, while frame structures are appearing, and brick.

Mountain simplicity and hospitality are illustrated by one man who said, "I want a good house; two rooms—one for the family and one for company, each big enough for a bed in every corner—and a lean-to cook room."

The following is a description from Professor Penniman's unpublished tales of the mountains: "Three days are ample to build a log-cabin twenty feet square. The part before the roof is called a 'pole pen'. This is run up in a few hours. The trees sufficient to build a cabin complete are often standing on an acre. With the roof up, and stone chimney on the outside, and the big fireplace opening into the room, the young people can begin housekeeping. A few saplings will make a bed frame fastened to the logs in one corner, and a bed without a tick, two feet thick, of fresh pine needles, gives a sense of luxury to the newly married pair."

Customs and Habits.

It must be remembered that there are all grades of society in the mountains, and that no general description can be applied to a specific case.

Woman is inferior to man in both number and position. Not only is she a household drudge, but a field hand as well (out-of-door work in itself, of course, does not constitute drudgery). She still follows behind him as they trudge over the mountain. A mountain boy, upon being asked how many brothers he had, answered promptly: "Two." But concerning the number of sisters, he drawled: "Oh, three or four." The modern woman movement hardly has penetrated into the hills, and when it does, it will meet orthodox opposition. However, women increasingly vote in school affairs in some districts. Furthermore, here and there a girl returns from Berea, or some other college, with ideas strange to her people.

Perhaps this explains the wide girdle, or other bit of modern adornment, now seen sometimes on the quaint costumes.

We were pushing through a deep forest in toiling over a ridge. Before us were two children, walking in single file, a boy of fourteen and a girl a year younger. Our youthful guide pointed in their direction and remarked, "They were married last spring. Some of us do get married that early hereabouts; but we who have been to the settlement school don't calculate to get married that soon."

"Store clothes" have displaced the homespun garments, the result being unfavorable in the appearance of the men. However, the settlement schools are reviving the home-weaving industry to some extent. The belt is beginning to rival the suspender on "Sunday" garments.

The quaint old English language also is disappearing, though slowly. It is becoming crystallized and is losing its flexibility whereby it was wont to be bent this way or that, to suit the fancy or fit the occasion. In a reminiscence of his boyhood, Professor Dizney tells of a minister in Dizney's valley, who, in preaching about apostasy, took as his text: "If they shall fall away", and who concluded in a high key: "'If they shall fall away', means that they cannot fall away, for anybody who knows anything about the English language knows that it is a verb *in the impossible mood and everlasting tense.*" There also comes to mind the following expression: "Law me, Honey, I'm glad to be back from the plains. Wooded mountains make the restinest place to lay your eyes on."

There is about to pass away a most interesting folk-song based upon English and Scotch ballads, and preserved verbally in the mountains with slight modification, from generation to generation. These songs of romantic love, hate, sacrifice, and revenge are sung in almost all of the log cabins. Thereby the visitor, who may have thought that the mountaineers neither weep nor smile, learns with delight that their natures are intensely fluid. The songs are sung in slow time, and in minor tones difficult to express in written music. An effort is being made to collect the words and write the music before it becomes too late.

The open hospitality, once common, is shrinking. An old man in his watermelon patch put it thus: "I used to raise melons for the whole valley, so that the folks would come to sit and talk with me on the porch while we ate them. But now too many foreigners have come in; they

would eat me out of home." Sometimes the mountaineer is disappointed in his hospitality to strangers.

There is a kindly, affectionate courtesy for one another among the people, which, it is hoped, will survive.

There is such a great need for improvement in sanitation that what has taken place is negligible.

The native is accustomed to work in his fields by seasons, with periods of rest between. During an "off" period in September but two men were seen at work in the field during eleven days of travel. It has been his wont to work during the favorable time, or when the larder is empty; or to rest during the unfavorable season, or while provisions are at hand. Therefore, in general, the population is unsuited to the routine of work in the mines, the manufacturing plant, and the lumbering camps, now appearing in the region under the control of outside capital. Furthermore, it is without a disposition to coöperate. Hence such workers are at once the despair and menace of the employer and the labor union. Consequently, foreign labor is imported, and the mountain man is in the way, as was the Indian. He will not necessarily become happy if, to meet modern industrial conditions, he throws off lightly his old attitude toward life gained through centuries of adaptation in the mountains. A few of the most versatile natives are profiting by the rapid changes; but the great majority, formerly independent land-owning farmers, are not. Many are seeking employment in mill or mine, or are contracting to the headwaters. It is significant that the leaders in the mountains, native and mission, deplore the *rapid* advance of industry into the region, and that they are bending every effort to prepare a civilization over a century in arrears, to meet the rude shock of the worst of our culture. In the 1911 term of court, Perry County, being invaded rapidly by railway construction, had nearly 600 cases; Owsley County, without access to railways, had less than 40 cases. A mountain guide in Pound Gap lamented, "The devil is coming into the mountains on wheels." Eight years ago I rejoiced with a clean cut, delightful, energetic man who was returning home from the Kentucky mountains buoyant because he had doubled his fortune by securing some of the primitive forest at an absurdly low price. He was bringing wealth and good cheer to his northern family. Now, with those slopes in mind, deforested, gullied, scorched, and sold ("unloaded"), I am glad that I did not smoke then, for I probably should have accepted some of his fine Havanas. The *rapid exploitation* of the natural

resources of a region by outside capital tends to harm the native, especially if his civilization is not modern. In this case the outcome is in the balance.

THE FUTURE.

If exploitation pure and simple continues, twenty-five years will bid fair to bring about the following results: The disappearance of this race of true Americans as a unit; the passing of the valuable timber; numerous forest fires in the region slashed over; greatly increased erosion of the steep hillsides with their soil already thin; short periods of flood within and below the region; long intervals of low water within and below the region; the reduction of fish and game; the introduction of a foreign mining element, also a foreign manufacturing body; and a region of great natural beauty changed to a region of squalidness. Presently, with the increase of population and the value of land in the United States, the region may be reclaimed at great cost.

Outside aid might do the following things: Regulate the exploitation of the coal and timber so that it will be gradual; aid the counties in building good roads; assist in educating the mountain people along broad lines to close the gap between them and us; help them to develop stock raising, fruit growing, scientific agriculture, and scientific forestry. Some of the results would be: The saving of the mountain race as a unit; the addition of a happy, prosperous, food supplying area to the United States; the prevention of the disasters of soil erosion and of flood, and the utilization of water power.

It is being pointed out that men break down under the tension of modern industrialism, unless they, somehow, are brought into contact with the beautiful, and get away for frequent moments of change and recreation. The government owns our national parks; but they are far out West, beyond the financial reach of the average worker. The government might also establish numerous small parks in the Southern Appalachian Highlands, which would become the recreation ground of millions of workers east of the Mississippi River.

CONSERVATION AND CIVILIZATION.

ARTHUR L. FOLEY, Head of the Department of Physics, Indiana University.

Not until recently has man begun to think of nature's resources in the light of the old saying that "one can not eat his cake and have it." Today the subject of the conservation of these resources is being discussed by men of science in every civilized country on the globe. Nevertheless it must be admitted that the full import of the question is not yet appreciated by many science men, while the public generally scarcely knows what the discussion is about. A few years ago the writer heard an address in which the speaker pointed out the importance of conserving the soil. A fellow citizen in speaking of the address said he did not understand what the speaker meant, that the earth is made of mud and that no one need be fearful of a shortage. The observation led to the reflection that people too are made of mud, some of it not very fertile. Perhaps our citizen did not know that the average productivity of the unfertilized soil of Indiana is but half of what it was when he was a boy. Perhaps he does not know that all animal life is dependent on plant life, and that plant life is dependent on a few soluble constituents of the soil which form but a small and diminishing per cent. of what he calls mud. Perhaps he does not know that the removal of timber and the cultivation of hillsides permit the rains to dissolve and carry away the soluble constituents and so impoverish the soil, and that every year thousands of acres of land, here in his own State, Indiana, are ruined in this manner. He does not know that every year the Mississippi River robs the Mississippi Valley of hundreds of millions of tons of that upon which its fertility depends, and that all other streams are doing relatively the same thing.

But I am not to discuss the conservation of our soil, nor the conservation of our timber or our food supply. I shall not discuss the conservation of air or water, although the time has passed when we can say "as free as the air we breathe or the water we drink." Good pure air is not free to everybody, by any means. If it were there would be no

"white plague." Nor is good pure water free. Sometimes it can not be obtained at a price, as some of us well know. New York City is just now completing an additional water system, this last system alone costing one hundred and eighty millions of dollars.

I wish to direct your attention to a phase of conservation that receives less consideration than is generally accorded the conservation of timber, food, and soil. I wish to consider our fuel supply, and its relation to civilization.

I shall begin by defining the word energy as the capacity for doing work; that is, the capacity for exerting force through space. Anything that can do work has energy stored in it, the quantity of energy being measured by the amount of work the thing can do. For instance, a clock spring or a clock weight has energy which it expends as it runs the clock; a battery has chemical energy which it expends when it rings a bell or drives a motor; a head of water has energy which it expends when it drives a turbine; gasoline has energy which it gives out as it heats a kettle or drives a motor car; a chunk of coal has energy which it expends when it heats our home or produces the pressure to drive a steam engine.

If we represent all the heat energy in the universe by the letter "h", all the chemical energy by "c", all the electrical energy by "e", and so on, using a different symbol for each and every one of the many forms of energy, then we may express the law of the conservation of energy in the form of an equation— $h+c+e+$ all other kinds of energy—a constant.

This law expresses the fact that energy is indestructible. We can neither create it nor destroy it. What then do we mean when we say that we should conserve and save our energy? To answer the question we shall need to consider two other laws or principles of energy, known as the Principle of Transformation of energy, and the Principle of Dissipation or Degradation of energy.

By transformation of energy we mean the conversion of energy of one form into energy of some other form. For instance, the energy of the coal is transformed in the boiler into the heat and pressure energy of steam, the engine converts this into a mechanical energy of motion; this may be used to drive a dynamo which converts it into electric energy, this may be passed through a lamp and be converted into light energy or it may be used to drive a motor which converts it again into mechanical energy, and so on. There is known no kind of energy that man can not

transform into any other form of energy. He may change almost at will the relative value of the energy terms on the left side of his energy equation, but he can not change their sum. No energy is lost in any transformation. The total remains constant.

In actual practice, however, when we attempt to transform or use energy, we find that more or less heat energy is generated in the process, and that this heat energy is dissipated through space. No machine is frictionless, no wire is without its electrical resistance. There is no motion of matter or electricity that does not result in the generation of heat which can not be utilized, heat which soon disappears forever in space. This energy is not destroyed, it is lost. It is easy to transform energy of any kind into heat energy, but it is impossible to transform all of any quantity of heat energy into energy of other kinds. Our best steam engines have an efficiency of some twenty per cent., the efficiency of the usual engine is not over ten per cent. This means that we waste from 80 to 90 per cent. of the coal and utilize from 10 to 20 per cent. Our best tungsten lamps give us only about 10 per cent. of the energy of the electric current required to operate them, and since the engine that drives the dynamo has an efficiency of, say 10 per cent., the tungsten lamp gives us as light only 1 per cent. of the energy of the coal, 99 per cent. being wasted in the form of dissipated heat. So we have the principle of the dissipation of energy; however we transform energy, a fraction of it—usually a large fraction of it—is always dissipated as heat and is forever lost. Thus while the total quantity of energy in the universe remains constant, the useful or available energy is rapidly diminishing. All forms of energy are tending to go into the form of heat, to run down hill as it were, as heat is regarded as the lowest form of energy. Consequently this Principle of the Dissipation is sometimes called the Principle of the Degradation of energy. It makes no difference what the final temperature of the universe may be; when all other forms of energy have been transformed into heat the heat energy will be useless. We can not use heat energy except as it runs down hill, from points of high temperature to points of low temperature. It is doing this all the time. Diffusion is a property of heat, and must result sooner or later in a state of uniform temperature and consequently in the disappearance of all available energy.

Perhaps you ask why we should worry about the condition of things a million years hence? The reply is that we need not do so, but that we may well take thought of what the condition may be before the twentieth

century shall have ended. No doubt the disappearance of all available energy is a matter of millions of years, perhaps of billions of years. But the disappearance of so much of our available energy that what remains may be entirely inadequate to supply the demands of a civilization such as we now have, is not a matter of millions of years, not even of thousands of years.

The progress of man has been proportional to his mastery of, and use of, Nature's resources. Thus we have the Stone Age, the Bronze Age, the Iron Age, and the Steel Age—the age of today. A 1914 model automobile, if made of bronze or iron, would not run a mile. The invention of the automobile could not have preceded the invention of steel. Steel made possible the light weight engine of high power, without which flying machines would be impossible. Without steel most of the weapons, and instruments, and machines of today would be impossible. Without power they would be useless. So this has been called the Age of Power or the Age of Energy, or better still, the Age of Coal Energy, for coal supplies almost all the energies required to do the work of the world. King Coal reigns with a lavish hand. We feast at the table, apparently unmindful of the fact that we are nearing the dessert course of his final banquet.

Our boasted triumphs over past generations are due to the fact that we have learned to use energy freely. We are not superior to those of earlier ages, in art, in architecture, in music, in intellect. We are vastly superior to them in our ability to make use of Nature's mineral resources. I might even say in our ability to use coal, for without coal the production of iron and steel would be practically impossible and the mineral resources of the world would remain undeveloped.

It is high time that we were awaking to the fact that civilization as we know it must disappear from the earth when the available energy has been exhausted. Concern over the social, intellectual, religious, and political state of future generations is of secondary moment compared with the question of the existence of civilization itself.

Each individual knows that he must die. But if he thinks the event somewhat remote he scarcely gives the matter a thought. He may even indulge in things that he knows will surely hasten the event. So with a race. We give no thought for the morrow, but continue to use and to waste Nature's resources, knowing full well that the death of the race is the inevitable result, and that our prodigality is speeding the day. We make the mistake of supposing that the day is indefinitely removed. We

calm ourselves with the thought that the human race has inhabited the world for thousands, perhaps millions of years, and still Nature is bountiful. But we should remember that not until this century has there been any considerable draft upon Nature's store of energy in the form of coal.

Coal was not discovered in the United States until some two hundred years after the discovery of America. It was seventy years after its discovery before it was commercially mined. For many years the output of the mines was very small. Lately, the disappearance of our forests and the astonishing increase in the use of machinery have combined to make enormous demands on our mines. The coal used in the United States during the past nine years is in amount equal to the total consumption up to the year 1895. The output of the mines for the year 1912 was 535 million tons—about five tons for each man, woman, and child in the United States. The figures relating to petroleum are just as significant. The industry began in 1859, and it was twenty-four years before the entire output was equal to that of one year now. The output of the last eight years equals all that produced before.

Natural gas is all but a thing of the past. When scientists foretold the speedy exhaustion of the supply and cried out against its criminal waste, their cry was unheeded and the waste went on. If all the gas wells had been properly cared for, and if all gas had been sold through meters, we should have had the blessings of natural gas for a century to come. People scoffed at the idea of natural gas failing. So did the newspapers throughout the entire gas belt. The introduction of meters was fought by papers and patrons, until the finish of natural gas was in sight. Instead of educating the people to economy and care, the papers incited them to extravagance and indifference.

We are now passing through a somewhat similar experience with petroleum and coal. Many of our oil fields have been exhausted and abandoned. No wonder, when we note that the production of petroleum of a single year is a quarter of a billion barrels. This enormous production has been made possible only because of the discovery of new fields to replace the exhausted fields. The discovery of new fields can not continue indefinitely. Most of our territory has been explored. It is only a question of time, and not a very long time, when oil too will have become a thing of the past. There will remain but one natural fuel, coal, to stand between us and a return to a primitive type of civilization.

The life of the coal beds has been variously estimated at from one hundred to five hundred years. The time will be longer or shorter, depending on our frugality or our prodigality. Yet our newspapers, among them the same ones that fought the use of meters for natural gas, fight the utilization of the power of Niagara Falls, calling upon state and national governments to preserve this wonder of Nature, the inference being that the use of the power of the Falls would mean the destruction of the Falls. Of course Niagara should not be exploited for the profit of individuals or corporations, nor should the Falls be destroyed. It might be arranged to permit, at certain times, all the water to go over the Falls for the delight of man. But in the opinion of the writer it is short-sighted, it is almost criminal, to permit millions of horse power of energy to go to waste, continually and continuously, merely for our enjoyment. Does the reader think it right to burn millions of tons of coal each year that might be saved for future generations, all in order that we—some of us—may see the glory of Niagara? Who is sordid? The man who is willing to forego a magnificent spectacle for the good of future generations, or the man who would feast his eyes and let future generations freeze? How does the Niagara waste differ in principle from the uncapping and lighting of a natural gas well with the gas under a pressure of hundreds of pounds per square inch, in order that people might hear the roar of escaping gas and see the heavens illuminated by a giant flame?

I remember that when the American Association for the Advancement of Science met in Indianapolis in 1890, the committee on entertainment arranged for an excursion through the Indiana gas belt and a natural gas display. At one city pipes were laid in the river and the gas liberated under the water. We saw the river, in appearance, converted into a seething cauldron. The sight was grand, but not pleasing. A man of science could not avoid the thought that we were being entertained at a fearful cost to future generations. Recently the writer's attention was called to the possibility of that display in the end conserving the gas supply instead of hastening its exhaustion. The display may have served to arouse sentiment against such wanton waste and consequently to hasten legislation prohibiting it. This may have been true in this particular instance, for those who saw the waste were those to whom such a thing would make a strong appeal. But people generally saw reckless extravagance on every hand and were a party to it. The writer recalls

a gas well within six miles of his father's home that was permitted to burn almost a year before the flow was stopped. The gas wasted from that well alone would be sufficient to supply a city of moderate size for a hundred years. Truly we are reaping where we have not sown, and are leaving but little of the harvest for future generations. To realize the truth of this statement you have but to consider the enormous development in the use of mechanical energy during this generation, and the necessarily enormous consumption of oil and coal required to supply that energy.

The one-horse buggy has been superseded by the thirty horsepower runabout, the two-horse carriage by the forty-horse touring car, the two-horse wagon by the sixty-horse auto truck, the two-horse stage coach by the five-hundred horsepower locomotive. The horse car has given place to the electric car, the sail ship to the steamship or dreadnought, the canoe to the motor boat, the bicycle to the motorcycle, the foot or hand press to the power press, the typesetter to the linotype, the tallow candle to the electric lamp.

Once man ate what his own fields produced; now much of his food comes to him from the ends of the earth. Once man was content to worship in the little church at the cross-roads; now he must attend conventions in Boston or Los Angeles. Once he thought twenty miles a journey; now he travels a thousand miles to see a ball game.

Now the house wife must have her electric irons and cookers, power washing machines, and vacuum cleaners. The farmer must have his feed choppers, shredders, threshers, and pumps, all operated by power, lately by gas engine power. The thousands of windmills that dotted the country twenty years ago have disappeared—replaced by gas motors. The grocer grinds the coffee by electricity and delivers it with an automobile. The absurd extremity to which we have gone in the application of power is illustrated when an auto delivery wagon calls for and delivers a ten cent package of laundry. These things are little things, but they illustrate the spirit of the age. We do nothing ourselves that we can get Nature to do for us. We give no consideration to the fact that we are burning the condensed sunshine of bygone ages. Our only question is, "What does it cost?" What does it cost *us*? Not what it has cost Nature, or what it will cost future generations.

The value of coal is fallaciously reckoned on what it costs to mine and transport it. The fact that coal represents energy stored by Nature

through countless ages of time is not given a moment's thought. Figuring this way, if we can manufacture ice a penny a ton cheaper than we can harvest the natural ice, we proceed to burn coal to make it. Think of the waste. Burning coal to make ice, when all the ice we need could be had for the harvesting. You say that natural ice is not produced in the tropics. Neither is artificial ice, in any quantities.

We flood our streets with oil, because we think it a cheap way of keeping down the dust; cheap only because we fail to consider the energy content of the oil and what it has cost Nature to produce it. The time will come when such extravagance will be prohibited by statute.

The fact is that we fail to realize that oil and coal are a legacy that has come to us from bygone ages, deposited in Nature's bank. We are spending our substance in riotous living, but unlike the prodigal have no place to go when it is all spent. Doubtless something will fall on our neck, but there will be no fatted calf.

The writer has painted a gloomy picture, such a picture as would have been painted twenty years ago, with dark clouds hanging everywhere about the horizon. However, the picture needs but one change to represent the conditions today. There is a rift, a small rift, in the clouds; a rift that may close and leave us again with leaden and ever darkening skies; a rift that may open wider and wider and leave us finally with the glorious sunshine of a cloudless sky. Whence the rift?

The energy content of matter depends on position and motion, not only on the position and motion of the mass as a whole, but upon the position and motion of the constituent parts. Experience tells us that the energy liberated during any change is relatively greater the smaller the parts taking part in the change. For instance: the energy required to change a gram of water into steam, a change of position of the molecules, is twenty times as great as the energy of a speeding rifle bullet of the same mass. To effect an atomic change, that is, to separate the hydrogen and oxygen atoms which form the water molecules, requires five times the energy involved in the molecular change. When the atom itself breaks up, disintegrates, relatively enormous quantities of energy are liberated.

Radium is a substance in which this electronic or sub-atomic change is going on continuously and spontaneously. It is continually throwing off or radiating minute particles, and so we say that radium is radioactive. A mass of radium gives off enough energy every hour to melt more than

its own weight of ice; and it does this day after day, year after year, and it will continue to liberate energy until the last trace of the radium has disappeared, a process that we have every reason to believe will require ages of time.

Many other substances beside radium are known to be radioactive. All substances may be more or less radioactive, the difference being one of degree rather than of kind. However this may be, we now know that there is stored within the atoms of matter quantities of energy, intra-atomic energy, beyond the powers of man to estimate. This is the rift in the clouds. It was produced by the discoveries of Becquerel and the Curies.

The rift in the clouds is not quite as wide as it was a few years ago, for so far man has failed absolutely to influence these radioactive processes in the slightest degree. Whether at the temperature of liquid air or the electric furnace, in boiling acids or alkalies, whether in a vacuum or at a pressure of a thousand atmospheres, whether inside or outside the strongest electric and magnetic fields man can produce, the rate of disintegration and consequently the rate of liberation of energy appears to be absolutely constant. Perhaps we may not hope to be able to control a change in the atoms themselves, for have not the atoms existed through countless ages and successfully withstood pressures and temperatures in Nature's laboratory exceeding any that man can bring to his service in the chemistry or physics laboratory?

That this intra-atomic energy exists is not theory. It is a fact that is as well established as any fact in science. Man hopes some day, somehow, somewhere, to unlock this infinite storehouse of energy. Today Nature stubbornly holds the key. The probability of man being able to wrest it from her is anything but bright. But we should not be, we must not be, discouraged, for it is our only hope. If the secret is ever discovered and we succeed in tapping this supply of energy no mind can imagine the heights to which civilization will mount by leaps and bounds. If the secret eludes us civilization is doomed to return to a primitive state from which it can never emerge.

Perhaps you urge that our estimate of the life of the coal beds is too short. If it were in error by one hundred per cent., and no authority claims as much, the depletion of our coal supply is simply moved forward a few generations. The ultimate outcome is unchanged.

Perhaps you say that the writer has failed to consider the possibility

of using the energy of the sun's rays. You should remember that sometimes we do not have enough sunshine in Indiana in a week to supply heat for a cup of coffee. It is a fact that where heat is most needed, and when it is most needed, to heat our homes and run our factories, there and then is the least sunshine. Imagine London depending on sunshine for heat and power. In winter when we need the most heat the sun shines the fewest hours per day, the fewest days per week, and the sun's rays are most oblique. Taking into consideration the necessarily low efficiency of any engine working between the temperature limits of an engine for using the sun's radiation, and the very large surface from which the energy would have to be gathered, men of science are agreed that the prospect of a practical sunshine engine are exceedingly remote.

Finally it may be argued that the writer has failed to see a rift in the clouds arising from the possibilities of water power. The answer is, there is no rift there. No doubt the use of water power will postpone the gathering of the clouds, but it will not disperse them. Leaving out of consideration the fact that water power is usually most abundant where least needed, that the available power varies greatly with the seasons, that the available water power is diminishing from year to year with the removal of forests and the draining of swamp lands, let us remember the fact that the total water power of the world is almost nothing compared with man's demands.

A single ocean liner burns fifty car loads of coal per day. To supply the power for such a liner would require ten such water power plants as the one on White River at Williams, near Bedford, which cost several hundred thousand dollars. Then, too, it would require all the ten plants to operate at full capacity, which the Williams plant can not do a considerable portion of the year, the supply of water being insufficient. The writer is informed that it is not using water power at all as this is being written.

Every fifteen days the new automobiles marketed by a single manufacturer of cars of low horsepower equals the entire water power development of the Mississippi River, at Keokuk. Every thirty days the new engines turned out by this one firm equal in power the total water power developed at Niagara. The total horsepower of the automobiles now registered in the United States is greater than the estimated total available water power of the country.

It would appear that one need not go further to show the utter

inadequacy of water power as a substitute for oil and coal. Those who think otherwise usually consider the question from the standpoint of factory power only, leaving out of consideration the enormous quantities of energy required to heat our homes, and to supply heat for such processes as ore smelting, cement manufacture, brick, tile and glass making, and thousands of others. To equal one ton of coal per month for heating purposes one would require the entire output of a fourteen horse-power plant, running twenty-four hours per day thirty days per month. If there are five hundred thousand families in Indiana and if each family consumes an average of two tons of coal per month during the winter season, the consumption is the heat equivalent of fourteen million horsepower. Remember, too, that Indiana is not a very populous State and that its climate is not severe.

Professor Soddy states the facts in his little volume on "Matter and Energy" when he says that "the age in which we live, the age of coal, draws its vivifying stream from a dwindling puddle left between the comings and goings of the cosmical tide."

We are to "witness a race, a race between science on the one hand and the depletion of our natural resources on the other hand." This race will be run chiefly by pure science, not by applied science. Engineers and inventors make their reputations and their fortunes by devising new and improved methods of using our natural resources; they are not concerned with the atom, the latest and the greatest energy reservoir discovered by man. We must look to such scientists as Becquerel, Curie, Rutherford, Ramsay. We must look to the humble, overworked, underpaid scholar toiling away in his laboratory. If he fails us, darkness comes.

WHY DO OUR BIRDS MIGRATE.

D. W. DENNIS.

It is insectivorous and "therefore" a migrant, is a common phrase in literature about migrants; it is the purpose of this brief paper to take the therefore out of of this sentence: to maintain that what a bird eats has nothing to do with the great bird movement from the south to the north in the spring and back again in the fall after breeding.

The Pennsylvania reed bird, the bobolink, doubtless stops at the reed swamps in Pennsylvania for refreshments on its way south; the South Carolina rice bird, another name for the bobolink, takes toll of the rice swamps; but no one thinks that the reeds or the rice are the cause of the migrations. Surely if they had not wings, they could hardly fly from the equator to Manitoba, but this does not make their wings the cause of the journey; nor is their food the cause.

It is stoutly maintained that climate is the cause. This, like wings and food, renders the journey possible; but it cannot in all cases cause it, for many water birds, like the gannet and the petrel, go to their breeding grounds from colder to warmer water and many from warmer to colder. They go to inhospitable, inaccessible rocks that they may nest in a place of safety, as I believe.

I was impressed at Wood's Hole in the summer of 1901 to see tern flying by in great numbers every morning. Later I visited their breeding grounds at Penikese; they were flying by Wood's Hole to get food for the day; they had not come to Penikese for food, for they came in such numbers that they overtaxed the fishing grounds for more than twenty miles to the eastward. They had not come for climate, for they had come from all available areas, colder as well as warmer. Perhaps it is admitted that they came to lay their eggs and rear their young safe from destructive mammals, including boys.

The facts about the blackpoll warbler sustain this theory almost as well as those about the tern, the gannet or any other water species.

The Blackpoll. It winters south of the equator and nests north of the Arctic Circle; its journey to its breeding grounds is a 10,000 miles round trip. It passes through Richmond about May 15 and returns September 15; its movements as it passes by us are deliberate. It cannot spend more than two months in its northern habitat; these must be very busy months. Nest making and family rearing are its chief business during these two months.

In a few minutes, or at least hours, the salmon prepares his nest and lays his eggs 1,000 miles up the Columbia from the Pacific, and we conclude he came for this.

In two months the blackpoll prepares its nest, lays its eggs, hatches its young, and rears them beyond the most critical periods of their existence, and starts back. Did it come to eat insects on the way, or to discharge this race duty? It is a ground nester; on or near the ground in that high latitude its eggs and family are safe from nest-robbing reptiles which abound in the warmer districts where it makes its winter home. Does it not make it wisdom's child, if it makes this long journey to nest in safety? If, as Aristotle said 2,500 years ago, the study of zoology is a study of fitness, it is real zoology to study the migrations of such birds as the blackpoll.

This argument applies to the water birds, which in countless numbers and numerous species fly over Indiana in early spring. The great majority of these nest on the ground near lakes and streams; some of them on floating islands in lakes, just the places where the eggs and young would be unsafe in their winter homes on account of reptiles.

The young of these birds swim almost from the shell, and would be reasonably sure to be eaten in southern waters.

The argument applies with almost the same force to all indefensible ground and low bush nesters, among which are the field sparrows, the vesper sparrow, dick-cissel, grasshopper sparrow, Savannah sparrow, bobolink, meadow lark, ground robin, brown thrasher, etc.

Nearly all our migratory birds show protective coloration, or sexual dimorphism; these are a confession of inability to take care of themselves or their homes, in fight. Those that exhibit sexual dimorphism are—

Bluebird,
 Robin,
 Redstart,
 Canadian warbler,
 Wilson's warbler,
 Hooded warbler,
 Yellow-breasted chat,
 Maryland yellow-throat,
 Mourning warbler,
 Connecticut warbler,
 Kentucky warbler,
 Prairie warbler,
 Pine warbler,
 Kirtland warbler,
 Black-throated green warbler,
 Blackburnian warbler,
 Blackpoll warbler,
 Bay-breasted warbler,
 Chestnut-sided warbler,
 Cerulean warbler,
 Magnolia warbler,
 Myrtle warbler,
 Black-throated blue warbler,
 Yellow warbler,
 Cape May warbler,
 Northern parula warbler,
 Parula,

Tennessee warbler,
 Orange crowned warbler,
 Nashville warbler,
 Golden-winged warbler,
 Blue-winged warbler,
 Prothonotary warbler,
 Black and white warbler,
 Summer tanager,
 Scarlet tanager,
 Dickcissel,
 Indigo bunting,
 Blue grosbeak,
 Cardinal,
 Towhee,
 Junco,
 American goldfinch,
 Redpoll,
 Purple finch,
 Pine grosbeak,
 Evening grosbeak,
 Baltimore oriole,
 Orchard oriole,
 Redwing,
 Yellowheaded blackbird,
 Cow bird,
 Bobolink,
 Humming bird,

fifty-four in all. The proof which all of them furnish is the same which the blackpoll warbler furnishes. They have, perhaps, come a less distance in all cases, and stayed a somewhat longer time.

All the balance of our migrating birds exhibit protective coloration, or are very inconspicuously colored—a confession of inability to protect the nest and an argument that birds migrate to protect it. A few conspicuous examples are:

Meadow lark,
 Vesper sparrow,
 Little brown creeper,
 Field sparrow,

Night hawk,
 Whippoorwill,
 Rails,
 Quail.

Our birds which build protected nests, or which are able to protect their nests are not migratory birds as a rule. I know of but one clear exception, the sapsucker. Our birds which build unsafe nests or which cannot protect them are migratory birds as a rule. I know of no exception that is clear. The Phoebe arrives when its food is scarce, and it leaves a land of plenty, a land of insects; food cannot be the attraction. That climate is not the compelling cause is shown by the fact that many birds arrive when the climate is very severe; it even kills thousands of them sometimes.

That birds are indigenous in the north; that they are migrating in the fall instead of the spring; that in the spring they are just going to their preglacial home; and that nostalgia is the real cause requires us to believe that birds have a way of preserving a record of their lost Atlantis that we do not possess, and may be dismissed as wholly psychological.

The salmon goes a thousand miles up the Columbia to spawn; the eel question has at last been solved; it goes to the deep sea to spawn. Sea-birds go to isolated rocks for the same purpose. It is the conclusion of this paper—there being no shred of evidence against it, and many weighty reasons for it—that our migratory birds go north for safety in nesting.

FLOOD PROTECTION IN INDIANA.

W. K. HATT.

Organized effort to study the causes and to lessen the effect of floods in Indiana begins with the appointment of the Indiana Flood Commission by Governor Ralston on April 20, 1914.

This commission is composed of one member from each congressional district and the personnel is as follows:

- Mr. E. W. Shirk, Peru, Chairman.
- Professor W. K. Hatt, Purdue University, Lafayette, Chief Engineer.
- Mr. Frank C. Ball, Muncie.
- Mayor Benjamin Bosse, Evansville.
- Mr. William Cronin, Terre Haute.
- Mr. Stephen B. Fleming, Ft. Wayne.
- Mr. J. H. Frederick, Kokomo.
- Mr. S. J. Gardner, New Albany.
- Mr. Victor M. O'Shaughnessy, Lawrenceburg.
- Mr. Joseph C. Schaf, Indianapolis.
- Mr. W. N. Showers, Bloomington.
- Dr. Chas. K. Stoltz, South Bend.
- Mr. Herman Trichler, Brookville.

The commission met first in Indianapolis, on April 30, 1914.

The purpose of this commission is to consider the extent of damages due to floods in the State of Indiana, and to report to the Governor what measures should be taken to provide relief in the future.

The commission expects to issue its final report in 1915. This report will contain a full presentation of the history of floods in Indiana, a summary of the causes of floods, a collection of available data on rainfall, river discharge and topography, a discussion of flood protection works and a discussion of the principles of legislation to provide for flood relief.

This present pamphlet is an abstract of the forthcoming report of the commission, prepared in non-technical style for general information.

FLOOD OF MARCH, 1913.

The appointment of this commission was the direct result of the flood of March, 1913, in which 467 lives were lost and over \$160,000,000.00 of property destroyed in the United States. The memory of this catastrophe is still fresh in the minds of the people of Indiana, in which State thirty-nine lives were lost, and over \$18,000,000 of property destroyed.

The total loss in the flood of March, 1913, can never be known. The interruption of transportation and of business, the destruction to farm lands by cutting of banks of rivers and covering of bottom lands with gravel, the loss of productive capacity of manufacturing plants, and the sickness following exposure, are not susceptible of exact computation.

Professor Beede of Indiana University reports a total damage of approximately one-half million dollars in seven counties in the lower White River basin, in which also nearly eight thousand acres of agricultural land were denuded of soil and some sixteen thousand acres of river bottoms were covered with soil and silt. He estimates the loss to agricultural land in this region as nearly \$250,000.00.

The loss reported by county auditors to county roads and bridges alone, was over \$3,000,000.00. Other tangible losses that have been determined are shown in Table 1. It is probable that the loss during the flood of March, 1913, in Indiana, may be estimated at over \$25,000,000.00.

Indeed the catastrophe was so general over the Ohio Valley that it excited the sympathy and support of the entire nation. The Governor of the State of Indiana received \$———— in subscriptions for the relief of flood sufferers in this State.

PART OF DAMAGES SUSTAINED IN THE FLOOD OF MARCH, 1913.

1. County highways and bridges.....	\$2,825,240 00
2. Railroads—steam	5,299,810 00
3. Electric railways	788,000 00
4. Buildings and personal property.....	8,104,250 00
5. Telephone and telegraph.....	17,510 00
6. Crops	735,700 00
7. Livestock	149,380 00
8. Farm lands	264,700 00
9. Suspension of business.....	582,000 00
Total	\$18,766,590 00

Counties not included in (1)—Cass, Clinton, Fayette, Floyd, Miami, Sullivan.

Railroads not included in (2)—Central Indiana R. R., Chicago and Wabash R. R., Cincinnati, Hamilton & Dayton R. R., Toledo, Peoria & W. R. R., Toledo, St. L. & W. R. R.

Electric lines not included in (3)—Marion and Bluffton Traction Co., Bluffton, Geneva & Celina Traction Co., Central Indiana Lighting Co., Indianapolis Street Railway Co., Louisville and Southern Traction Co., Louisville and Northern Railway and Light Co., Vincennes Traction Co., Washington Street Railway Co.

(5) Includes Indianapolis Telephone Company only.

Counties in flood districts not included in (4), (6), (7), and (9)—Adams, Blackford, Cass, Clark, Clay, Clinton, Fayette, Floyd, Fountain, Franklin, Gibson, Grant, Greene, Harrison, Howard, Huntington, Jay, Jefferson, Ohio, Parke, Perry, Putnam, Randolph, Ripley, Scott, Sullivan, Switzerland, Tippecanoe, Vanderburg, Vermillion, Vigo, Wabash, Warrick, Wells, White, Whitley.

(8) Includes loss only in 230 miles of East and West Forks of the White River through Morgan, Owen, Greene, Daviess, Knox, Jackson, Lawrence, and Martin counties.

First there are six main problems to be solved before our Indiana communities can protect themselves against floods.

FIRST PROBLEM.

Flood Flow.

First there must be proper information as to the amount of water carried safely in a channel. To determine this amount we must first know the rainfall that may reasonably be expected at a time not too remote, and the rapidity with which this rainfall runs down the watershed.

In considering flood protection in Indiana we are barred at the outset from a sure solution at present, first, on account of a lack of rainfall records over a sufficiently long time; second, by a lack of stream gaging to determine the amount of water which does run down our streams during heavy rains; and third, by a lack of surveys of watersheds.

In other words, a heavier storm than any that has been recorded in the last thirty years of our rainfall records, may come in the future, but our records do not serve to determine the probable extent of this storm.

Again, we have not gaged our streams to know the relation between the runoff and the rainfall. Such records as are gathered in other communities will not apply to our peculiar conditions, that is, two watersheds of equal area, one long, narrow and V-shaped, and the other broad and flat, will yield very different flows in the streams. Again, the character of the surface, whether of rocky formation or swamps or farmland, will change the conditions.

Therefore, to obtain an exact solution of our flood problems we must first of all get accurate surveys and determine the flow of our streams. This cannot begin too soon. For this reason, the Indiana Flood Commission recommends an early beginning of this work of surveys and stream gaging.

These surveys are most important for another purpose, namely, to determine if the water of the upper reaches of the rivers can be held back for a time in reservoirs. For instance in the case of the Wabash River at Logansport, which carries the floods from the upper Wabash, the Mississinewa, the Salamonie, and the Eel River, we would like to know if it is possible to find reservoir sites in the valleys of these tributaries, so that the flood flows may be controlled. Each tributary flood might be held back to the proper amount, and for the proper time, so as to let these flood flows by Logansport one by one.

For example, in Ohio, it was found that by reservoir control, flood protection could be obtained for the cities of the Great Miami Valley at a cost of \$17,000,000.00, whereas the total sum of the cost of the individual protection schemes gotten up by each city acting separately was over \$100,000,000.00. The study of reservoir protection for the Miami Valley was made by the use of the topographic maps of the State of Ohio from which reservoir sites were planned and preliminary estimates worked up. Later on, detailed surveys showed that the preliminary work was accurate to within one per cent. The topographic survey of Ohio is 87 per cent. complete, whereas the Indiana survey is only 9 per cent. complete. If we were fortunate enough to possess topographic maps of the State of Indiana, we could go ahead immediately to study flood protection in a more complete manner.

The topographic map of the State is not only necessary for complete flood protection studies, but it is of use in the following:

(1) As a preliminary map for planning extensive drainage projects, showing areas of catchment for water supply, sites for reservoirs, routes of canals, etc.

(2) For laying out highways, electric roads, railroads, aqueducts and sewerage systems, thus saving the cost of preliminary surveys.

(3) In improving rivers and smaller waterways.

(4) In determining and classifying water resources, both surface and underground.

(5) In determining routes, mileage, location of road-building material, and topography in country traversed by public highways.

(6) In classifying lands and in plotting the distribution and nature of the soils.

(7) As base maps for the plotting of information relating to the geology and mineral resources of the country.

Our first problem is therefore to gather reliable information as to stream flow and topography.

The Indiana Flood Commission, however, realizes that critical conditions exist in several cities which can not wait the ten or twelve or fifteen years required for the completion of such surveys. The commission has therefore made the best solution it can, and has studied all available records, has computed rainfall and runoff, and determined to the best of its ability, the amount of water which an Indiana city may expect to take care of during future flood time.

Briefly, the records of the heaviest storms in the Ohio Valley region have been studied and the relation between the drainage area and the inches of rainfall worked out for these storms. Several of these storms have been studied, notably those of October, 1910; January, 1913, and February, 1884. For instance, it was found that the center of the storm in January, 1914, was over Southwestern Kentucky; the center of the storm of March, 1913, was over a line from Mt. Carmel, Ill., to Richmond, Ind. It is reasonable to expect as a matter of chance, that similar storms in the future will be centered fifty to one hundred miles from its former center. Cities must therefore reasonably expect to take care of such storms.

The result of the study is equivalent to fixing a future expected rainfall as equal to that of the storm of March, 1913, plus one-third additional in the White River Valley, and one-fourth additional in the Wabash watershed. Small drainage areas are yet to be studied. The river dis-

charge resulting from the specified rainfall is determined from river gagings at selected points during the flood stages of March, 1913. Adjustments are made for various rainfall and channel slopes directly as the rainfall and as the one-fourth power of the slope.

To determine the area of channel or bridge opening to carry this flow, the commission suggests tentatively six feet per second as a flood velocity through an improved channel, and not over eight feet per second as a velocity through bridge openings. In any particular case, special study of channel conditions must be made. The Indiana Flood Commission has thus proceeded with the compilation of recommended bridge openings throughout the various parts of the State, as an approximate solution of our present difficulties. A survey of actual bridge openings through the State accompanies this study.

SECOND PROBLEM.

Design of Works.

The second problem is to design flood protection works to take care of the water which is recommended to be carried. This is not a difficult problem, involving only good engineering knowledge and judgment.

These flood improvements will consist in improvement of the channels of the rivers involving cleaning and straightening the river bed and lengthening the bridges, and removing obstructions, and secondly the building of levees to retain the flood heights. If proper surveys exist, reservoir control may be studied.

The Indiana Flood Commission has gathered together a number of plans that have been drawn for the Indiana cities, and it is in a position to assist communities that desire advice on the nature of flood protection works.

THIRD PROBLEM.

Construction Work.

After complete information has been gathered, and the best engineering skill has been operating, a third and most important step must be taken. There must be some organization to finance and build flood protection works. In other words some legislative action must be taken, some so-called enabling legislation. In any community some agency must be created to determine the necessity of improvements, to direct their construction, and to establish an assessment roll for benefits and damages

within a district defined in advance. And this agency must be appointed and directed by the courts or by a State board.

This is the crucial problem. It involves the coördination of several, at present unrelated agencies, as for instance the city government, the county commissioners, and the railways.

Of what benefit is it to a city like Peru, to spend \$350,000.00 on a levee, if this scheme demands for its proper action the lengthening of a county or railway bridge, when the county commissioners or railway officials refuse to coöperate.

It must also be remembered that we all have gone ahead creating new obstructions in the flood plain and in the channel which interfere with the flow of our flood waters. Railways, cities and county commissioners are responsible for the conditions. Channel obstructions must be removed, and either the State or the Federal Government must take action. Some control must be exercised over present as well as future constructions in the channels.

FOURTH PROBLEM.

Valley Protection.

When we take a wider view than that of the specific problem of a single city, we must consider a flood protection scheme from the standpoint of the watershed as a whole. One city in Indiana has made flood protection plans which deflect the water around the city, and throw it around in increased volume on its neighboring down stream. Cities often content themselves with sluicing the water through the cities and pile them up on communities below. Here is again the problem of state action to protect the whole people. Fortunately this is not merely an action of control, but means a wider viewpoint that may disclose a cheaper and better method of protecting the whole valley.

FIFTH PROBLEM.

Maintenance.

After these works have been constructed, we have a fifth problem in their maintenance. It must be recollected that these works are built to protect against floods which happen only once or twice in a generation. Naturally such works as levees and reservoirs will tend to be neglected during this unused interval. If people construct dwellings and operate industries in a space supposedly protected by improperly maintained reser-

voirs, or levees, they are in jeopardy. In this case the State must exercise some power to protect the people and see that these works are maintained.

SIXTH PROBLEM.

Federal Action.

In considering the question of floods the view is successively of city, of county, of watershed, of State; and finally the rights and duties of the Federal Government come into view. Our present problem is to delimit and properly apportion the action and responsibility as between the States and the Federal Government. At present the Federal Government controls all openings and obstructions in navigable streams. The logic of the situation would extend this to the upper reaches, because what happens there will affect navigation below.

For instance, if, due to obstructions, bars pile up on bridges and soil is washed down and creates bars below, there is a real connection between the upper reaches and the lower parts of the river.

Again, the Weather Bureau is in the best position to take observations of rainfall, and the Geological Survey can best and does make the topographic surveys, and the stream gaging.

Thus in this problem, the complex question of the division of water control, as between the States and the Federal Government, is to be determined in the future. A watershed is a natural unit, and not a political unit. There should be some coördination between the States in the Ohio Valley, whose problems are very similar.

AN APPARATUS FOR AERATING CULTURE SOLUTIONS.

PAUL WEATHERWAX.

A number of experiments on various phases of plant physiology, in each of which it was necessary to secure a constant stream of air continuing for several days, has led to the construction of a very efficient piece of apparatus for that purpose. The apparatus used by Prof. D. M. Mottier several years ago for aerating artificial cultures of algæ was modified by F. L. Pickett and used in a series of experiments on desiccation; and the writer has made some further changes in the construction of the apparatus shown in the figure and described below. This is now being used very successfully in the aeration of culture solutions.

The principle employed is that of the Sprengel mercury pump (water being used as a liquid in this case) by which bubbles of air are entangled in a stream of liquid which flows into a closed vessel. The only thing that remains to be done is to separate the air and the liquid, which are under slight pressure, and convey them from the reservoir by separate tubes.

The first problem is that of getting a stream of water that will flow uniformly. An attachment to a water pipe is usually sufficient for this. If this is not satisfactory, however, a siphon may be arranged to give a uniform flow. D, in the figure, is an ordinary battery jar provided with a siphon, B, which has an adjustable stopcock. A, which taps a water pipe and has an adjustable stopcock, supplies the jar with water a little faster than it is taken out by the siphon, B. Another siphon, C, removes the excess and keeps the water always at the same level, determined by its outer end, thus assuring an even flow, which should be just fast enough to cause the water to fall as a succession of drops.

The funnel, E, made by fitting a stopper into the end of a short piece of glass tubing about 1 cm. in diameter, has the end of the slender tube, F, extending 2 or 3 mm. above the cork. By means of this arrangement the water dropping into the funnel is caused to descend through the tube as a series of drops separated by spaces filled with air. Thus, if no escape is allowed, the reservoir, K, is filled with water and air under a pressure


equal to that of the aggregate length of all the water drops in E at any one time. But the air escapes through H, under control of a pinchcock, and the water is forced out through G. The waste water escapes through J.

The flow of air through H must be so regulated that water is forced out through G just a little faster than it enters from F. This provides for an occasional release of surplus pressure by the escape of air through G and prevents the filling of K with water, as will be the case if the air is allowed to escape too fast through H. The only irregularity of flow is at the time of the release of pressure through G; but the air stream is seldom interrupted for more than a few seconds, and by careful adjustment the frequency of these interruptions may be reduced to a minimum. Perfect adjustment would entirely eliminate these irregularities by allowing the water to escape through G just as fast as it enters through F; but, perfection being impossible, it is better to have the interruption occur as an escape of air through G than of water through H.

Theoretically the pressure of the air issuing from H, and consequently the depth to which a solution can be aerated, is determined by the vertical distance from the level of the water in K to the outlet of G. In practice, however, the apparatus falls somewhat short of this, due to friction of the air through H and the capillarity of the liquid to be aerated. The density of the culture solution is, of course, a determining factor also.

The efficiency of the apparatus depends largely upon the nature of the tube F. If it is of too small bore, the friction is too great; and if it is too large, the water has a tendency merely to run down the inside surface and fails to carry any air with it. A very satisfactory size of tube is one having an internal diameter of 2 to 4 mm. If a larger quantity of air is needed at H, the pressure to remain the same, it is better to use two tubes for F than to try to increase the capacity by substituting one larger tube. If the pressure is to be increased and the amount of air to be delivered in a given time is to remain the same, G must be lengthened, and this may necessitate the lengthening of F also, for F will carry air only so long as the aggregate length of its water column is greater than that in G. In adjusting the apparatus, glass or metal stopcocks have been found more satisfactory where the flow of water is to be regulated, while pinchcocks on pieces of rubber tubing have been found best for regulating the stream of air.

When well adjusted and in good working condition the apparatus is economical. Tests on the one now in use have shown that it can be made



An Apparatus for Aerating Culture Solutions.

to deliver 50. c. c. of air per minute at a depth of 15 cm. in a $\frac{1}{2}$ per cent. Knops solution at the expense of 50 c. c. of water. At this rate less than 20 gallons of water per day would be used with the apparatus running continually.

The apparatus as now in use is designed for the aeration of water cultures, but its wide range of adjustment and its economy will permit its being used for many other purposes. Various devices may be attached at H for changes of temperature, humidity, or chemical nature of the air, provided that allowance be made for the increased pressure that may be necessary.

Where it is desired that the stream of air be carefully guarded from outside contamination, this apparatus is clearly the superior of any by means of which the air is drawn through tubes by an aspirator at the end, for it is a decided advantage in such cases to have the pressure, which determines the direction of any possible leakage, outward rather than inward.

*Indiana University,
Bloomington, Indiana.*

ANTAGONISM ON *B. FLUORESCENS* AND *B. TYPHOSUS* IN CULTURE.

P. A. TETRAULT.

It is a fact long established that when two organisms live together in close relationship, the association will be one of tolerance, of mutual benefit or of one-sided injury. The term antagonism as used in this paper has more the meaning of one-sided injury. The phenomenon, for the bacteria, was recognized as far back as 1888 when Freudenreich and Garre, working independently, demonstrated specific antagonisms between given bacterial forms. The last named worked especially with *B. typhosus*. It was found that the typhoid organism did not thrive in a medium where certain other bacteria had previously grown; in other words, the cell secretions were toxic for *B. typhosus*.

W. D. Frost,* working on this same problem, discusses a number of theories advanced to account for this phenomenon.

One theory is that of the exhaustion of the food supply. All the available food has been extracted from the medium by the first organism growing on it. This was controverted by Olitzky by demonstrating that *Micrococcus aureus* would grow on a medium which had nourished a previous crop of bacteria but which did not permit the growth of *B. typhosus*.

Another theory was that of enzyme action. This, Frost says, could not hold in this case because enzymes are colloidal in nature and could not pass through a collodion membrane.

A history and comparison of the different cultures used in my work is given below.

All the cultures came from The Museum of Natural History, New York. No. 29 was obtained originally from the University of Chicago and was isolated from the swimming pool. No. 469 came from the Kral laboratories, Germany. No. 31 also came from the University of Chicago.

*The Antagonism Exhibited by Certain Saprophytic Bacteria against the *B. typhosus* Gaffky. Jour. of Inf. Diseases. Nov. 5, 1914.

was isolated from the Mississippi River and was labeled *B. fluorescens liquefaciens*.* No. 502 came from the University of Vermont and was labeled *B. fluorescens tenuis*.

When grown on various media these cultures gave the following reactions:

TABLE I.

Media.	No. 29.	No. 469.	No. 502.	No. 31.
L. milk	Digested.	Digested.	No reaction.	No reaction.
Lac. broth	No gas.	No gas.	No gas.	20%.
Nitrate broth.	Nitrates reduced.	Nitrates reduced.	Nitrates reduced.	Nitrates reduced.
Pep. broth	No indol.	No indol.	No indol.	No indol.
Gelatin.	Liquefied.	Liquefied.	Not liquefied.	Not liquefied.†

The only appreciable differences between these two groups of cultures lie in the litmus milk and gelatin reactions. This would suggest that the process was one of digestion, but by direct microscopic methods it could not be determined. When *B. typhosus* was mounted in some of the sterilized *B. fluorescens* filtrate and examined under the microscope, no agglutination was observed.

When the two organisms *B. fluorescens* and *B. typhosus*, are grown in parallel streaks on solid media, it is found that there always remains a zone between the two where no growth occurs, *B. fluorescens* gives off a pigment which facilitates the study of this phenomenon by microscopic methods. *B. typhosus* never trespasses over the green border line put up by *B. fluorescens*. This suggested a further study of the two organisms in liquid media. The method was practically the same as that used by W. D. Frost in his work on "The Antagonism Exhibited by Certain Saprophytic Bacteria against the *B. Typhosus Gaffky*" and described in his article on "Collodion Sacs."†

A gelatin capsule is fastened onto the end of a glass tube by heating the tube slightly before applying the capsule. The capsule and part of the tube are then dipped into collodion and allowed to harden. After a few dippings the sac is strong enough to stand without the aid of the gelatin. The gelatin is dissolved by means of hot water and the sac is ready for use. The sac is filled with nutrient broth and inserted into a flask con-

*Although labeled *B. f. liquefaciens*, No. 31 failed to liquefy gelatin.

†Reports and Papers of the Am. Pub. Health Assn. Vol. 28.

taining the same kind of medium. After sterilization in the autoclav, the sac and flask are inoculated with the different cultures to be studied. The sac prevents the bacteria from mingling but, being permeable, permits the diffusible products of metabolism to distribute themselves uniformly throughout the liquid medium. By taking samples from both tube and flask and plating, it becomes a rather simple matter to determine whether the life or the growth of either organism is affected by the manufactured products or wastes of the other.

The experiments were run in series. Each series consisted of five flasks. Four of these contained *B. typhosus* in the sac and one of the cultures of *B. fluorescens* in the flask. The fifth was used as a control and contained only *B. typhosus*. Four of these series were run simultaneously. The temperature was 37 degrees Centigrade. The experiments ran through a period of twelve weeks.

Of the strains of *B. fluorescens* used, cultures No. 29 and No. 469 imparted a very deep color to the medium after growing for twenty-four hours; No. 31 and No. 502 imparted very little color.

The next table shows a certain correlation between the elimination of *B. typhosus* by *B. fluorescens* cultures secreting a deep colored pigment as compared with those cultures secreting very little pigment.

TABLE II.

After twenty-four hours incubation.

Flask-containing <i>B. fluorescens</i> .	Sac containing <i>B. typhosus</i> .
No. 29	Growth.
No. 469	Growth.
No. 502	Growth.
No. 31	Growth.

After forty-eight hours incubation.

No. 29	No growth.
No. 469	No growth.
No. 502	Growth.
No. 31	Growth.

After seventy-two hours incubation.

No. 29	No growth.
No. 469	No growth.
No. 502	Growth.
No. 31	Growth.

The above table shows that the secretions of *B. fluorescens* in the flask penetrate the collodion sac containing the typhoid culture. There takes place, then, not merely an inhibitory action, but an actual bactericidal one. The secretions have to be of a certain concentration before this action takes place. It is found that the above is not true for the cultures showing slight chromogenesis.

In the next experiment, *B. fluorescens* was planted and allowed to grow before *B. typhosus* was introduced into the sac. The following table shows the results obtained.

TABLE III.

After growing *B. fluorescens* for twenty-four hours, inoculating the sac with *B. typhosus*, and again incubating for twenty-four hours:

<i>B. fluorescens.</i>	<i>B. typhosus growth.</i>
No. 29	Absent.
No. 469	Absent.
No. 502	Present.
No. 31	Present.

After growing *B. fluorescens* for forty-eight hours, inoculating the sac with *B. typhosus*, and again incubating for twenty-four hours:

<i>B. fluorescens.</i>	<i>B. typhosus growth.</i>
No. 29	Absent.
No. 469	Absent.
No. 502	Present.
No. 31	Present.

This table shows that once the toxic substances are produced in sufficient quantities and time enough is given for them to penetrate the sac, the typhoid organisms will not grow.

In the next experiments the fluorescens organisms were grown for ten days, then filtered and the filtrate sterilized for ten minutes at a pressure of fifteen pounds in the autoclav. The sterilized filtrate was then inoculated with *B. typhosus* and at the end of twenty-four hours samples were plated with plain agar and incubated.

The results are shown in the following table:

TABLE IV.

<i>B. fluorescens.</i>	<i>B. typhosus</i> growth.
No. 29	Absent.
No. 469	Absent.
No. 502	Present.
No. 31	Present.

The active substance is not destroyed at the temperature of live steam under fifteen pounds pressure for fifteen minutes.

CONCLUSIONS.

The specific antagonism of *B. fluorescens* for *B. typhosus* is due to a substance secreted by the first named organism.

This antagonism is not characteristic of all *B. fluorescens* cultures. There seems to be a correlation between the intensity of the color and this property.

The action is bactericidal.

B. fluorescens cultures with slight pigment production do not prevent the growth of *B. typhosus*.

The metabolic substances must reach a certain concentration before they become effective.

The toxic substances secreted by *B. fluorescens* have the following properties:

1. They are thermo-stabile.
2. They are diffusible through a collodion sac.

Although the growth of *B. fluorescens* in milk would suggest a digestive process, the typhoid bacilli are not agglutinated when grown in a sterilized filtrate of the first named organism.

NOTES UPON THE DISTRIBUTION OF FOREST TREES IN INDIANA.

STANLEY COULTER.

It is recognized at the outset, that even in a restricted area, such as that under consideration, much of uncertainty is given to any conclusions drawn because of lack of exact data covering every part. The richness of the tree flora in a given county as contrasted with that of adjoining counties, is usually to be explained by the presence of a skilled and persistent worker. It is very certain that no county in the State is entirely destitute of trees, in spite of the fact that not a single species may be reported as occurring within its bounds in any published report.

It frequently happens, also, that reports covering certain counties have been based upon the work of untrained observers who have failed to discriminate closely related species or who have made incorrect determinations, in either case confusing the situation and necessitating a careful revision of the data. While effort has been made to eliminate errors arising from such causes it is more than probable that some have escaped detection, but in the main as regards the species discussed the data seem accurate and complete.

Of the one hundred twenty-six indigenous species mapped in connection with this study, the larger part by far are probably of general distribution throughout the State. Variation in the density of the stand and in the size and form of individual trees are of course found, but the non-occurrence of any one of the species of this group in any county, under favorable conditions, would be more notable than its presence.

A relatively small group is confined to the extreme northeastern counties of Lake, Porter and Laporte.

Quite a large group is restricted to the southern counties or those lying in the first three tiers north of the Ohio River. A peculiar tree flora within this group is that of Posey, Gibson and Knox counties, lying along the lower stretches of the Wabash River. Some of the trees reported from this subdivision no doubt are of wider range than indicated, since few

counties in the State have been worked with such persistence and with such painstaking accuracy. Others of the species are doubtless restricted to the rich alluvial soils of the lower Wabash Valley.

To these might be added a small group showing a distribution so unconnected as to be extremely difficult of explanation. An instance is found in *Pinus Strobus* L., which is found in Lake, Porter and Laporte, extreme northwestern counties; Warren and Montgomery, western-central counties; and in Clark in the southeastern part of the State. Its occurrence in the northwestern counties affords no difficulty; the western-central location might be explained as a continuation of the former area; but the recurrence of the species in Clark County on the Ohio River in the eastern part of the State furnishes a difficult problem. In the case, also, of *Crataegus coccinea* L., we find equally unrelated areas, this species being reported only from Noble and Steuben, extreme northeastern counties, and Floyd, a southern county on the Ohio River. The genus *Crataegus*, however, presents such difficulties in the discrimination of species that the case just cited may possibly be due to lack of accuracy in determination.

The species limited to Knox, Gibson and Posey counties or to some part of the area are as follows:

- Taxodium distichum* (L.) L. C. Richards.
- Hicoria Pecan* (Marshall) Britton.
- Quercus lyrata* Walter.
- Quercus Michauxii* Nuttall.
- Quercus falcata* Michaux.
- Celtis mississippiensis* Bosc.
- Crataegus viridis* Linnæus.
- Crataegus nitida* (Engelmann) Sargent.
- Gleditsia aquatica* Marshall.
- Ilex decidua* Walter.
- Forestiera acuminata* (Michaux) Poinet.

In addition to these species, which seem strictly limited to the region named, two others have been reported from a single additional county:

- Catalpa speciosa* Warder. Gibson, Knox, Posey and Vigo.
- Fraxinus Michauxii* Britton. Gibson, Posey and Marion.

The Bald cypress (*Taxodium*) is a southern swamp form, which finds in the Indiana locations its extreme northeastern limits. In Indiana it is

found only along the wet (often submerged) banks of streams or in river swamps or sloughs. The local distribution has been carefully worked out by Deam.¹ While not as large a tree as in the South Atlantic and Gulf States the species as found in Indiana often reaches a height of 145 feet and a diameter of six feet. The species may be regarded as having entered the State in the period of flooded streams, maintaining its foothold in situations unfavorable for the ordinary species of this latitude. The areas in which the species occurs in the State are being rapidly reduced by agricultural operations and its disappearance from the tree flora of the State seems inevitable.

The Pecan (*Hicoria Pecan*) nowhere wanders far from the lowlands adjoining river courses. The species has been so largely cultivated in the State both for ornament and fruit that its original locations in the State are difficult to make out. Unquestionably its mass occurrence was in Knox, Gibson and Posey counties. Its occurrence in Vigo County, reported by W. S. Blatchley, is unquestionable, but may be regarded as exceptional. A record of its occurrence in Fountain County recorded in Indiana Geological Report, 1882. Vol. 11, p. 122, is of doubtful validity and may safely be disregarded. The remaining citation from Jefferson County (A. H. Young) is based upon a single tree located in the river bottoms near Hanover. It stood alone in a large bottom land, otherwise destitute of trees. It was near a dwelling and for this reason and because of its small size, it is a fair inference that it was a cultivated form. It is to my mind certain that the pecan as a member of the Indiana tree flora is mainly confined to the three southwestern counties, but extending in greatly reduced numbers northward as far as Vigo County. In any event, the northeastern limit of the species is reached in these locations. The trees are smaller than those of more southern and western localities and according to Deam, "only about one-fourth of the native trees ever bear fruit and only about one out of every ten trees is a profitable nut-bearing tree."² This southern and western form probably entered our area at about the same time and under the same physical conditions as the Bald cypress.

The Over Cup Oak (*Quercus lyrata* Walt.) according to Sargent occurs in "river swamps and small deep depressions on rich bottom lands, usually wet throughout the year."³ This would explain the close restriction of the

¹Eleventh Annual Report. Indiana State Board of Forestry, 1911. p. 108.

²Deam:—Op. cit. p. 138.

³Sargent. Manual of the Trees of North America. p. 269.

species to Gibson, Knox and Posey counties. The species is mainly coastal, occurring along the Atlantic and Gulf coasts, finding its way into the interior along the valley of the Mississippi and its tributaries. The occurrence of this coastal species, and of others which might be cited, in regions so markedly interior, suggests the thought that they entered our flora at the time when the great northward arm of the Gulf of Mexico practically divided the area of the United States into distinct eastern and western regions. A comparison of the boundaries of this arm of the Gulf of Mexico with the distribution of many species of plants gives striking support to such an explanation. In Indiana, the species is not ordinarily separated from the Burr oak (*Quercus macrocarpa* Michx., which it closely resembles.

The Cow or Basket Oak (*Quercus Michauxii*, Nuttall) so far as our records go occurs only in Knox and Gibson counties, where it is found in low, rich bottom lands. This restriction of area may be due in part to the close resemblance in size and habit of this species to the Swamp White Oak (*Quercus bicolor* Willdenow) and a consequent failure to distinguish it from that species. This oak, also, is more or less coastal in its mass distribution, finding its way into the interior along the valley of the Mississippi; while the Indiana localities represent its northeastern limit the species maintains its normal size and habit. Its ability to maintain a foothold in localities often covered with water serves to explain its persistence in the counties named.

The Spanish Oak (*Quercus falcata* Michaux) is reported definitely from Knox, Gibson and Posey counties, with an additional citation from Fountain County (Brown) which the writer has not had opportunity to verify. Some difficulty arises in this case because of the question of the validity of the species. According to Sargent *Quercus falcata* is separable into two species, *Quercus digitata* Sudworth, and *Quercus pagodaefolia* Ashe. According to Gray's Manual⁴ *Q. falcata* is found on dry or sandy soil; upon the authority of Sargent *Q. digitata* grows in similar soils on dry hills; while *Q. pagodaefolia* occurs on rich bottom lands and alluvial banks of streams. In Indiana the form in question "is usually found in low ground, associated with *Quercus bicolor*, *Q. palustris*, *Q. Schneckii*, *Q. stellata* and *Q. velutina*. The whole of the township is low."⁵ The habitat as well as considerations of distribution would seem to indicate the species to be *Q. pagodaefolia* Ashe instead of *Q. falcata* Michaux. In the Proceedings of

⁴Gray's New Manual of Botany. Seventh Edition. p. 343.

⁵Deam. Op. cit. 207.

the Indiana Academy of Science, Vol. 11, page 142, and Vol. 12, page 299, I have previously expressed the opinion that the species in question is *pagodæfolia*. If this is true, this southern form came into our area along the valley of the Mississippi, and a species which is a large and valuable "timber tree in the river-swamps of the Yazoo basin, Mississippi, and of Eastern Arkansas"⁶ might reasonably be expected in the low, wet bottom lands of the lower Wabash Valley. The Indiana station represents the northeastern limit of the species, a fact reflected in its sparing occurrence and reduced size. Deam's collection of 1915 show the occurrence of this species in Jefferson County.

The Yellow Hackleberry (*Celtis Mississippiensis* Bosc.) is frequent or common along streams and in the lowlands of Gibson, Knox and Posey counties. It is a southern and western form, reaching a height of from 60 to 80 feet and a diameter of from 2 to 3 feet in the basin of the lower Ohio River. In Indiana which represents its northern limit the tree "is inclined to grow scrubby and crooked." (Deam.) It is medium sized, rarely exceeding a diameter of 18 inches. Its occurrence within our area is easily explained, since the counties named are not especially far removed from the center of its maximum development both as to size and numbers. It is a little difficult, however, to explain why it has not spread more widely in the State.

The Southern Thorn (*Cratægus viridis* Linnæus) is distinctly southern and somewhat western in its mass distribution, reaching its greatest abundance and largest size in western Louisiana and eastern Texas. It is found along stream borders and the margins of swamps in moist soils, doubtless finding its way into our area when such conditions were practically continuous.

The Shining Thorn (*Cratægus nitida* (Eng.) Sargent) is said in Sargent's "Trees of North America" to occur on the "bottoms of the Mississippi River in Illinois opposite the city of St. Louis." The species occurs in rich bottom lands in both Gibson and Posey counties in fair abundance as a small tree, from 20 to 30 feet high and with a broad and handsome crown. As a result of the recent work in the segregation of species in the genus *Cratægus* it is practically impossible to form any definite notion as to the range of any particular form. Much field work will be necessary before we can determine just what species of this puzzling genus are members of our flora. No opinion is expressed, therefore, regarding the source from

⁶Sargent. Op. Cit. 245.

which *C. nitida* came into the State. There is as much reason for regarding the Illinois station as the westward extension of the Indiana station as the reverse.

The Water Locust or Thorn Tree (*Gléditsia aquatica* Marshall) is found in a few localities in Gibson, Knox and Posey counties in sloughs and cypress swamps. This is a northern and eastern extension of a definitely southern species which must have entered our flora at a time when the swamp areas of the river bottoms were practically continuous and which has been able to maintain itself only in occasional deep river swamps within our boundaries. In Indiana the species is both rare and local and one which will, in all probability, soon disappear.

One of the Hollies (*Ilex decidua* Walter) occurs occasionally in the three southwestern counties, being invariably restricted to the borders of ponds and sloughs near water courses. Although at times it forms fairly dense thickets, it rarely, in our region, reaches tree size. The distribution as given in Sargent's "Trees of North America," p. 618, is significant in this connection: "Borders of streams and swamps in low moist soil; southern Virginia to western Florida in the region between the eastern base of the Appalachian Mountains and the neighborhood of the coast, and through the Gulf States to the valley of the Colorado River, Texas, and through Arkansas and Missouri to southern Illinois; usually shrubby east of the Mississippi River and only arborescent in Missouri, southern Arkansas and eastern Texas." It is merely another instance in which an essentially coastal form has found its way deep into the interior. When considered in connection with other cases, some of which have been cited, the conclusion is almost inevitable—that the only adequate explanation is to be found in relating it to the northward stretching arm of the Gulf of Mexico.

Pond-bush (*Forstiera acuminata* (Michaux) Poiret) is another species strictly limited to Gibson, Knox and Posey counties where "it is found in swamps, on the borders of ponds and on low river banks. It is very tolerant of shade and is frequently found growing in a thick stand of tall trees."⁷ The Indiana stations represent the extreme northeastern limit of this species, which extends westward to Missouri and south to Texas. In Indiana it is ordinarily a shrub, at times forming almost impenetrable thickets. It is impossible to determine from the data at hand as to whether this is a western or southern form. In either case its habitat,

⁷Deam. Op. Cit. 342.

"low, wet river banks and swamps" suggest the same reasons for its occurrence in the Indiana flora as has been suggested for the preceding species.

Hardy Catalpa (*Catalpa speciosa* Warder) is a tree of the borders of streams and ponds and of fertile often flooded bottom lands. According to Sargent it is probably found in its greatest abundance and of the largest size in southern Illinois and Indiana, extending to western Kentucky and Tennessee, southeastern Missouri and northeastern Arkansas. In Indiana it is confined to Knox, Gibson, Posey and Vigo counties as a member of the original forests. Its occurrence in other counties is due to its widespread cultivation for post material or for ornamental purposes. Deam says,⁸ "In Indiana it was found along the valley of the Ohio River as far east as Rockport and in the valley of the Wabash as far north as Vigo County. The mass of its distribution was west of a line connecting Terre Haute and Rockport." The citations given are, however, all that can be considered as verified. In the catalpa we evidently have another case in which the distribution is easily explained if it is related to a northward extension of the Gulf or to a condition of flooded rivers.

The Swell-butt Ash (*Fraxinus Michauxii* Britton) usually grows in low grounds which are inundated for several months during the year. As its common name indicates the swollen base is characteristic of this species. It has been collected in Gibson, Posey and Marion counties by C. C. Deam. The Gibson and Posey County stations represent normal conditions for the species; the Marion County collection is in different case. The tree, which was of medium size, was growing in moist soil by the roadside. The known care and accuracy of Mr. Deam preclude any doubt as to the determination, so that the occurrence of the species in this station must be referred to some accidental means of transportation or to what is perhaps more probable, the incorrect labelling of material furnished by some nursery for roadside planting. As a component member of our native forests the species is undoubtedly confined to Gibson and Posey counties. As this is a species but recently segregated its distribution is not yet thoroughly known. It, however, is known to range from New York to North Carolina and Louisiana and west to Missouri.

This is very evidently another case of a species of coastal distribution with a seeming extension well into the interior.

If we summarize these thirteen species, peculiar to our southwestern counties we find them all to be swamp forms or those growing in bottom

⁸Deam. Op. Cit. pg. 347.

lands frequently inundated during the year, or in low moist localities. We find that the larger part of them, in their mass distribution follow the swamps of the Atlantic or Gulf coast, or of both. It is very evident also that the extension of range northward must have occurred when similar physical conditions existed; that is, either at the time the Gulf of Mexico stretched an arm far into the north, or if a later date is preferred in the time of the flooded rivers and lakes of the Champlain period. Occasional means of transportation may serve to explain occasional cases, but where species become component parts of a forest in a region apparently remote from their mass distribution a different explanation must be sought.

Six species, so far as the records go, are confined to Lake, Porter and Laporte counties or to some one of them. In this region, also, extremely skillful and persistent work has been done by Rev. E. J. Hill, a fact which should be taken into account. The species peculiar to this region are the following:

- Pinus Banksiana* Lambert.
- Thuja occidentalis* Linnaeus.
- Betula populifolia* Marshall.
- Betula papyrifera* Marshall.
- Alnus incana* (Linnaeus) Muenchhausen.
- Celtis pumila* (Muhlenberg) Pursh.

The Jack or Scrub Pine (*Pinus Banksiana* Lambert) occurs in Lake and Porter counties, where it is fairly common on the sand dunes bordering Lake Michigan. The general range of this species is decidedly northern, the Indiana stations representing in all probability its extreme southern limit. In our area it is an undersized, rather shrubby form, maintaining itself with difficulty. The continuity of waterways is the evident explanation of the occurrence of this species in the Indiana tree flora.

The Arbor-Vitæ or White Cedar (*Thuja occidentalis* Linnaeus) apparently occurs native only in Lake County. This characteristic species of northern swamp regions is found only in cold swamps of our area. There seems no good reason why it should not be found in similar situations in other counties bordering Lake Michigan. The form has been so extensively planted for windbreaks and for ornament that many incorrect citations are on record. Its presence as a member of our flora is evidently referable to continuous waterways furnished by the Great Lakes.

The Gray or White Birch (*Betula populifolia* Marshall) is found in

scant numbers and of small size in Lake, Porter and Laporte counties. This typical northern species may also be regarded as one that has retained a foothold in isolated localities after the recession of the shores of Lake Michigan and the disappearance of the bordering swamps. The reported occurrence of the species in Tippecanoe County (Golden) "in sparing numbers along the Wabash River" demands further study. The well-known difficulty of discriminating the species of *Betula*, due to seasonal and age changes in appearance and habits, suggests that a closer study may prove the reference an error.

The Paper or Canoe Birch (*Betula papyrifera* Marshall) is found in Lake and Porter counties, always being reported as rare and of small size. This is another species definitely northern in its mass distribution, the Indiana stations standing as its extreme southern limit. It is probably another of the species which entered our area in the time of flooded rivers and lakes of the Champlain but one which has been able to maintain a precarious foothold up to the present time. Its early disappearance from the tree flora seems inevitable.

The Tag or Speckled Alder (*Alnus incana* (Linnæus) Muenchhausen) is found in Lake and Porter counties between dunes near to the lakes. This is the common alder found in swamps and on the borders of streams further north. It has been able to maintain itself in our area in greater numbers and with less reduction of size than any other one of these extreme northern species.

A dwarf shrubby form of Hackberry (*Celtis pumila* (Muhlenberg) Pursh) is included. In both Gray and Sargent the form is regarded as a variety of *C. occidentalis* Linnæus. Its general range is in the South Atlantic States ranging westward to Missouri, Colorado, Utah and Nevada. It has been reported only from Lake County near the Calumet River at Millers. Its occurrence in such a restricted locality is rather puzzling and as yet no satisfactory explanation has been reached. The form in the greater part of its range occurs on rocky banks of streams—a condition not even approximated at its Indiana station. The temptation to regard it as an ecologic variant of the very common *Celtis occidentalis* is almost irresistible.

Of the six species just discussed, five are definitely northern in their mass distribution. Their presence as members of our flora is very evidently referable to the continuity of waterways existing during the

Champlain period. The more difficult problem is the explanation of their persistence.

A consideration of a few other species will serve to emphasize the point in mind.

The Larch or Tamarack (*Larix laricina* (DuRoi) Koch) is found in Porter, Marshall, Kosciusko, Noble, Steuben, DeKalb and Blackford. An examination of the older shore lines of Lake Michigan gives a sufficient explanation. Even the Blackford County citation, which seems well to the south, is made clear when the ancient bay of Lake Michigan extending southward through Allen County in the neighborhood of Fort Wayne is recalled.

Thus also the eastern Peach-leaved Willow (*Salix amygdaloides* Anderson) found in Lake and Kosciusko finds ready interpretation, as does also the case of the Wild Red Cherry (*Prunus Pennsylvanica* Linnæus fil.) occurring in Lake, Porter and Kosciusko counties.

Any one who maps some of the more widely ranging species of the State will be immediately impressed by the close relations existing between the distribution of the species and the course of waterways. In some instances the distribution follows a single waterway, in others it seems to follow not merely the main stream but also all of the tributaries. Indeed, by far the most striking feature in the series of one hundred twenty-six maps is the definite way in which this relationship stands out. The most cursory inspection of the maps reveals it and serves to suggest at least a possible causal relation.

In the opinion of the writer the occurrence of given species in widely separated localities without intervening stations will be found to be due to the existence at some time in the past of practically continuous waterways connecting these now separated localities. Further, that such connections, in so far as the region under consideration is concerned, are mainly to be found in the Champlain period, although perhaps in some cases this connection was furnished by the northward stretching arm of the Gulf of Mexico. If the shore lines of streams and lakes of the Champlain period could be drawn upon our present maps many of our problems in Phytogeography would solve themselves. In confirmation of this view is the dominating influence of continuous waterways or of streams in the distribution of species clearly shown by any careful study of present range extensions.

In the main, widely ranging species, at least among trees, do not have

this wide range because of any perfection of seed dissemination, nor because of occasional means of transportation. It has more probably been brought about by a former connection of these separate regions, making them practically continuous. Such practical continuity may have been secured by means of the flooded rivers and swollen lakes of the Champlain period. This is not offered as a solution of all of the problems of plant distribution, but in the firm belief that in many of these problems the solution is to be sought in former physical conditions.

It was the original intention that the present paper should also include a discussion of the species restricted to the southern tiers of counties, but the time required to work our former shore lines and water-levels for that region was too great. The preliminary work, however, seemed to indicate confirmation of the conclusions indicated in this paper.

CORRECTIONS TO THE LISTS OF MOSSES OF MONROE COUNTY, INDIANA, I AND II.

MILDRED NOTHNAGEL AND F. L. PICKETT.

At the winter meetings of the Indiana Academy of Science in 1912 and 1913, the authors presented lists of the Mosses of Monroe County, Indiana. Since the publications of these lists, several corrections in identification have been made and these will be listed below. The correct names will appear first, the former name in parentheses after the accession number.

Corrections to the 1912 list of Monroe County Mosses, I.

Fissidens minutulus Sull. (44.) (*F. bryiodes*.)

Determined by G. B. Kaiser.

Orthotrichum lescurii Aust. (3). (*O. porteri*.)

Determined by G. B. Kaiser.

Brachythecium campestre B. & S. (16). (*B. plumosum*.)

Determined by G. B. Kaiser.

Eurynchium hians (L.) B. & S. (17). (*Brachythecium rutabulum*.)

Determined by G. B. Kaiser.

Plagiothecium geophilum Aust. (22). (*P. deplanatum*.)

Determined by G. B. Kaiser.

Corrections to the 1913 list of Monroe County Mosses:

Barbula unguiculata (Huds.) Hedw. (71). (*Didymodon ru-beillus*.)

Determined by G. B. Kaiser.

Funaria hygrometrica (L.) Sibth. (79). (*F. flavicans*.)

Determined by G. B. Kaiser.

Aphanorhegma serratum Sull. (122). (*Physcomitrium immersum*.)

Determined by G. B. Kaiser.

Amblystegium orthocladon (Pb.) Lindb. (98). (*A. fluviatile*.)

Determined by G. B. Kaiser.

Amblystegium varium (Hedw.) Lindb. (93). (*A. serpens*.)

Determined by G. B. Kaiser.

Hypnum curvifolium Hedw. (85). (*H. fertile*.)

Determined by G. B. Kaiser.

H. curvifolium Hedw. (88). (*H. pratense*.)

Determined by G. B. Kaiser.

Indiana University Botanical Laboratory.

THE MOSSES OF MONROE COUNTY, INDIANA, III.

F. L. PICKETT AND MILDRED NOTHNAGEL.

The collection and classification of the mosses of Monroe County, as has been reported by the authors at the winter meetings of the Indiana Academy of Science for 1912 and 1913, was again resumed in the spring of 1914. The present list will include a few forms from Hamilton and Lake counties of Indiana, and Berrien County of Michigan.

The 1914 list of mosses includes twenty-nine new species, from eleven families and twenty-six genera, of which are representatives of one family and thirteen genera not reported in former reports. Material has been prepared, as described in former lists and left in the herbarium of the Botany Department at Indiana University with notes as to time, place, and habitat of the collection as well as condition of specimens. The entire collection in the herbarium now contains specimens, many of them in duplicate from different collections of ninety-five species and five varieties, representing fifty-three genera and seventeen families of the Bryales. In this report, as in former ones, the numbers within the parentheses after the name of the specimen is the accession number.

A great number of the specimens collected this year have been sent to A. J. Grout of Brooklyn, N. Y., or to G. B. Kaiser of Germantown, Pa., for verification or identification and the credit of such will be given in the list.

Order SPHAGNALES.

Family *Sphagnaceae*.

Sphagnum acutifolium Ehrh. (189). Determined by G. B. Kaiser.

In swamps south of Gary, Lake County, Ind. Sterile.

Order BRYALES.

Suborder NEMATOBONTEE.

Family *Buxbaumiacae*.

Buxbaumia aphylla L. (134).

Matures spores from December to April. On sandy soil in moist ravine.

I. C. pond. Fruiting.

Webera sessilis (Schmid.) Lindb. (145).

Matures spores in spring and early summer. On clay in moist ravine,
I. U. pond. Fruiting.

Suborder ARTHRODONTÆÆ.

Family *Dicranaceæ*.

Ceratodon purpureus (L.) Brid. (163, 199). Verified by G. B. Kaiser.
Matures in spring and early summer. On sandy soil about Gary, Lake
County, Ind. Common. Fruiting.

Dicranella heteromalla var. *orthocarpa*. (Hedw.) E. G. B. (136). Ver-
ified by A. J. Grout.

Capsules mature in November and December. On clay on hillside
northeast of Bloomington, Ind. Fruiting.

Dicranum fulvum Hook. (151). Determined by G. B. Kaiser.

Yellowish green mats on beech tree on south side of Griffy Creek,
north of Bloomington, Ind. Sterile.

Family *Grimmiaceæ*.

Hedwigia albicans (Web.) Lindb. (142, 160). Verified by G. B.
Kaiser.

Matures spores in spring. All material collected was sterile. In dark
brownish green mats on decaying wood near I. U. pond.

Grimmia doniana Smith. (130). Determined by G. B. Kaiser.

In thin hoary patches on limestone, I. U. Campus.

Family *Tortulaceæ*.

Tortella cæspitosa (Schwaegr.) Limp. (291). Verified by G. B. Kaiser.

Matures spores in spring. Specimen robust and fruiting. Found on
moist rocks above spring, Clear Creek, Ind., Monroe County.

Family *Orthotrichaceæ*.

Drummondia clavellata. Hook. (120, 135).

Matures spores in spring and summer. Found fruiting on bark of tree.
Hamilton County and Monroe County, Ind.

Orthotrichum porteri Aust. (123). Determined by Mrs. E. G. Britton.

Matures spores in the spring. In black-green mats upon limestone.
Fruiting. Rare.

Orthotrichum schimperii Hamm. (173). Determined by G. B. Kaiser.

Matures spores in spring. In dark green cushions upon trees and
stumps in dry places.

Family *Bartramiaceæ*.

Philonotis fontana (L.) Brid. (186). Determined by G. B. Kaiser.

Matures spores in May and June. Found on moist rocks above spring, Clear Creek, Monroe County, Ind.

Family *Bryaceae*.

Mniobryum albicans (Wahl.) Limp. (178). Determined by G. B. Kaiser.

In thick green mats or tufts on rocks near water, northwest of Harrodsburg, Monroe County, Ind. Sterile. Fruits infrequently.

Bryum caespiticium. L. (172). Determined by G. B. Kaiser.

Spores mature in spring. On a decayed log in tufts. Bloomington, Ind. Common.

Family *Leskeaceae*.

Anomodon minor (P. B.) Förn. (133). Verified by G. B. Kaiser.

Sterile specimen. Found on retaining wall along north pike, Bloomington, Ind.

Family *Hypnaceae*.

Rhytidium rugosum (L.) Kindb. (141). Verified by G. B. Kaiser.

Sterile form. In dense light green masses on stone, Harrodsburg, Monroe County, Ind.

Brachythecium rivulare B. S. (132, 185). Verified by G. B. Kaiser.

Sterile, never found fruiting here. In very moist places or in water.

Bryhnia graminicolor (Brid.) Grout. (154). Determined by A. J. Grout. Sterile form, rarely found fruiting. On moist clay and stones just above water line in Griffy Creek, northeast of Bloomington, Ind.

B. Novae-angliae (S. & L.) Grout. (197). Determined by G. B. Kaiser. On edge of swamp, north of Gary, Ind., in loose light green mats. Sterile.

Cirrophyllum bosci (Schwaegr.) Grout. (138, 181, 182). Verified by G. B. Kaiser.

Spores mature in autumn. Found on soil in moist ravine in bright green mats. Sterile form.

Cratoneuron filicinum (L.) Roth. (179). Determined by G. B. Kaiser. Sterile specimen found in bluish green tufts just above water line in shaded spring.

Amblystegium irriguum (Wils.) B. HS. (180). Verified by G. B. Kaiser.

Sterile specimen. Found just above water line in shaded spring in deep olive to blackish green mats on rocks and clay.

- Hypnum molluscum* Hedw. (166). Determined by G. B. Kaiser.
Sterile form. On soil on wooded hillside, I. U. dam.
- H. patientiae* Lindb. (176). Verified by G. B. Kaiser.
Matures spores in spring. Fruiting. Yellowish green mats on decayed
log, near Buchanan, Berrien County, Mich.
- Plagiothecium deplanatum* (Sch.) Grout. (170). Determined by G. B.
Kaiser.
Sterile. On limestone in open ravine north of Bloomington, Ind.
- Amblystegiella adnata*. (Hedw.) Nichols. (170, 184). Determined by
G. B. Kaiser.
Matures in summer. Specimen sterile. Found on limestone and on
base of tree in dark green mats, about Bloomington, Ind.
Family *Leucodontaceae*.
- Leucodon brachypus* Brid. (152).
Matures spores in winter. Specimen sterile. In blackish green mats
resembling *Hedwigia albicans*. Found on log north of Bloomington,
Ind.
- Forsstroemia trichomitrium* var. *immersum* (Sulliv.) Lindb. (204).
Verified by G. B. Kaiser.
Matures spores in summer. Fruiting. On tree in ravine southeast of
I. U. dam.
- Indiana University Botanical Laboratory.*

A NEW ENEMY OF THE BLACK LOCUST.

GLENN CULBERTSON.

During the latter part of June and during July of 1914, the leaves of the greater number of the locust trees in Switzerland, Jefferson, Clark, and Floyd counties, of southern Indiana, were observed to be losing their greenish appearance, and upon closer examination the chlorophyll of the leaflets was found to have been largely consumed. The foliage appeared as though dried up as a result of a severe drouth. Here and there individual trees, at a distance from groves, were unaffected, but the trees of practically every grove, at least among the hills of the Ohio and tributaries were seriously affected. So evident was this that the brown and sere appearance of the groves was noticeable as far as they could be seen.

The infected trees were found to be alive with a small beetle, which Professor Enders of Purdue classified as *Chalcopus dorsalis* of Blatchley's "Coleoptera of Indiana": "This beetle is from 6 to 6.5 mm. long, wedge shaped and rather broad, bluish black, thorax red, with black sutural stripe. Found throughout the State, but much more abundant in the southern counties. Occurs on flowers of black locust, in the leaves of which the larvæ mine. Hibernates beneath the locust bark."

On striking the trees the beetles could be heard falling to the ground by the scores. They could be seen in large numbers on the foliage, as many as five were counted on a single leaflet.

The eggs of this beetle are deposited late in April or early in May, and by the 20th of May the young larvae are at work between the coverings of the leaflets, destroying all the inside portion. In some cases several larvae may be seen at work within a leaflet. When mature the larvae stop eating and remain enclosed within the leaf coverings until metamorphosis is completed, when they emerge, usually about the 20th or 25th of June, and for several days feed upon the upper leaf surface of any green foliage that may remain.

The writer is of the opinion that many of the locust groves in the northeastern part of Jefferson County were badly infected with this beetle

during the summer of 1913, and possibly previous to that, as several groves of that region were observed with foliage that seemed to be drying up, as though injured by a serious drouth. The infected area seems to be rapidly increasing, and the annual defoliation of the trees must in time prove a serious injury, since the foliage on the majority of the infected trees was practically useless by the first of July and in many cases even earlier. What the favoring circumstances have been that have caused this remarkable increase in the numbers of this insect is at present a mere conjecture. It may be that the unusual heat and drouth of May, 1914, and of the summer of 1913, may have caused their rapid multiplication, or that the relatively rapid increase of locust trees, their favorite food supply and breeding place, has augmented their numbers.

What the future may bring no one knows, but if this beetle continues in as great numbers in succeeding years, they will prove a very serious menace to locust groves, and the fence and telephone post industry of southern Indiana. Judging from the undoubted rapid increase in the past, the future is not promising.

As a remedy Professor Enders and others recommend spraying with arsenate of lead or other arsenical compounds. This no doubt would be in a measure effective, if applied within a few days after the emergence of the mature beetle June 25th to July 5th, and could be done on level or moderately level ground, but since the tens of thousands of volunteer locusts are on slopes so steep that they are almost inaccessible, it would prove a difficult task indeed to get at them with a spraying outfit. It is not probable that the pest, if it proves to be a serious one, will be eradicated in that way. It is very difficult to get farmers to spray orchards, much less locust trees scattered far and wide over rough, hilly land. It is to be hoped that an efficient remedy may be provided, for, if not, this defoliator, in addition to the borer, will probably end the locust industry.

A NEW LEAF SPOT OF VIOLA CUCULLATA.

H. W. ANDERSON.

A leaf spot on *Viola cucullata* has been prevalent in Indiana and neighboring States for a number of years. It is especially noticeable during the early spring months. Collections of leaf spots on this host in different parts of the country have been made from time to time and have been filed away in the herbarium without being classified or wrongly labeled as the *Phyllosticta* leaf spot. A careful examination during the past year has revealed the fact that this particular leaf spot is caused by a *Colletotrichum* which has never been described as occurring on this species. Since this disease is widespread it was thought worth while to make a careful study of the causative organism.

Of the violets which occur in this region only *Viola cucullata* has been found to be attacked by this particular fungus. It is interesting to note that while this species is attacked *V. palmata* is apparently immune. *V. cucullata* was formerly considered a variety of *V. palmata* and the immunity of the latter emphasizes the specific difference. However, only a limited number of plants of *V. palmata* have been observed and these in a region where the disease was not common on the other species.

Cultivated violets have been examined only in the local greenhouses. It is probable, however, that all cultivated species are immune, otherwise the disease would have been observed and reported by those especially interested in violet diseases.

Macroscopical Appearance.—The fungus produces a typical leaf spot. The earliest indication of infection is a pale area with a definite dark green border. Later the area in the center of the spot dies, turns white, grey or light brown, a dark brown ring appears about the edge, forming a definite, regular spot. At an early stage the acervuli appear as dark brown dots on the lighter central area. They are irregularly arranged and occur on both sides of the leaf. The dark color of the acervuli is due, in part, to the numerous setæ. Later the center of the spot becomes very thin and papery and may fall out, thus giving a shot hole effect. When badly in-

fect the spots are occasionally confluent. Usually there are only a few spots on a leaf.

Etiology.—The fungus gives rise to numerous acervuli, which are dark brown or black, irregularly scattered, varying greatly in size (50-200 microns). They are beset with dark brown setæ. The setæ are numerous, arising from any part of the acervulus, 1-4 septate, dark brown, sharp-pointed, straight or slightly curved above the base. The base is usually bent or curved in various ways.

The spores are hyaline, non-septate, slightly curved averaging 4.5x25 microns. They are borne on short, hyaline conidiophores. In some cases at germination a delicate septum was observed in the middle of the spore. This is by no means always present.

Nomenclature.—There has been, in the past, some confusion in regard to the limits of the genera *Colletotrichum*, *Vermicularia*, *Volutella* and *Chaetostroma*. This has been due, largely, to the lack of care exercised by investigators when species of these genera have been studied. Sections carefully made clear up generic confusion very easily. The fungus described above is undoubtedly a *Colletotrichum* since there is no pycnidium, the spores being borne on short conidiophores in a setose acervulus.

In 1899, Dr. Ralph Smith¹ described a leaf spot of pansy caused by a *Colletotrichum*. The type material of this fungus has been examined and found to differ from the *Colletotrichum* under discussion in the size and shape of the spores, the shape of the setæ and the character of the spot produced.

Dr. Peck² in 1878 described a leaf spot of *Viola rotundifolia* as follows:

"*Vermicularia concentrica* Peck and Clinton n. sp. Perithecia small, black, beset with straight, rigid bristles, concentrically placed on arid, orbicular spots; spores oblong, slightly curved, pointed at each end, colorless, .0008'-0.001' long.

"Living leaves of *Trillium erythrocarpum*. Pine Valley, Clinton, July:

"The tissues at length fall out from the affected spot, leaving apertures through the leaf. The perithecia are less regularly disposed near the center of the spots. Judge Clinton also sends a variety on the leaves of *Viola rotundifolia* in which the concentric arrangement of the perithecia is not at all preceptible, but I detect no other difference."

(¹) Botanical Gazette. 27: 203-204, Mar. 1899.

(²) Report of the N. Y. State Botanist 1878. 29th Annual Report of the N. Y. State Museum of Natural History, pps. 47-48.

In his "Sylloge", Saccardo³ changed the specific name of the fungus to *Peckii*. His nomenclature is as follows:

"*Vermicularia Peckii* Sacc.

V. Concentrica Peck, 29th Ann. Rep. N. Y. State Mus. Nat. Hist. 47-48, 1878. (Not Lev., Ann. Soc. Nat. 66. 1845.)

"*V. Peckii* var. *Viola rotundifolia* Sacc."

Through the kindness of Dr. H. D. House, State Botanist of New York, I was able to secure authentic specimens, collected and determined by Dr. Peck subsequent to his description of the fungus. These specimens were from *Trillium erythrocarpum* Michx. (*T. undulatum* Willd.) and from *Viola rotundifolia*. A careful examination of these specimens, together with some recently collected material from Dr. House, was made. Some of the spots were embedded in paraffin, sectioned and stained. From these examinations it was concluded that the fungus occurring on *Viola rotundifolia* was not a *Vermicularia* but was identical with the *Colleotrichum* occurring on *V. cucullata*. The spore measurements and general characters of the acervulus, setæ and conidiophores of the fungus on *Trillium* were also identical but, as stated by Dr. Peck, the acervuli of the former occur in definite concentric circles in the spot, while in the latter no such arrangement is noticeable. Whether or not the species on *Viola cucullata* is identical with the one on *Trillium* can only be determined by cross-inoculation. Up to the present time the author has not had an opportunity to complete his investigations along this line. It would be unusual, however, to have a fungus of this type parasitic on hosts so widely separated as *Trillium* and *Viola*. The identity of the fungi on the two violet species can hardly be questioned. The nomenclature of the fungus on *Viola rotundifolia* is so awkward and incorrect that a change should be made. However, this is not advisable until the relationship to the fungus occurring on *Trillium* is definitely settled.

Life History of the Fungus.—The field observations of this fungus have been limited to a single year. The disease appears at a very early period in the spring on leaves that have evidently lived over winter. The earliest collections were made in the first week in April before the plants had time to develop leaves. It is probable therefore that the fungus lives over winter on the old leaves of the plant, although it has not been observed during the winter months.

(3) Saccardo, P., Sylloge Fungorum. 3: 232. 1884.

The spread of the disease in the early spring is evident from an observation of an infected plant. A single plant often has at first only one or two infected spots. As the new leaves develop they are seen to be free from the disease, but a few days after a damp period these leaves are badly infected. This disease is, in many respects, an early spring one, for the most luxurious growth of this species of violet is during April and May and the fungus seems to thrive best during cold, damp weather. At no time do the number of spots become so great as to kill the leaf. The damage done by this fungus is evidently negligible so far as this species is concerned.

The fungus spores are probably distributed through the usual agents. They are produced in great numbers on both sides of the leaf and they germinate readily in tap or rain water.

Artificial Inoculations.—A detailed report of the experiments on inoculations made would be out of place at this time. Summarizing these, it was found that infection by sprayed spores on the leaves was readily brought about if the plants were kept very moist for several days. The number of infections were small on each plant. Cross-inoculations on *V. pubescens* and *V. striata* gave negative results.

Cultural Characters.—The fungus is easily isolated by poured plates of the spores. Cultures were made on various media, but a description of the development on dextrose-potato agar will be sufficient to show the general character of the growth. The mycelium is at first hyaline, mostly confined to a thick growth on the surface of the agar, with scanty aerial mycelium. At the edge of the advancing mycelium the agar takes on a pink and then a red color which is very striking. Later the mycelium darkens and in some cases within four days conidia are to be found. The darkening of the mycelium continues and within two weeks a heavy, black, stroma-like crust is formed over the surface of the agar. On this stroma hyaline, gelatinous areas of conidia are developed.

Conclusion.—This preliminary report on a leaf spot of *Viola cucullata* is given in order to stimulate others to examine the plants of this species during the coming spring, thus establishing more definitely its range and determining whether or not it is confined to this one host. An effort will be made by the writer to determine the relationship of this species to the one occurring on *Trillium* and on *V. rotundifolia*.

OAT SMUT IN INDIANA.

F. J. PIPAL.

In the winter of 1914, the writer, representing the Botanical Department of the Indiana Agricultural Experiment Station, conducted, in cooperation with the Extension Department of Purdue University and the county agricultural agents, a series of meetings at which demonstrations were given of the formaldehyde treatment of seed oats and potatoes. The meetings were held in Benton, Lake, Porter, Jasper, Pulaski, Laporte, Elkhart, Grant, Madison, Randolph, Clinton and Montgomery counties, which are among the largest oat-growing counties in the State. According to the report of the last census these twelve counties raised over thirty-two per cent., in acreage, of the entire oat crop of the State. It may be of interest, therefore, to report some facts resulting from these meetings, since they furnish fairly reliable data as to the oat smut situation throughout the State.

A most striking thing has come to light in connection with this campaign. It has been learned that out of 3,168 persons reached through the meetings less than a dozen farmers previous to that time had ever used the formaldehyde treatment for their seed oats. The use of formaldehyde as a general disinfectant and a specific fungicide for potato scab was originated, about eighteen years ago, in the Botanical Department of the Indiana Agricultural Experiment Station, by Dr. J. C. Arthur. It was then applied as a disinfectant for oat smut and the stinking smut of wheat by Professor H. L. Bolley, formerly assistant to Dr. Arthur. It remains to the present date the simplest, cheapest and most effective seed grain disinfectant in use. A large majority of the farmers of the State, however, evidently have not, for some reason, taken advantage of this discovery, and still allow the smut disease to reduce the oat yield by several million bushels every year.

One of the reasons for this neglect evidently is the fact that most farmers do not fully realize the extent to which the oat smut occurs in their crops. About thirty years ago, Dr. Arthur, then a botanist for the

New York Agricultural Experiment Station, at Geneva, demonstrated that oat smut is not readily visible to the unpracticed eye unless ten or more per cent. of the crop is affected. The smutted stalks are, to a large extent, considerably shorter than the sound stalks, and can not usually be seen except upon close examination of the field. And again, most of the smutted masses are blown away by harvest time and only bare stalks remain, leaving nothing conspicuous to indicate the amount of damage done.

Dr. Arthur found nine and one-half per cent. of smutted plants in fields at the Geneva Station in which the presence of smut could scarcely be detected without close examination. In the third annual report of the New York Experiment Station he remarks in this connection: "The appearance of smut as one passed through the fields was no greater than is usually to be seen in any part of the country, * * * and the result of the count * * * is as much a surprise to the writer as it will doubtless be to others."

E. S. Goff, of the Wisconsin Experiment Station, estimated the loss from oat smut in that State, in 1896, at about nine per cent.

Bowman and Burnett, of the Iowa Experiment Station, found, in 1907, an average of seven and nine-tenths per cent. of smutted heads in twenty fields examined.

Kellerman and Swingle estimated, in 1888 and 1889, that Kansas lost annually over eleven per cent. of the oat crop from smut.

In bulletin No. 37, of the Ohio Experiment Station, J. F. Hickman says: "In passing through one of our oat fields last summer I observed what seemed to be a smutted head here and there, but upon careful examination I found more than seven per cent. of this variety smutted."

In order to demonstrate the importance and the value of the formaldehyde treatment as effectively as possible the county agents in a number of counties made arrangements with some of the farmers to treat all their seed oats except a small portion to serve as a check on the treatment. It may be well to state here that most of the farmers who agreed to make the tests were under the impression that their oat crops of the previous seasons were comparatively free from smut. The test fields were distributed over Madison, Grant, Laporte, Pulaski and Benton counties.

When the oats headed out the county agents counted the smutted heads and figured out the percentage of smut on the treated and untreated plots. In Madison County, where the writer assisted the county agent,

Mr. W. R. Butler, in this work, counts of smut were also made in several fields where no treatment had been tried.

The following table shows the results of the tests as reported by the county agents.

TABLE 1.

RESULTS OF THE FORMALDEHYDE TREATMENT FOR OAT SMUT ON TEST FIELDS
IN FOUR COUNTIES.

COUNTY.	Number of Test Fields.	Reported by.	Average Per Cent. of Smut on Treated Fields.	Average Per Cent. of Smut on Untreated Fields.
Madison.....	15	W. R. Butler...	.3	12.0
Grant.....	4	O. Crane.....	.8	13.0
Pulaski.....	7	W. V. Kell.....	.1	11.7
Benton.....	6	J. W. McFarland	.2	11.0
Average.....			.3	11.9

In Laporte County, Mr. L. B. Clore, the county agricultural agent, arranged for a test of the formaldehyde treatment on the county poor farm. The manager of the farm was very reluctant at first to make the test, claiming that there never had been any oat smut on the farm. When the smut was counted, however, it was found that fifty-two per cent. of the crop was smutted on the untreated field and only about one per cent. on the treated field.

The results demonstrated to the farmers beyond any doubt the value of the treatment. The treated fields were practically free from smut, while those not treated had, individually, from one to fifty-two per cent. of the crop destroyed by the disease. Three fields in Madison County had thirty or more per cent. of smutted heads, and one field in Pulaski County showed a loss of forty-five per cent. The average percentage reported from Madison, Grant, Pulaski and Benton counties correspond closely, indicating that the prevalence of oat smut is fairly uniform throughout the sections these counties represent.

In addition to the data obtained from the test fields further reports on the prevalence of oat smut were received from seven counties as shown in the next table. The figures submitted in these reports were secured by the county agricultural agents and other men who made, in most cases, careful observations and counts of oat smut in their respective counties.

TABLE 2.

AVERAGE PERCENTAGE OF SMUT FOUND IN THE OAT CROP OF 1914 IN SEVEN COUNTIES.

COUNTY.	Reported by.	Average Per Cent. of Smut.
Randolph.....	C. A. Mahan.....	15
Whitley.....	W. C. Dilts.....	01
Montgomery.....	R. A. Chitty.....	15
Starke.....	H. R. Smalley.....	10
Lake.....	S. J. Craig.....	20
Gibson.....	H. F. Buk.....	10
Jefferson.....	G. Culbeertson.....	15
Average.....		13.5

As shown in the table the average per cent. of smut reported from the seven counties corresponds closely with the average figures from the counties mentioned in Table 1. Leaving out the report from Laporte County, which can not be considered representative owing to the high per cent. of smut obtained in the single test, the grand average for the counties under consideration is practically 13 per cent. This no doubt is a fairly accurate figure representing the loss from oat smut in the State. It corresponds closely with the estimate of Dr. Arthur who placed the loss in the State, figured from general observations, from eight to twelve per cent.

According to the crop statistics, compiled by the United States Department of Agriculture, Indiana devotes annually about 1,735,000 acres (average of 1909 to 1913 seasons) to the production of oats. The average yield for the State has been about thirty bushels per acre. It may be considered, therefore, that the average annual production of oats in Indiana is, in round figures, about 52,000,000 bushels. Considering that smut destroys about thirteen per cent. of the crop the above yield represents

only eighty-seven per cent. of the full crop. Figuring on this basis the annual loss from oat smut amounts to 7,770,113 bushels. This is more than the total yield of Benton, Allen and Tippecanoe, three of the largest oat-growing counties in the State. At the average price of oats of thirty-five cents per bushel the loss in cash value equals \$2,719,539. The cost of treating seed oats with the formaldehyde solution would be about two cents per acre, or \$34.00 for all seed sown in the State. The net profit resulting from the treatment would be, therefore, considerably over two and one-half million dollars. To gain this amount every year by practicing the treatment is certainly worth the effort, and practical instructions and demonstrations along this line in all oat growing sections of the State are highly desirable.

The formaldehyde treatment of seeds oats, as recommended by the Indiana Agricultural Experiment Station, is briefly as follows:

Spread out the seed on a floor and sprinkle with a solution of one pint of 40 per cent. formaldehyde to 50 gallons of water until thoroughly moist. Shovel over repeatedly to distribute the moisture evenly, then shovel into a pile and cover with sacks or canvas for at least two hours. The seed may be sown as soon as dry enough to run without clogging the drill. If to be kept longer than one day, grain should be dried as rapidly as possible by spreading in a thin layer and stirring occasionally with a rake. Avoid reinoculating with smut from smutting sacks or bins after treatment. One gallon of the solution will treat a little more than one bushel of oats.

In order to facilitate the work of treating the grain, machines have been invented which much simplify the labor and enable one to treat large quantities of grain in a comparatively short time. Several types of these machines are now on the market selling for twenty dollars or more each.

If total destruction of the oat crop in three counties occurred, it would arouse the farmers of the State to action. Why should not the loss of more than two and one-half million dollars distributed over the State do so? If all farmers in Benton County treated their seeds oats they would save enough in one season to build at least eight township schoolhouses, each costing not less than twelve thousand dollars. And then they could save enough every year to pay the salaries of all their school teachers. Many other counties in the oat-growing sections could do equally well.

In some townships the formaldehyde treatment would save the farmers enough money to pay for the building of miles of stone roads. Should not these facts stir the farmers to some concerted action by which they would banish the smut disease from the State? The grain treatment is simple, cheap and easy of application. It is up to the oat growers in the State to make up their minds and do the right thing. A man in Madison County, on whose farm a test of the formaldehyde treatment was made this spring, was very much pleased with the results, and he said in substance: "Why it's a very simple thing. There's very little work connected with the treatment and the cost can almost be disregarded. I treated my seed for less than twenty cents. I wonder why I haven't been practicing it long before."

PLANTS NEW OR RARE TO INDIANA. NO. V.

 CHAS. C. DEAM.

Specimens of the species reported are deposited in my herbarium under the numbers indicated. The Gramineæ were determined by A. S. Hitchcock; the Carices by K. K. Mackenzie; the *Juncus* by H. H. Bartlett; and the Antennariæ by M. L. Fernald.

Panicum Werneri Scribn.

Floyd County, June 8, 1913. No. 13,256. In a sterile white and black oak woods on the "knobs" about one mile west of New Albany.

Muhlenbergia foliosa Trin.

Grant County, September 4, 1914. No. 15,279. Low border of the lake located about five miles northeast of Fairmount.

Whitley County, August 23, 1914. No. 14,562. Low border on the west side of Round Lake.

The only reference to this species occurring in the State is in Rhodora, Vol. 9:19:1907, in an article by Lamson-Scribner on "Notes on Muhlenbergia", in which he refers to "No. 68, by H. B. Dorner from Indiana."

Apera spicaventi (L.) Beauv.

Orange County, August 1, 1914. No. 15,561. Frequent over an area of five or six acres about one mile west of Leipsic. Reported by Prof. M. L. Fisher of Purdue University.

Bromus arvensis L.

Jefferson County, May 28, 1911. No. 8,486. In a woods along the roadside about one-half mile south of North Madison.

Bromus hordeaceus L.

Laporte County, May 28, 1913. No. 13,031. Frequent along the roadside east of the water works at Michigan City.

Carex Leavenworthii Dewey.

Shelby County, June 8, 1913. No. 13,193. Collected by Mrs. Chas. C. Deam in a dry woods one and a half miles west of Morristown.

Vermillion County, May 8, 1910. No. 5,819. In dry soil in a wooded ravine about one mile northwest of Hillsdale.

Carex louisiana Bailey.

Gibson County, June 10, 1913. No. 13,297. In a low flat woods on the east side of Foote's pond.

Carex projecta Mack.

Hendricks County, July 13, 1913. No. 13,677. Collected by Mrs. Chas. C. Deam in a swamp about five miles northwest of Danville.

Carex suberecta (Olney) Britt.

Noble County, June 26, 1914. No. 14,350. Low border of Engle Lake.

Shelby County, June 8, 1913. No. 13,204. Collected by Mrs. Chas. C. Deam in the Milburn Swamp about one mile west of Morristown.

Tipton County, May 24, 1913. No. 12,909. Collected by Mrs. Chas. C. Deam in a low place along the Lake Erie Railroad about two miles west of Goldsmith.

Carex picta Steud.

Brown County, May 21, 1910. No. 6,412. Somewhat frequent on the oak ridges between Helmsburg and Nashville. This sedge has the habit of growing in bunches. When established for some time it advances in the form of a circle, the center of which is bare. Areas two feet in diameter have been noted with a ten-inch bare center.

Morgan County, August 12, 1913. No. 13,958. Collected by Mrs. Chas. C. Deam on a wooded hillside about one and a half miles northeast of Martinsville.

Gray's text-book of botany gives this species as occurring in a ravine near Bloomington upon the authority of Dudley.

Carex impressa (S. H. Wright) Mack.

Allen County, June 14, 1914. No. 14,258. Creek bottom about seven miles south of Ft. Wayne.

Gibson County, June 9, 1913. No. 13,339. In a low flat woods about four miles west of Patoka.

Grant County, June 16, 1907. No. 2,057. Along a creek about two miles northeast of Van Buren.

Hamilton County, May 19, 1912. No. 10,549. Collected by Mrs. Chas. C. Deam in a swamp near Carmel.

Hancock County, June 2, 1912. No. 10,892. Collected by Mrs. Chas. C. Deam in a ditch along the C. H. & D. Railroad near Reedsville.

Huntington County, July 4, 1907. No. 2,141. In a wet woods about two miles north of Buckeye.

Johnson County, June 8, 1912. No. 11,088. Collected by Mrs. Chas. C. Deam along Young's Creek south of Franklin.

Marion County, May 30, 1911. No. 8,522. Collected by Mrs. Chas. C. Deam in a ditch near Irvington.

Posey County, May 24, 1911. No. 8,350. In a wet woods near Goose Pond.

Wells County, May 22, 1908. No. 3,040. Abundant in a wet woods just south of Bluffton.

Carex atherodes Spreng.

Tipton County, July 9, 1913. No. 13,632. Collected by Mrs. Chas. C. Deam in a prairie habitat along the Lake Erie Railroad about two miles west of Goldsmith.

Juncus brachycarpus Engelm.

Crawford County, July 13, 1899. In a valley near Wyandotte Cave.

Marshall County, July 2, 1911. No. 9,021. In a ditch along the railroad about one mile south of Culver.

Alsine graminea (L.) Britt.

Laporte County, May 22, 1910. No. 6,440. On the bank of an open ditch through the prairie just west of the State Prison.

Ranunculus cymbalistes Greene.

Floyd County, April 20, 1913. No. 12,565. Type locality about one-half mile south of the Southern Railroad on the top of the first wooded ridge west of New Albany. Associated with *Pinus virginiana* Mill. and *Quercus prinus* L.

Physocarpus opulifolius var. *intermedius* (Rydb.) Rob.

Starke County, September 1, 1914. No. 15,152. A large colony on the south side of Bass Lake.

Sanguisorbia minor Scop.

Lawrence County. Introduced in grass seed in a field about 80 rods from the boundaries of Orange and Washington counties. Reported by Prof. M. L. Fisher of Purdue University.

Meibomia illinoensis (Gray) Kuntze.

Benton County, July 31, 1912. No. 11,839. Roadside about four miles north of Fowler.

Jasper County, July 30, 1912. No. 11,789. Roadside about four miles west of Remington.

Lagrange County, August 29, 1914. No. 14,939. Roadside about three miles northeast of Howe. Associated with *Kuhnia eupatoroides*, and *Drymocallis agrimonioides*.

Lake County, July 29, 1912. No. 11,750. Roadside about four miles southeast of Cedar Lake.

Marshall County, July 2, 1911. No. 8,995. Open wooded hillside about a half mile south of Culver.

Steuben County, August 11, 1903. In woods near Gage Lake. In the same county on a gravelly wooded hill on the east side of Hog-back Lake in 1906. No. 1,265.

Hartmannia speciosa (Nutt.) Small.

Floyd County, June 8, 1913. No. 13,260. Abundant in an alfalfa field and on both sides of an adjoining road on the "knobs" west of New Albany, just beyond the terminal of the traction line to Silver Hills.

Solidago hispida Muhl.

Steuben County, September 20, 1914. No. 15,504. In white oak woods and along a wooded roadside on the east side of James Lake.

Antennaria neodioica Greene.

Wells County, May 28, 1914. No. 14,190. Growing with white oak and *Antennaria fallax* on the high clay bank of the Wabash River about one mile below Vera Cruz.

Antennaria Parlinii Fernald.

Brown County, May 21, 1910. No. 5,985. Frequent on sterile wooded hills between Helmsburg and Nashville.

Clark County, May 8, 1912. No. 10,496. In sterile white and black oak woods on the Forest Reserve.

Decatur County, May 5, 1912. No. 10,479. Abundant on the top of the sterile wooded bank of Flat Rock River about half a mile above St. Paul.

Jennings County, April 28, 1912. No. 10,429. Top of the high bank of the Muscatatuck River between Vernon and North Vernon.

Martin County, May 20, 1913. No. 12,872. On top of a wooded bluff along White River about two miles above Shoals.

Montgomery County, May 16, 1913. No. 12,750. In a white oak woods along Sugar Creek near the "Shades".

Ripley County, May 19, 1912. No. 10,570. On dry clay bank of Baupst Creek about one mile west of Morris.

St. Joseph County, May 29, 1914. No. 14,204. Dry woods about five miles southwest of South Bend.

Antennaria Parlinii var. *arnoglossa* (Greene) Fernald.

Vermillion County, May 8, 1910. No. 5,837. Top of the bank of a wooded ravine about a mile and a half northwest of Hillsdale.

Antennaria solitaria Rydb.

Floyd County, April 20, 1913. No. 12,552. Growing under a beech tree in the "knobs" about a mile west of New Albany, also on the top of one of the "knobs" with *Quercus Prinus*.

SOME PECULIARITIES IN SPIROGYRA DUBIA.




PAUL WEATHERWAX.

A form of *Spirogyra* found on the campus of Indiana University early in the spring of 1913 has shown, in its natural habitat, as well as when subjected to new physiological conditions, some phenomena of growth that are not only irregular for *Spirogyra* but also seem to be confined rather closely to the one species.

The plant does not agree exactly with the description of any species given in the literature available, but it conforms fairly well with the description given by Wolle (1) and also the one given by Collins (2) for *S. dubia* Kg. This species, according to these descriptions, has two spirals, or "more rarely three", and the fruiting cell is described as being slightly inflated. The plant observed here had regularly three chloroplasts, and the fruiting cells were not at all swollen. Wood (3) notes this same difference in the sporangial cell and suggests other variations but concludes that these characteristics are not sufficiently different to justify the description of a new species. A form showing a physiological peculiarity similar to one shown by this plant, and probably from the same general location, is identified by Pickett (4) as *S. clongata* (Berk.) Kg.

When first found the plant formed a thin, green coating on a piece of rusty sheet iron lying in running water. Most of the filaments were only one to three cells in length and were probably developing from zygotes, but the striking thing noted was the highly differentiated basal cells (Fig. 3) by which the filaments were attached to the mud on the iron, and, in many cases apparently to the rough surface of the iron itself.

Conditions were favorable for rapid growth, and ten days later the filaments were three or four inches in length and composed of many cells, but still as firmly attached as would have been filaments of *Cladophora* of the same size. The root-like basal cells had grown very much longer and had assumed a variety of peculiar shapes. Their walls had thickened, and their contents were just beginning to show signs of decomposition (Fig. 5).



A quantity of the material was put into a shallow dish of distilled water and placed in a north window in an attempt to cause it to conjugate, but indications of an unhealthy condition soon became apparent. As a first indication of this condition the chlorophyll bands became more slender and the pyrenoids very prominent. Soon after this the filaments began to break up by the decay of some of the cells, so that but few segments could be found that were more than seven or eight cells in length, and the majority of them were made up of but one or two cells. In the meantime these cells that seemed to have greater vitality began to develop a number of branches as shown in Figs. 1 and 2. The cytoplasm usually followed out into these branches, often taking a loop or an end of a chloroplast with it. In many filaments that showed the peculiarity the branches all arose from the same end of the cells, suggesting a continuation of the condition of base and apex that was made so evident by the highly specialized basal cells of the younger plants. So far as was observed, the branches always remained continuous with the cells from which they arose, no new cells being cut off on branches. Filaments of other species of *Spirogyra* often broke up into segments on being put into similar conditions, but no branching was observed.

The decay of the plant was probably started, in some instances at least, and very evidently greatly aided, as soon as the vitality of the alga had been slightly impaired, by the growth of a fungus, *Aphanomyces phycophilus* De Bary, one of the few parasitic forms of the *Saprolegniaceae*, which has already been described (5). Other species of *Spirogyra* seemed immune to the attack of this fungus.

In some conjugating material of this same species of *Spirogyra* similar physiological peculiarities were noted. This latter material had been preserved for class use, and the exact locality of its collection is not known, but it was probably found in the same general locality as was that first mentioned. The filaments of this material showed also, but in a less marked degree, the same unhealthy condition. Some typical branches found are shown in Fig. 4. These branches seemed to serve as "hold-fasts" for attaching the filament to other filaments of *Spirogyra* or probably to other things in the water.

Any attempt to get at the meaning of the branches found in either instance must maintain a degree of consistency with two or three prominent points observed. The branching described is associated with a pathological condition and is characteristic of this to a more marked

degree than of any other species of *Spirogyra* that was tested. Since the phenomenon was observed once in a physiological condition that had been made favorable for conjugation and again where conjugation was actually taking place, it was at first thought that the branches were exaggerated attempts at conjugation, and, in some instances, this may have been the case. But the filaments were usually close enough together that such long tubes would not have been necessary, and no actual union of gametes as a result of any such activity was at any time observed. Moreover, the filaments shown in Fig. 4 illustrate a condition noted in two or three cases, where filaments having mature zygotes in some of their cells were attached by these branches to others also containing zygotes. The filament shown in this figure as holding to another by means of the foot-like branch was a long one and had at another place mature zygotes that had been formed as a result of conjugation with some other filament. If these branches were modified conjugating tubes, a relation of this sort would be out of harmony with the tendency toward bisexuality that is usually exhibited by the plant.

Indiana University,

Bloomington, Indiana.

LITERATURE CITED:

- (1) Wolle, Rev. Francis. Fresh water algæ of the United States. 1887.
- (2) Collins, Frank S. Green algæ of North America.
Tufts College Studies, Vol. II, No. 3. 1909.
- (3) Wood, Horatio C. Jr. A contribution to the history of the fresh water algæ of North America.
Smithsonian Contributions, No. 241, Vol. 19. 1872.
- (4) Pickett, F. L. A case of changed polarity in *Spirogyra elongata*.
Bul. Tor. Bot. Club, Vol. 39. 1912.
- (5) Weatherwax, Paul. Aphanomyces phycophilus De Bary.
Proc. Ind. Acad. of Sc. 1913.

REPORT ON CORN POLLINATION IV. (FINAL).

M. L. FISHER.

The work under this head has been reported in the 1908, 1910, and 1911 proceedings. The reports have dealt mostly with cross-pollinating with pollen from a variety of a different color or race. One of these crosses—sweet, male, and Reid's Yellow Dent, female—was selected to be carried out to the end to see if a new variety could be produced.

In the third year two types of sweet corn were distinguishable, one a large ear with whitish kernels and white cobs like the original Stowell's Evergreen, and the other, a smaller ear with yellowish kernels and red cobs. These two types were planted the season of 1911, but through poor management no seed was saved. Enough ears were obtained to see that the types were fairly well fixed. Old seed was used in 1912 and hand pollinations were made on each type. A few good ears of each kind were obtained. The kernel and cob characters came true to the original selections. Upon being cooked as roasting ears, both types were found to be of excellent quality—the yellow kernel and red cob type being slightly sweeter. The mature ears showed some dent kernels, but not many. A good quantity of seed was obtained. It may be said further that the yellow-kernel red-cob type was somewhat earlier than the white-kernel white-cob type, the latter inclining to be late.

In the season of 1913 three plantings were made, one in the writer's garden, another in the garden of the foreman of the Station Experimental plats, and the third in the trial gardens of D. M. Ferry & Co., Detroit, Mich. In the two garden trials, the white-kernel white-cob type was used on account of its promising greater prolificacy. The corn in the writer's garden was almost ruined by a hail storm, and that in the foreman's garden was somewhat injured. Such ears as were obtained for use as roasting ears were declared to be of superior quality. The writer saved no seed, the foreman was able to save a good quantity and planted again in 1914.

The corn planted on the plats of Ferry & Co., was reported on as follows:

"The salient features of our reports are to the effect that neither of your selections seems as yet well enough fixed in type to be ready for presentation. Both show a large percentage of reversion to plain parent stock. They are both late and half of the ears in our trial were irregularly and poorly filled. Quality seems excellent, but the color of the red cob shows badly in cooking.

From a seedsman's standpoint we do not believe the strains to be as yet of any value."

The foreman mentioned above planted a small patch the past season (1914), but drouth and hot winds ruined the entire planting. However, enough seed remains for another planting and a replenishment is hoped for. Although results have been somewhat discouraging, it is believed that a successful and fixed variety may yet be developed.

STOMATA OF TRILLIUM NIVALE.

F. M. ANDREWS.

Gümbel¹ was the first to make known the presence of twin stomata. Since that time Pfitzer² and others have shown the presence of stomata in groups of two or more on the leaves of various plants. In *Saxifraga sarmantosa* stomata are arranged "in circular groups"³ in considerable number. In various species of *Begonias*, as De Bary states⁴ in *Begonia manicata*, *B. spathulata*, *B. Dregei* and *B. heracleifolia* two or more stomata are arranged over one respiratory cavity.

This occasional grouping of the stomata in certain plants is even more strikingly shown in *Trillium nivale*. The stomata are often found on the leaves in pairs over a common respiratory cavity, but frequently in numbers up to ten or more. In opening and closing they act just as a single stoma does.

The presence of more than one stoma over a common respiratory cavity is also shown on the sepals and petals. Figure 1 shows part of a sepal of *Trillium nivale* in which the stomata are in pairs in one case and in threes in another case over a common respiratory cavity. These arise from the successive division of a common mother cell. The stomata on the sepals and petals are frequently lateral or diagonal as regards one another, but in every case their origin from one mother cell is the same. The arrangement in groups of as many as ten or more over one respiratory cavity on the sepals or petals is also met with.

Figure 2 shows a case, taken from the outside of a sepal, where only one guard cell, A, is fully formed. There is only a remnant of a second guard cell, B. The same thing has also been observed on the inside of the petal.

¹Gümbel. *Jahr. für wiss. Bot.* Bd. 7, p. 551.

²Pfitzer, E. *Jahr für wiss. Bot.* Bd. 7, pp. 532-560.

³*Treviranus. Verm. Schritten*, IV. 30. Quoted from DeBary, A. *Comparative Anatomy of Phanerogams and Ferns.* 1884 p. 47.

⁴Vivani. Quoted from DeBary as above.

In Figure 3 a pair of stomata is shown in which only three guard cells were formed. In this case the apertures are closed by the movement of the two outer guard cells only.

These deviations from the general order, position and number of stomata in *Trillium nivale* also obtains, but to a less extent, in other species of the genus *Trillium*. It is also in keeping with other deviations, for which the genus *Trillium* is noted, such as monstrosities in the leaves themselves and in the parts of the flower. Interesting questions are connected with the twin, triple and grouped stomata of *Trillium nivale* and other plants as to their complete development, the real causes of their arrangement and their physiological reactions.




FIG. 1. *Trillium Nivale*. Stomata from outside of sepal showing double and triple groups over one respiratory cavity. x ca. 100.




FIG. 2. *Trillium Nivale*. Stoma with only one fully developed guard-cell. x 45.




FIG. 3. *Trillium Nivale*. Stomata from sepal. x 45.

THE PRIMROSE-LEAVED VIOLET IN WHITE COUNTY.

LOUIS F. HEIMLICH.

On the afternoon of June 2d last, I started out from Reynolds, Indiana, to add a few specimens to my herbarium. Following the Pennsylvania lines east, I noticed many familiar plants, among them being *Viola lanceolata* L., the lance-leaved violet, growing in great abundance in the wild grass along the right of way. A goodly number of *Viola sagittata* Ait. were often close neighbors to *lanceolata*.

After digging up a few very fine specimens of these two violets I climbed over the fence and went up on one of the sand ridges so characteristic of the country about Reynolds. I had crossed this area very often and knew that *Viola pedata* L., the bird-foot violet, grew here. Only a few of their flowers remained, the seedpods on some being already of good size.

Passing over the edge of this oak-forested sand ridge, I descended into what was once a swamp area. The soil suddenly becomes mucky, mixed with sand and late decayed leaves. Here, to my surprise, I discovered a violet which I had never seen before. I knew it was a violet. I felt sure of that, and so remarked to my two companions. We looked and found more of them nearby. They spread from the lower limit of the sand ridge out to a little beyond a fence-row, covering an area of about 40 by 125 feet.

These violets, which I later found to be *Viola primulifolia* L., the primrose-leaved violet, seemed to seek the shade. Most of them grew along the fence-row in wild grass, together with some weeds and small brush. Those which grew out in the open short grass were low spreading and less succulent plants. The season for flowering was about over and the cleistogamous capsules were making their appearance.

Viola primulifolia L., varies from about 5 cm. to 20 cm. in height, bearing from a few to a dozen or more primrose-like leaves. The plant is stemless, the leaves rising from a medium sized rootstock or runner. The lower leaves are oval to almost round. The upper, larger leaves are ovate.

Viola Primulifolia L.

—DRAWN FROM A
MOUNTED SPECIMEN—

$\frac{3}{4}$
NATURAL SIZE



J. F. Heimlich 14
-Nov. 17-1914-

some oblong-ovate, with acute apexes and long tapering bases. Some of the leaves may be somewhat sub-cordate. Both lower and higher leaves are more or less crenate, mostly glabrous, with slight pubescence along the midrib and the edges of the petiole.

The flowers are small, the largest being hardly a centimeter in width when full blown. The petals are white with several purple stripes on their inner surfaces. The pedicels may be 15 cm. or 18 cm. long or as long as the leaves. One or two small bract-like leaves appear about half way up the pedicels. If two are present they may be opposite each other or a little apart. The various botanies do not mention these structures nor does the




illustration in Britton and Brown show them on this particular violet, although they are shown on a number of the other accaulescent species.


Stolens are common and give off vigorous new plants. The small reddish-brown seeds are scarcely more than a millimeter in length.

Viola primulifolia L., occurs in the eastern United States. Its range is given in the various botanies as in moist or almost dry soil from New Brunswick to Florida along the coast. In so far as I know it has not been reported in any other place in this botanical region. How it happened in White County, Indiana, or how general its distribution here is, I do not know. Mr. Deam, who later visited the place with me, thought it might occur also farther north in the State. I have looked for it, specifically,



SOME DETAILS CONCERNING VIOLA PRIMULIFOLIA L.



1. Characteristic leaf - showing pubescence. NATURAL SIZE.



2 & 3 Lower Leaves - showing variation in outline NATURAL SIZE.




4 Seed-pod bursted. TWICE NATURAL SIZE.



8. Seeds - color: reddish-brown. FOUR TIMES NATURAL SIZE.



5. Pedicel and flower. Flower - white with purple stripes. NATURAL SIZE.



9. Stolon - showing formation of a new plant.

L.F. Heimlich 44
-NOV-17-1914-

several times since in different localities where I thought it might occur, but so far have not found it. It was rather abundant in the very limited area during the early summer, but the drouth which followed killed every one of them. At present I have several potted specimens growing nicely along with several other species of violets.

This paper does not aim to give a complete study of the plant. It is intended to give merely a brief account and description of the plant as I know it, and to announce definitely that *Viola primulifolia* L., does occur in Indiana.

CONTINUOUS RUST PROPAGATION WITHOUT SEXUAL REPRODUCTION.

C. A. LUDWIG.

With the demonstration of heterœcism in the rusts, the teliospore came to be looked upon as primarily a resting spore, for heterœcism was first proved for the black rust of grasses, in which the teliospore is a true resting spore. It was therefore believed by implication, indeed often stated, that the teliospore is the means of carrying rust fungi over unfavorable weather conditions and is especially equipped structurally for that function. The other spore forms and the mycelium, except in the comparatively rare cases of a mycelium diffused through and perennial in the tissues of the host, were not supposed to be able to survive such adverse conditions. Later when modern cytological methods were applied to the rusts, it was found, as some leading uredinologists had already suspected in a rather vague way, that the essential feature of the teliospore is that it is the structure in which is begun the series of nuclear phenomena which close the sporophytic stage and precede the gametophytic stage with its resulting sexual fusions. With this latter idea goes the rather common belief that no type of life, plant or animal, except perhaps the very lowest, can long maintain a high degree of vigor without at least occasional sexual fusions. It is this idea that has given rise to the belief held in many places that if all the barberry bushes could be destroyed, the black rust of cereals would not be able to maintain itself more than a few years. Thus we have superimposed one upon the other in the minds of many men these two ideas: (1) that the teliospore is necessary to the continued existence of the rust because it is the means of passing the winter or other unfavorable season, and (2) that it is necessary to the vigor of the fungus because it is the structure in which are initiated those changes which culminate in sexual fusions and without which such fusions would not take place. The continued prevalence early in the season of the grain rusts at great distances from any possible æcia of the species, however, led some investigators to doubt the validity of the beliefs just recorded. In consequence, a number

of investigations were made which showed, as suspected, that the rusts in question do have other means than the teliospore of surviving the winter and that in all likelihood they are able to propagate themselves indefinitely in an asexual manner, and that without serious impairment of vigor. In this paper it is proposed to give the results of this work and to present some field data, chiefly from Indiana, which was made available to the writer by his access to the Arthur herbarium. This data goes to indicate that what is true of the grain rusts is equally true of some others which because of their lack of economic importance have thus far escaped this sort of investigation.

DURATION OF VIABILITY OF UREDINIOSPORES.

Before entering on the discussion proper, however, it seems best to treat here two points which have a bearing on what is to follow; namely, the duration of the viability of urediniospores and the distance to which they may be blown and produce infection. As to the duration of germinability in the urediniospore, more work has been done with the grain rusts perhaps than with any other. In the case of *Puccinia graminis* Hungerford reports¹ finding germinable urediniospores at Madison, Wis.; on timothy in October, November, December, January, and March; but it does not seem to be at all certain that the spores used on the last named date were wintered spores. On the contrary, it is altogether possible that they were but recently produced. Mercer,² however, was not able to find germinable urediniospores of the same rust on the same host during the winter in North Dakota; and Eriksson and Henning,³ as the result of several experiments with the rust on different hosts, came to the conclusion that the fungus does not pass the winter in the uredinial stage in Sweden. They also came to the same conclusion in regard to *P. glumarum*.⁴ Bolley⁵ reported the germination of 8-15% of urediniospores of *P. graminis* after twenty-one days in dry air in August. The same investigator has shown⁶ that the urediniospores of the leaf rust of wheat can be used for successful infection material after thirty days' exposure to the outside air in July, while Freeman and Johnson consider it possible for urediniospores of *P. graminis* and *P. rubigo-vera* to survive the winter in Minnesota, North

¹Phytopathology 4:337-338. 1914.

²l. c. 20-22.

³Die Getreideroste 38-47. 1896.

⁴l. c. 153-159.

⁵Centralblatt für Bakt. Par. und Infekt. 4: 893. 1898.

⁶Agricultural Science 5:263. 1891.

Dakota, and Wisconsin.⁷ Fromme has shown⁸ that the period of viability in the leaf rust of oats, *P. coronata*, may be as extended as eighty-four days. The spores in this case were stored dry in a gelatin capsule. Marshall Ward⁹ succeeded in securing germination of unrediniospores of *P. dispersa* which had been for sixty-one days in dry air in the diffused light from a north window. Thus, while urediniospores are capable of germinating as soon as mature, they are capable under proper conditions of maintaining their viability for a period of two to three months and probably more. One of these conditions seems to be dryness. Probably the most common limiting factor to long life of urediniospores in nature is a combination of warmth and moisture. In such a case germination probably takes place, thus of course forestalling any long duration of life in the spore. There seems to be no good reason, however, if germination can be avoided, why urediniospores might not survive the winter. That coldness of weather does not destroy ability to germinate is attested by the fact that a number of investigators—Hungerford,¹⁰ Ward,¹¹ Carleton,¹² and others—have collected viable urediniospores of various grass rusts during the winter months.

DISTANCE WHICH WIND BLOWN SPORES TRAVEL AND PRODUCE INFECTION.

It seems to be a fact, although from the nature of the case not fully proved, that urediniospores of the rusts may travel long distances by the wind and produce infection. This, of course, is to be expected of a structure which can stand drying for so long a time and is so light in weight. Klebahn¹³ calls attention to a sand storm which arose in northern Africa and progressed northward over Europe, transporting various mineral particles to various places in Europe. He adds that without doubt the rust spores, which are much lighter than the mineral particles, are much easier transported by air currents. Under the circumstances they would remain suspended much longer and could be carried at least as far and perhaps farther than the mineral particles. The same investigator¹⁴ proved the presence of spores in the air high off the ground by constructing traps for the spores and exposing them in trees and on buildings. He was able in

⁷Bur. Plant Ind. Bull. 216:52. 1911.

⁸Bull. Torrey Club 40:518. 1913.

⁹Ann. Mycol. 1:138. 1903.

¹⁰Phytopathology 4:337-338. 1914.

¹¹Ann. Myc. 1:132. 1903.

¹²Div. Veg. Phys. and Path. Bull. 16:44. 1899.

¹³Die wirtswechselnden Rostpilze 68. 1904.

¹⁴I. c. 68.

this way to capture a large number of spores, many of which were urediniospores of the rusts. It is clear, therefore, that urediniospores may be carried many miles in the air; and there is apparently no reason for thinking that they may not then start infection. Doubtless they do start infection and in this way produce a number of isolated areas of rusted plants.

As for epidemics, however, there is evidence that they are not caused by spores brought from a great distance. One item of this evidence is furnished by Pritchard,¹⁵ who was unable to capture any urediniospores of *P. graminis* at Fargo, North Dakota, in a series of trials extending over a period of nearly a month, until the rust was common on wheat in the neighborhood. Pritchard's trap was a dish set on a five-foot post and containing a little water. Another item is the fact that in the spring, as described by Christman,¹⁶ the earliest outbreak of leaf rust on wheat is a rather heavy one on the old, wintered leaves. The old leaves then die and a period of approximately four weeks follows in which little or no rust can be found, followed by another free infection. If the epidemic were initiated by spores blown from far away, we would not expect a heavy infection early in the spring. Instead we should expect a very light early infection increasing gradually to the full epidemic later on. Further evidence of the limited distance to which spores blow in anything like sufficient quantity to produce an epidemic is found in recorded observations where it was possible to know the source of the spores producing the infection. Perhaps the species which are most limited in this respect so far as observations recorded up to the present time go are *Uromyces andropogonis*, *Puccinia andropogonis*, and *P. ellisiana*, all of which, according to observations made by Long¹⁷ are limited to an area within six feet distant from the æcia. These observations are in harmony with our own general experience in collecting rusts near Lafayette and elsewhere. The æciospores cause free infection to a distance from their source depending upon the height of the source from the ground, but the distance is almost always small. As to the distance to which æciospores of *P. graminis* blow in considerable quantity there are published observations by Arthur,¹⁸ Pritchard,¹⁹ Mercer,²⁰

¹⁵Bot. Gaz. 52:183. 1911.

¹⁶Trans. Wis. Acad. Sci. 15:106. 1905.

¹⁷Jour. Agr. Res. 2:303-304. 1914. Phytopathology 5:170. 1912.

¹⁸The æcidium as a device to restore vigor to the fungus. Proc. 23rd meeting Soc. Prom. Agr. Sci. p. 3. 1903.

¹⁹Bot. Gaz. 52:178. 1911.

²⁰Phytopathology 4:22. 1914.

and others. The greatest of these values is that reported by Pritchard, whose statement is here quoted: "The rust was abundant within 25 yards of the barberry bushes, but practically disappeared at a distance of 60 yards. The most persistent searching was required to discover a single pustule beyond 80 yards." Another observation by Pritchard²¹ indicated that urediniospores of *P. graminis* are carried only short distances in sufficient number to cause an epidemic. We have the published record, however, of rust spores having been blown as much as a mile and producing infection; and, strange as it may seem, the spores in question are the smallest, most delicate ones in the life cycle of the rusts if we omit the non-functional pycniospores, namely, the basidiospores. E. T. Bartholomew²² gives a table which shows that 59.1% of the leaves on apple trees near cedars were infected with rust. A quarter of a mile away it was 55.4% and a mile away it was 6.5%. All this does not show, of course, that rust spores are not carried by the wind for long distances in a vigorous condition, but it does show that the distance for abundant infection from any spore producing center is not great.

With this as a basis it should be possible to obtain an idea of the maximum distance a rust might be expected to progress in a season. The greatest distance recorded above is one mile, but those spores would doubtless travel a mile and a half farther (or two and a half miles) and produce infection. As a factor of safety, let us double this value; and as a further factor of safety, let us double this latter value. This gives us ten miles. A rust generation, according to Freeman and Johnson,²³ takes eight to twelve days, and more in cold, bad weather; and our own results at the Purdue Experiment Station agree very well with those figures. Assuming, then, ten days for a rust generation, ten miles of migration per generation, a growing season from the middle of April to the middle of October, approximately 180 days, and good weather with no interruption to the growth of the fungus, we should expect it to migrate for a distance of 180 miles. This value will be used presently in comparing the telial distribution of some of the rusts with their possible aërial distribution.

PROPAGATION OF INDIVIDUAL SPECIES.

The black rust of grasses was among the first to be observed living, and apparently thriving, at long distances from any of its aëria. There

²¹Bot. Gaz. 52:184 1911.

²²Phytopathology 2:255-6. 1912.

²³Bur. Plant Ind. Bull. 216:45. 1911.

seems now not to be the least doubt but that it can pass the winter in the uredinial stage; in fact, McAlpine²⁴ claims that in Australia the æcia of the rust do not exist and that the rust will not infect the barberry. As to the exact method of wintering there is some difference of opinion. The contention by Pritchard²⁵ that teliospores or mycelium in the seed grain have something to do with its propagation in wheat is different from all the others in the suggestiveness that a sexual process may be involved even in the absence of the æcium. All other theories which have been advanced assume a strictly asexual propagation, and probably Pritchard's theory does not really assume otherwise either. Perhaps the most famous theory is Henning's now discredited mycoplasma theory. The real means by which the rust passes the winter is probably mycelium in the leaves of the host plant. The presence of this mycelium during the winter months has been shown by Hungerford²⁶ and by Johnson²⁷ in the leaves of timothy.

That the leaf rust of wheat is carried through the winter in the same way is shown by the findings of Bolley,²⁸ Carleton,²⁹ and Christman.⁶⁰ This method of carrying the fungus over accounts satisfactorily for the heavy early infection, followed by a period of little or no infection, which is in turn followed by the epidemic proper. The old leaves, which are infected from the autumn, carry the first epidemic and then die, the mycelium, of course, dying with them. In the meantime the new leaves have been infected; and in about four weeks, which as has been shown by Freeman and Johnson,³¹ and by Christman,³² is the approximate incubation period for that time of year, the uredinial stage breaks out freely on them.

Aside from the work with the grain rusts, not much has been done in the way of determining the method of passing the winter by rusts in regions remote from their æcia. Carleton,³³ however, states that *Puccinia montanensis* on *Elymus* winters in the uredinial stage, and calls attention to the situation with regard to the bluegrass rust. This rust, *Puccinia Pourum*, is found over most of North America. Only in the far west, however, does it produce teliospores, and so only in this region can it have

²⁴Rusts of Australia 66-67. 1906.

²⁵Bot. Gaz. 52:169-192. 1911. Phytopathology 1:150-154. 1911.

²⁶Phytopathology 4:337-338. 1914.

²⁷Bur. Plant Industry Bull. 224:12-13. 1911.

²⁸Microscopical Journal Mch., 1890: 59-60.

²⁹Div. Veg. Phys. and Path. Bull. 16:21. 1899.

³⁰Trans. Wis. Acad. Sci. 15:98-107. 1905.

³¹Bur. Plant Ind. Bull. 216:56. 1911.

³²Trans. Wis. Acad. Sci. 15:106-107. 1905.

³³Bur. Plant Ind. Bull. 63:20. 1904.

æcia. It is so common throughout central and eastern regions early in the season, as well as later, that the idea of any seasonal migration from its region of possible æcia is clearly absurd. It must pass the winter in the uredinial stage, and it probably does so as mycelium in the leaves of the host. We have here a case of a rust which certainly maintains itself in a fair state of vigor for some years without the intervention of æcia and probably maintains itself indefinitely. Of course it is possible that it is constantly being renewed in vigor in the west by the presence of the æcia, and that the fungus thus renewed in vigor is slowly but continuously migrating eastward; but such a hypothesis strikes one as being fanciful rather than likely to be true.

Puccinia Sorghi, the corn rust, is another species with a wide distribution. It is usually not difficult to collect in any field of corn after tasseling time. The infection is usually not heavy, however. The æcia occur on *Oxalis*, but they occur so seldom that they seem to have little to do with the actual propagation of the rust. It is probably carried over from one year to the next by urediniospores which survive the winter or by the uredinial stage in living plants in southern regions. The latter source of infection seems more likely for this rust than for wheat rust because of its later appearance and less severity.

Puccinia Asperifolii, the leaf rust of rye, is a rust which has no known æcia in this country. Its case in America is therefore comparable with that of the bluegrass rust in this region or of *P. graminis* in Australia. It has to maintain itself by the sporophytic stage only.

Uromyces caryophyllinus on carnation is another rust which has no æcia in this country. There is no direct evidence that it can maintain itself over winter, for it usually appears in greenhouses; but it must have passed through thousands of uredinial generations since it was introduced, yet it seems to show no particular loss of vigor.

Puccinia Chrysanthemi is a Japanese species which has been introduced into America and Europe. It attacks cultivated chrysanthemums, chiefly in greenhouses. It has now been known in this country for about a decade and a half, and during this time it has never, so far as is known, produced a teliospore, although in northern Japan and in the mountains of Japan they are common. During this time no great impairment of vigor seems to have taken place, although chysanthemum growers are able to keep it in check by the use of resistant varieties and by the exercise of care in watering.

The rusts so far considered are some of the more ordinary species, belonging to the Aecidiaceæ. There are species in both the Uredinaceæ and the Coleosporiaceæ, however, which seem to have the same ability to maintain themselves indefinitely in the uredinial stage.

Among the Uredinaceæ two of the most common rusts are *Melampsora Medusæ* on *Populus* and *M. Bigelowii* on *Salix*, both of which have æcia on *Larix*. The æcia are so much alike that it is impossible with our present knowledge to tell them apart. It seems well here, therefore, to consider the two species together, although there are definite morphological characters in the urediniospores which mark them as clearly distinct from each other. The collections of *M. Bigelowii* in the Arthur herbarium show its presence in nine counties of the State, the first collection being made in 1887 and the last one in 1914, both in Tippecanoe County. The rust is common and the epidemic is usually severe. The only explanation which seems reasonable for not having collections from all counties in the State is in the lack of collectors being at work in those not represented. *M. Medusæ* is represented by collections from five counties in Indiana, and the same remarks as to prevalence and severity that were applied to *M. Bigelowii* apply to this species also. Both of these rusts have a range also far to the southward and westward of this region. Their æcia, to the present time, have not been collected nearer this region than New York and Wisconsin. However, it is likely that they do occur nearer because the larch has a range extending as far south as northern Illinois and northern Pennsylvania. It also occurs occasionally as an ornamental tree at various places in the State. The æcia probably occur within the hundred eighty mile distance from the northern half of the State and perhaps from all parts of the State. It does not seem reasonable to think, however, that æcia occur within several hundred miles of the southern range of the fungi. The natural assumption is, therefore, that they are able to pass the winter in the uredinial stage.

Bubakia Crotonis, on *Croton monanthogynus*, has been taken four times in Indiana, from at least three counties, and over a period of time extending from 1896 to 1912. It also extends as far north and west as Nebraska. No æcium is known for the rust, but the nature of the fungus suggests a Pinaceous host and a cœmoid æcium. *Cœoma strobilinum* on *Pinus palustris* and *Pinus taeda* has been suggested by Arthur.³⁴ Neither of

³⁴Bull. Torrey Club 33:519. 1906.

these species of pine, according to Sudworth,³⁷ has a range extending farther north than the southern border of Tennessee. This distribution is well outside the 180 mile limit established earlier in this paper. Two other species of pine, *P. echinata* and *P. virginiana*, have a distribution which might possibly meet the requirements, but there seems to be no evidence other than their distribution that they carry the æcia of this rust. Since *Cronton monanthogynus* is an annual, the evidence seems to favor the idea that the urediniospores are able to survive the winter.

Two species of *Pucciniastrum* occur in Indiana, *P. Agrimonie*, on *Agrimonia*, and *P. Hydrangeæ*, on *Hydrangea*. The former has been taken in five counties in the State at various times since 1896, and usually the infection is severe. *P. Hydrangeæ* has been taken three times in Tippecanoe county only. No æcia are known as yet for either of these species, but the æcia of the different species of *Pucciniastrum*, so far as known, are species of *Peridermium* on leaves of *Abies* and *Tsuga*. Judging by the distribution for these trees given by the manuals, Indiana is probably just outside of a 180-mile zone south of their distribuion. These trees are often planted for ornament, however, and the possibility exists that the æcia are to be found in the State. The rust occurs, however, as far south and west as the state of Mexico in the country of Mexico, and it is not to be expected that a species can travel so far in a season.

Among the Coleosporiaceæ, there are at least four species which have been collected in the State under conditions which lend color to the idea that they were carried over the winter in the uredinial generation. The rusts of the genus, *Coleosporium*, have their uredinia and telia on various broad leaved plants. Their æcia are leaf inhabiting species of *Peridermium* on pines. *Coleosporium Terebinthinaceæ* was collected in the autumn of 1912 and 1914 on *Silphium terebinthinaceum* in a restricted area near Lafayette. In the latter season, the species was limited to a patch a few rods in extent; other *Silphium* plants in the same patch were unaffected; and no affected *Silphium* plants could be found across a small ravine, although unaffected ones occurred in abundance. Other plants a mile or so away in two directions were examined but were found uninfected. The æcial stage of this rust is not known, and so it is impossible to say positively how near to this locality the æcia may approach. The nearest collection of *Peridermium* on pine leaves to be found in the Arthur herbarium is an undetermined collection on *Pinus virginiana* from Mammoth

³⁷Forest Atlas. Geographic distribution of North American Pines. Part 1, Maps 25 and 35. 1913.

Cave, Ky. Mammoth Cave, as well as can be told by scaling on the map, is approximately 215 miles from Lafayette. It is conceivable, of course, that a wind-borne spore from such a *Peridermium* could have started the infection of *Silphium* plants each year; but when we consider the likelihood that the two species do not belong together, and the fact that the rust was found in practically the same place both times, together with the fact that the host is a perennial plant, it seems more reasonable to think that the original infection was started by a stray spore, and that its further propagation and carrying over the winters was accomplished in the uredinia! stage, either by surviving spores, or by mycelium in the living host.

A somewhat similar case is that of *Coleosporium Ipomoeæ*, which has been collected repeatedly in Tippecanoe County since 1895 on *Ipomoea pandurata*. It occurs in great abundance and is doubtless to be found in practically all parts of the State where this host is found. The same thing is true for this species as for the preceding regarding the alternate stage and the possibility of the epidemics being started by æciospores, with this addition, that because of the more general distribution and greater commonness of the fungus, it is much less likely to be started each year by æciospores.

Coleosporium Vernoniæ, on different species of *Vernonia*, has for its æcial stage *Peridermium carneum* on *Pinus Elliottii* and *P. palustris*. It has a very wide distribution in the State, being represented in the Arthur herbarium from eight counties. The hosts of the æcia according to Sudworth³⁶ and Small³⁷ are both confined to an area south and east of central North Carolina and the north third of Alabama. This distance from Lafayette, as scaled on the map, is approximately 430 miles, a distance about 2.5 times as large as our maximum distance which we might expect a rust to migrate in a season. Moreover, it has been collected at Lafayette in different years as early as July 18 and July 24, which dates are early enough in the season to render it even more unlikely that the infections were developed, even indirectly, from æciospores of the same season.

Coleosporium Campanulæ is a species occurring in Indiana on *Campanula americana*. The æcium is known as *Peridermium Rostrupi* and occurs on *Pinus rigida* in eastern Ohio. The closest approach of the range of the host to Lafayette, according to Sudworth's map³⁸ is in eastern Ohio, which is

³⁶l. c. Map 35.

³⁷Flora of the Southeastern United States 33. 1913.

³⁸Forest Atlas. Geographic distribution of North American Pines. Part I. Map 26. 1913.

approximately 250 miles distant. That the fungus at least sometimes winters over is evidenced by the fact that it has been collected in the vicinity of Lafayette on rosettes of the host as early in the season as May 6. There is little or no doubt that it had wintered in the unredinal stage, probably as mycelium in the living leaves of the host.

Perhaps the clearest indication of the survival of the winter by uredinospores or mycelium outside of the Aecidiaceæ occurs in *Colcosporium Solidaginis*, on *Solidago*, *Aster*, and a few other Carduaceous hosts. This species is very widespread throughout the United States and is exceedingly common. Its exceeding commonness is attested by the fact that its Indiana distribution is represented in the Arthur herbarium by 44 mounted collections and a few unmounted ones, from 10 counties, and extending over a period of time from 1890 to the present. The æcial stage, *Peridermium acicolum*, occurs on *Pinus pungens* and *P. rigida*, with a distribution from Massachusetts and central New York to central North Carolina. According to Sudworth's maps³⁹ *Pinus rigida* is the one of these two æcial hosts which is nearer this section. Its nearest approach, as already shown, is eastern Ohio, which is approximately 250 miles distant from Lafayette. This is a greater distance than we would expect the fungus to migrate in one growing season; but the fungus extends also much farther to the west and northwest, so far, in fact, that it seems almost absurd to think it could have spread so far from its æcial base in a season. Furthermore, the writer on the first and second of July in 1912 made collections in eastern Indiana which show that the species was already well established for the season in a region a mile or more in extent. For such an infection, spores must be present in some quantity or must be present very early. But this is not the most convincing evidence at hand. There is a collection from Lafayette on *Solidago ulmifolia*, made June 25, 1896, and one on *S. serotina* made May 15, 1901. There is also one on *Aster cordifolius* made May 30, 1896, and one on *Aster* sp. indet. made May 12, 1902. This last collection is on the rosette leaves of the plant which were practically in contact with the ground, and the rust is well developed. The collection was actually made earlier in the season than any æcial collection of the rust at hand except one, which was made at Durham, N. C., May 3, 1910. The range for the æcial collections is May 3 to July 6; and it was clearly impossible for this specimen to have resulted from infection tracing back to æciospores of the same spring. The circumstance seems to be much more easily ex-

³⁹l. c. Maps 26, 30.

plainable by assuming that some urediniospores or mycelium survived the winter.

It seems fair, then, to judge from the foregoing that a good many rusts can pass the winter and propagate themselves for a long time, and probably indefinitely, without the intervention of sexual reproduction. This is in line with the experience of Freeman and Johnson,⁴⁰ who carried *Puccinia graminis*, *P. rubigo-vera*, and *P. simplex* through 52 uredinial generations without apparent degeneration, and of Fromme,⁴¹ who similarly carried *P. coronifera* on oats through thirty-seven uredinial generations, and of carnation raisers generally, who still find the carnation rust an enemy to be fought although it has in all probability never produced an æcium on this continent.

The evidence, therefore, which is to be gained from the behavior of the rusts concerning the question as to whether or not a plant species can long maintain a high degree of vigor without sexual reproduction is quite definitely in favor of the idea that it can. True it is that in the long cycled rusts an effect of stimulation follows the stage in which the sexual fusions take place, but this effect becomes dispelled by one or two uredinial generations, so that the rust is then back at the old level of vigor; and it remains there through an indefinite number of uredinial generations.

Purdue University,

Lafayette, Indiana.

⁴⁰Bur. Plant Ind. Bull. 216:34. 1911.

⁴¹Bull. Torrey Club 40:510-511. 1913.

CORRELATION OF CERTAIN LONG-CYCLED AND SHORT-CYCLED RUSTS.

H. C. TRAVELBEE.

When in 1897 Dietel, in his work "The Uredinales" for "Die natürlichen Pflanzenfamilien" of Engler and Prantl, pointed out the remarkable similarity between the teliospores of *Puccinia Mesneriana* Thüm., on *Rhamnus* and those of *Puccinia coronata* Cda. and *Puccinia coronifera* Kleb. on grasses, which have their æcia on *Rhamnus*, he established the first observation on correlations between rusts of widely different species. He also called attention to the fact that a similar condition obtains between the teliospores of *Puccinia ornata* Arth. & Holw. on *Rumex* and the teliospores of the grass rust, *Puccinia Phragmitis* (Schum.) Körn. which has *Rumex* for its æcial host. In both of these cases we note the teliospores of a short-cycled rust appearing on the æcial host of a long-cycled heteroecious rust. The teliospores of the two species are morphologically alike although appearing on host plants of quite different families.

About this same time (1898) Fischer stated* that quite independently of Dietel, he found by his researches a list of similar relationships. He reported five heteroecious species of *Puccinia*, two of *Chrysomyxa*, one of *Melampsora* and one of *Coleosporium*, all having short-cycled forms appearing on their æcial hosts, agreeing with their teliospores. He also listed three *Uromyces* and one *Puccinia* which show this sort of a relationship with certain micro- or hemi-forms.

It is worthy of note here that the complete life history of all the forms correlated in this manner were known at the time the observations were made.

When in 1903 and 1904 Tranzschel connected into a heteroecious life history two rust forms which until that time had never been suspected of bearing any relationship to each other, he made a wonderful advance along the line of this sort of investigation. His method was as unique as it was important, and on account of the interesting field it opens for investigators is worthy of detailed mention.

He had an unconnected *Aecidium*, *A. punctatum* Pers., on *Anemone ranunculoides*, and was endeavoring to find its alternate host. He observed that on *Anemone nemorosa* there appeared a short-cycled *Puccinia*, *P.*

*Beiträge zur Kryptogamenflora der Schweiz. 1:109. 1898.

fusca (Pers.) Wint., whose morphological aspects were strikingly similar to those of the *Aecidium*, i. e., the sori of the two rusts were arranged in the same manner; the effects on the host plant were the same, and the macroscopic characters of the two were alike. He concluded that the two forms were closely related phylogenetically. Then his problem was: How find the alternate stage of the *Aecidium* form?

On examining the teliospores of the short-cycled form microscopically he found them to possess very striking features, having a roughly warty wall, and being strongly constricted at the septum. A careful examination of his unconnected *Puccinias* revealed one having spores similar to those of the short-cycled form on *Anemone*, *Puccinia Pruni-spinosæ* Pers., the plum rust. He cultured the æciospores from *Anemone* on the leaves of plum and peach trees and grew the uredinia and telia of the plum rust. In his investigations at this time he combined, in this way, five heterœcious species. Three of these species he proved by cultures; namely,

1. *Puccinia Pruni-spinosæ* Pers. with *Aecidium punctatum* Pers.
2. *Uromyces Veratri* (DC.) Schroet. with *Aecidium Adenostylis* Sydow.
3. *Uromyces Rumicis* (Schum.) Wint. with *Aecidium Ficaræ* Pers.


The correlation, then, of two rusts of distinct species depends upon several things: first the family relationship of the host plants. The rusts are usually on the same or closely allied species of host plants. The macroscopic characters of the two rusts are also of decided importance. They can usually be expected to have similar effects on the common host plant; and the location and disposition (i. e., whether they are amphigenous, epiphyllous, hypophyllous, caulicolous; numerous or few, scattered or crowded; circular, oblong or irregular, etc.) of the sori are important factors.

The most important thing, however, is the agreement of the microscopic characters of the analagous spore forms. The teliospores of the short-cycled and long-cycled forms are compared for thickness, color and markings of the walls; the condition of the apex (thickened or not); measurements of the spores; length, color and type of pedicels; and the general conformation of the spores.

Owing to the fact that the knowledge of the rust fungi of very many regions is not complete, the geographic distribution can not be considered as extremely important. Nevertheless many interesting comparisons are shown by a study of the distribution of the correlated forms.

For some time past the writer has had access to the Arthur herbarium, and at the suggestion of Dr. J. C. Arthur, made a list of the heterœcious species of *Puccinia* having their teliospores on grasses, whose complete life histories are known. The teliospores of the short-cycled forms appearing on the æcial hosts of these rusts were compared with the teliospores of the long-cycled forms with the intention of compiling such a list as Fischer published. The following combinations were found to appear as good correlations:

LIST OF CORRELATIONS.

	RUST.	TELIAL HOST.	ÆCIAL HOST.
1.	{ <i>Puccinia Crandallii</i> Pammel & Hume <i>Puccinia Symphoricarpi</i> Hark.	{ <i>Festuca</i> <i>Poa</i> <i>Symphoricarpos</i>	<i>Symphoricarpos</i>
2.	{ <i>Puccinia simillima</i> Arth. <i>Puccinia Anemones-virginiana</i> Schw. ...	{ <i>Phragmitis</i> <i>Anemone</i>	<i>Anemone</i>
3.	{ <i>Puccinia Stipæ</i> Arth. <i>Puccinia Grindeliæ</i> Peck.	{ <i>Koeleria</i> <i>Oryzopsis</i> <i>Stipa</i> <i>Bigelovia</i> <i>Chrysopsis</i> <i>Chrysothamnus</i> <i>Grindelia</i> <i>Gutierrezia</i>	{ <i>Aster</i> <i>Bigelovia</i> <i>Chrysopsis</i> <i>Chrysothamnus</i> <i>Erigeron</i> <i>Grindelia</i> <i>Gutierrezia</i> <i>Lygodesmia</i> <i>Nothocalais</i> <i>Senecio</i> <i>Solidago</i>
4.	{ <i>Puccinia Andropogonis</i> Schw. <i>Puccinia Seymeriæ</i> Burrill.	{  <i>Andropogon</i> <i>Dasystema</i>	{ <i>Castilleja</i> <i>Dasystema</i> <i>Penstemon</i>
5.	{ <i>Puccinia pustulata</i> (Curt.) Arth. <i>Puccinia Comandræ</i> Peck.	{ <i>Andropogon</i> <i>Comandra</i>	<i>Comandra</i>
6.	{ <i>Puccinia monoica</i> (Peck) Arth. <i>Puccinia Holboellii</i> (Horn.) Rostr. ...	{ <i>Kæleria</i> <i>Trisetum</i> <i>Arabis</i>	{ <i>Arabis</i> <i>Parrya</i> <i>Schænocrambe</i> <i>Smelowskya</i>
7.	{ <i>Puccinia Agropyri</i> Ell. & Ev. <i>Puccinia DeBaryana</i> Thuem.	{ <i>Agropyron</i> <i>Anemone</i>	<i>Anemone</i>
8.	{ <i>Puccinia rubella</i> Arth. <i>Puccinia ornata</i> A. & H.	{ <i>Phragmitis</i> <i>Rumex</i>	{ <i>Rheum</i> <i>Rumex</i>
9.	{ <i>Puccinia rhamni</i> Wettst. <i>Puccinia Mesneriana</i> Thuem.	{ <i>Avena</i> <i>Rhamnus</i>	<i>Rhamnus</i>

The last two combinations in this list are the ones noted by Dietel.

Orton¹ dealt with quite a different type of correlation when he reported in detail the similarities between six species of heteroecious *Uromyces* and six species of heteroecious *Puccinia*. He also extended this study to include autoecious species of *Uromyces* and *Puccinia*. In every instance the host plants of the two rusts are of the same species, or of species closely related morphologically and phylogenetically. Because of the cellular difference in the teliospores (*Uromyces*, one-celled; *Puccinia*, two-celled) Orton laid special emphasis on the agreement of the microscopic characters of the aeciospores and urediniospores of the two rusts, remarking only in a general way similarities between the teliospores.

When we consider the differences in the number of spore forms in the life cycles of the various species of rusts, and take into consideration the morphological variation of the analogous spores, it is apparent that the possibilities of correlation are numerous. There are many problems presented in connection with such correlations. The choice of host plants, the similarity of analogous spore forms, and the like effects on the host all point to a common ancestor. What then is the primitive form? What is the evolutionary history of the derivative species? How great a range may be expected in the variations of correlated species? These and similar questions arise when a theoretical consideration of the condition is undertaken.

The practical application of knowledge gained by correlation studies will be along the lines of culture work, especially in forecasting the alternate host plants of unconnected aecial or telial forms.

Mycologia, IV: No. 4, July, 1912.

Purdue University,
Lafayette, Ind.

SOME SPECIES OF NUMMULARIA COMMON IN INDIANA.

CLAUDE E. O'NEAL.

The difficulty of distinguishing the various species of *Nummularia*, and even the genus itself from the genus *Hypoxylon*, is quite evident to anyone who has made any attempt at their classification. In a paper entitled "A Monograph of the Common Indiana Species of *Hypoxylon*,"* C. E. Owens, by the aid of plates and an artificial key, sets forth the characteristics of the common species of the latter genus. The purpose of this paper is to do a similar work with the available species of the genus *Nummularia*.

In the study of *Nummularia*, attention is first directed toward the stromata, which appear as blackish or brownish incrustations on the dead trunks and limbs of our common deciduous trees. In form, the stromata vary greatly, but in general, they belong to two types, one of which may be described as cup-shaped, and the other as convex. The former type is usually orbicular or elliptical in shape, while the latter may be either orbicular, elliptical, or broadly effused with an irregular outline.

The stromata arise beneath the epidermis of the substratum where they may remain concealed for some time, but sooner or later, the epidermis is broken through and the spore-bearing surface is exposed. Sometimes in old specimens the entire epidermis may be removed and the erumpent characteristic overlooked. In some cases the entire cortex may decay and fall away, leaving the stromata standing out on the decorticated surface. Again, some of the more resistant ones may be found in good condition on a log that is ready to drop to pieces from decay.

In width the various Indiana species range from a few millimeters in the cup-shaped forms to several centimeters in the broadly effused types. The thinnest ones that the author has found have been about one-half millimeter thick while some of the cup-shaped forms may have a thickness of

*Proceed. Ind. Acad. Sci. 1911, p. 291.

half a centimeter or more. In the early stages of development in the species described the stromata may be lightly colored, but when the sporidia are mature the color of the fertile layer is black. Sometimes the substratum is stained by the fungus. One species is readily recognized by the characteristic orange color imparted to the wood beneath the stromata. Another species gives a peculiar ring-like marking to both wood and bark.

If a stroma be cut through, the flask-like perithecia are found deeply embedded in it. They are arranged in a single row and open by minute pores (ostiola) on the upper side of the stroma. The ostiola in some species may be rather prominently raised giving the fertile surface a pimpled appearance, or they may be sunken; while in some forms they may be so obscure as to be passed over unnoticed. The edges of the stromata are usually sterile.

The perithecia contain many eight-spored asci. These asci are cylindrical in shape and bear the spores in a single row. They are readily distinguished from the paraphyses with which they are found by their shorter length and the spores which they contain.

The spores of the *Nummularias* vary greatly in size, color, and shape. The largest ones that the author has observed were about 16 microns long and one-half as broad, while the smallest were about 5 microns long and about $2\frac{1}{2}$ microns wide.

The shape varies from elliptical to orbicular. When the spores are young they are usually hyaline, but as they become older, they turn brown and in some species they finally become opaque.

In general, the Indiana species of *Nummularia* are saprophytic in habit and consequently have but little economic importance. Under certain conditions, however, *N. discreta** becomes parasitic and causes considerable damage to poorly kept apple trees. To many fruit growers it is known merely as apple canker; others distinguish it as the blister canker. The fungus gains a footing in a wound or in a decayed portion of the tree and spreads to the living parts. The stromata arise upon the mycelia beneath the epidermis of the host. The overlying epidermis shortly becomes dry and papery and sooner or later it is torn and drops away leaving the

*Canker of Apple—Hasselbring, Ill. Exp. Sta. Bull. 70.

Fungous Diseases of Plants—Duggar, pp. 282-284.

Apple Blister Canker and Methods of Treatment—W. O. Gloyer, Ohio Exp. Sta. Cir. 125.
The New York Apple Tree Canker. Bulls. N. Y. Ag. Ex. Sta. Nos. 163 and 185. Paddock.
The Control of Canker in the Orchard—J. R. Cooper, Neb. Hort., Vol. 3, 1913.

cankerosus stromata exposed. Since a great deal of damage is done before the stromata are formed, it is necessary to identify the disease from other characteristics. The remedy for this parasite is obviously the proper treatment of wounds and the removal and destruction of affected parts. Care in pruning would in many cases prove a sufficient preventive.

This fungus has been reported on several other hosts including *Ame-lanchier* but in these cases the economic importance is insignificant.

KEY TO THE COMMON INDIANA SPECIES.

- I. Stroma cup-shaped with perithecia opening on the concave side. (A).
- II. Stroma convex or plane. (B).
 - A. Stroma prominent, orbicular, with a thick, raised margin; stroma 4 to 7 mm. across; spores globose or subglobose, 10-16 microns1. *N. discreta*.
 - A. Stroma erumpent-superficial, either orbicular or elliptical; margin not so thick nor so regularly bulging as in the preceding species; concave part $\frac{1}{2}$ to 1 cm. across. Spores subinequilateral, 10-15x5-6 microns2. *N. repanda*.
 - B. Stroma dull black, orbicular, elliptical, or broadly effused; ostiola rather prominently raised, 3-5 per linear mm. Spores 12-16x6-8 microns3. *N. bulliardi*.
 - B. Stroma shiny black, more or less furrowed, staining the wood of the substratum orange color. Spores 12-16x5-7 microns. .4. *N. tinctor*.
 - B. Stroma thin, orbicular, suborbicular or linear; ostiola depressed; ranging from $\frac{1}{2}$ to 1 cm. across. Spores $4\frac{1}{2}$ -5x2-2 $\frac{1}{2}$ microns.....5. *N. microplaca*.

DESCRIPTIONS.

1. *Nummularia discreta*, (Schw.) Tul. Plates I and II.
Sphaeria discinola, Schw. Syn. Carp. No. 63.
Sphaeria discreta, Schw. Syn. N. Am. 1249.
Sphaeria excavata, Schw. l. c. 1250 (Sec. spec. in herb. Schw.)
Nummularia discreta, Tul. Sel. Carp. II, p. 45.

Stroma erumpent, circular or subcircular, sometimes uniting to form elongated patches, cup-shaped when mature, with a thick raised margin; ashen or grayish yellow, becoming black; the concave surface at first white

punctate from the minute ostiola which are hardly visible when mature. The bark and wood beneath the stromata are marked with a black circumscribing line. Perithecia arranged in a single row, oval or ovate-cylindrical, about one millimeter long, usually rather abruptly contracted above into a short neck and extending to the base of the stroma. Asci cylindrical, 140-170x10-15 microns (E. & E., 110-120x10-12); spore bearing part, 110-125x10-15 microns. Paraphyses, long and filiform. Sporidia subglobose, almost hyaline at first, finally becoming opaque, 10-16 microns in diameter. (11-13 Gloyer, 10-12 E. & E.)

On dead trunks and branches of *Pyrus malus*; quite common on the living trees as well. Practically every orchard visited in Hendricks, Putnam and Monroe counties, Indiana, as well as Delaware County, Ohio, showed traces of this fungus. Gloyer reports it especially abundant in southern Ohio. Reported on *Amelanchier canadensis*, Newfield, N. J., and on *Gleditschia triacanthos*, Ohio, (Morgan); also (See Saccardo in Syll.) on *Sorbus*, *Ulmus*, *Cercis* and *Magnolia*.

On apple trees this fungus usually attacks the trunks and larger limbs, making somewhat sunken, cankerous areas several inches in length. The dead bark is separated from the sound by a distinct line and cracks occur along this boundary. At the beginning, living spots within the cankerous area give the affected parts a mottled appearance. This distinguishes it at this stage from other cankers.

2. *Nummularia repanda*, (Fr.) Nke.

Sphaeria repanda, Fr. S. M. II, p. 346, Obs. Mycol. I, p. 168.

Hypoxyton repandum, Fr. Summa Veg. Sc. p. 383.

Nummularia pezizoides, E. & E. Bull. Torr. Club, XI, p. 74.

Nummularia repanda, Nitsch. Pyr. Germ. p. 57.

Exsic. Fekl. F. Rh. 2178. Thun. M. U. 1460.

Stroma erumpent-superficial, orbicular or subelliptical, $\frac{1}{2}$ to 1 cm. in diameter, concave and often with a thin, erect, rather broad margin, reddish-gray at first, finally black; disk mammillose from the projecting ostiola. Perithecia monostichous, immersed, ovate-oblong, $\frac{1}{2}$ to $\frac{3}{4}$ mm. long, crowded, causing the sides to be somewhat compressed. Asci cylindrical, sessile, eight-spored, 110-120x8 microns, with long filiform paraphyses. Sporidia obliquely uniseriate, narrow ovate, obtuse, subinequilateral, dark brown, $8\frac{1}{2}$ -14x4-7 $\frac{1}{2}$ microns. (E. & E. 11-14x4-5 microns; Sacc. in Syll., 15-16x6-7 microns.) Readily distinguished from *N. discreta* by its differently shaped spores and its mammillose disk.

On *Hicoria*, Clark County, Indiana, (Van Hook); on wood and bark, Topeka, Kans. (Craigin), and on bark, Ottawa, Canada (Macoun); on bark of *Ulmus americana* Missouri, (Demetrio). On *Sorbus aucuparia* in Europe.

3. *Nummularia bulliardi*, Tul. Plate III, Figs. 1, 2 and 3.

Hypoxyton nummularium, Bull. Champ. tab. 468, fig. 4.

Sphaeria nummularia, D. C. Flore Fr. II, p. 290.

Sphaeria anthracina, Schm. & Kze. Mycol. Hefte 1, 55.

Sphaeria clypeus, Schw. Syn. N. Am. 1219.

Nummularia clypeus, Cke. IX, 507.

Exsic. Ell. N. A. F. 85. Rab. F. E. 2956. Rehm, Asc. 977.

Illustrations. (See Sacc. XX, p. 202, for list of.)

Stroma at first covered by the epidermis, soon erumpent, almost superficial and free, convex, orbicular or oval, sometimes irregular in shape or broadly effused, black inside and out, punctulate from the slightly prominent ostiola, clothed at first with a reddish or rusty layer of conidia. Perithecia rather large, ovate, black, loosely included in the packed cells of the stroma. Asci cylindrical with very short stalks, spore-bearing part 115-140x7-10 microns (E. & E. 100-115x10 microns), with long, stout paraphyses. Spores eight, uniseriate, elliptical, hyaline becoming opaque, 10-23x5-10 microns, mostly about 15-20x6-8 (E. & E. 12-15x7-9. Sacc. 12-14x9-10 microns.

In the field this species is liable to be confused with certain species of *Diatrype* (Fig. 2, Plate III) but may be readily distinguished from them by the color of its spores. Collected in abundance in Brown, Clark, Hendricks, Monroe and Putnam counties, where it usually attacks the beech and more rarely the maple. Reported common on the dead trunks and limbs of various deciduous trees in Europe and North America. According to Ellis and Everhart, it occurs for the most part on oak in the vicinity of Newfield, N. J.

4. *Nummularia microplaca*, (B. & C.) Cke. Plate IV, Fig. 4.

Diatrype microplaca, B. & C. Journ. Linn. Soc., X, p. 586.

Anthostoma microplacum, Sacc. Syll. I, p. 298.

Nummularia microplaca, Cke. Syn. 837.

Exsic., Rav. Fungi Car. IV, 39. Rav. Fung. Am. 355. E. & E.

N. A. F. Second Ser. 1556.

Illus., Revue Myc. VII, (1885) tab. 52, fig. 3.

Stroma orbicular to subelliptical, $\frac{1}{2}$ to 1 cm. across, or elongated 1-4x $\frac{1}{2}$ -1 cm. or by confluence extending for long distances in grooves of the bark. It forms a thin carbonaceous crust, black, arising beneath the epidermis but soon becoming bare, surface even, faintly punctulate from the minute ostiola, which are not prominent but slightly depressed, the opening at first filled with a white farinaceous matter. Perithecia ovate-globose, small (less than one-half mm. across), arranged in a single row. Spore-bearing part of the ascus 40-50x4 microns (E. & E. 25x3 microns), or with the base about 60-80 microns long (Sacc. 37-50x4.5. E. & E. 45-50 long). Spores uniseriate, ends mostly slightly overlapping, elliptical, inequilateral, pale brown, 5-7 $\frac{1}{2}$ x2 $\frac{1}{2}$ -3 microns (E. & E. 4 $\frac{1}{2}$ -5x2-2 $\frac{1}{2}$. Sacc. 5-6x3 $\frac{1}{2}$ -4).

Should not be confused with *Hypoxylon Sassafras* which has very prominent perithecia while *N. microplaca* appears smooth and has stroma depressed.

Abundant near Bloomington, Indiana, on *Sassafras officinale*; reported on the same host in South Carolina (Ravenel) and in Ohio (Morgan and Kellerman); on *Persea*, Georgia (Ravenel).

5. *Nummularia tinctor*, (Berk.) E. & E. Plate IV, Figs. 1-3.

Sphaeria tinctor, Berk. Lond. Jour. Bot. IV, p. 311.

Hypoxylon tinctor, Cke. Syn. 996.

Diatrype?? tinctor, (Berk.) Sacc. Syll. I, 200.

Exsic., E. & E. N. A. Fungi, Second Ser. 1789.

Stroma very hard and brittle, much effused, showing the irregularities of the surface on which it grows, 1mm. thick, black, with surface almost smooth, but distinctly papillose from the projecting ostiola as seen under the hand-lens, wood beneath the stroma stained a beautiful reddish-orange color, and rendered very hard. Perithecia monostichous, crowded, elongated ($\frac{3}{4}$ mm. in length), covered above with the stromatic layer. Asci 100-140x6-10 microns (E. & E. 112x7-8). Spore-bearing part of ascus 75-120 long (E. & E. 90-100). Filiform paraphyses in abundance. Spores uniseriate, pale brown, conspicuously uniguttulate, oblong navicular, 13-20x5-8 microns (E. & E. 15x6).

On *Platanus*, *Fagus*, *Acer*, *Ulmus* and *Cercis* in the vicinity of Bloomington, Indiana (Van Hook). Occurs throughout the Mississippi valley and in the south as far east as Florida.

In this paper, free use has been made of North American Pyrenomyces by Ellis and Everhardt. The descriptions have been re-written to suit the material at hand. Especially has it been found necessary to revise the measurements of parts as seen under the microscope. The author wishes also to make due acknowledgment to Prof. J. M. Van Hook, of Indiana University for material and valuable assistance in the preparation of this paper.

*Indiana University,
Bloomington, Indiana.*

PLATE I.

Figures 1 and 2. Where *Nummularia discreta* thrives. Much of this is due to *N. discreta*. (From photos by the author in Hendricks County, Indiana.)



PLATE II.

Figure 1. *Nummularia discreta* on decorticated apple limbs. (Reduced.)

Figure 2. Same natural size.

Figure 3. Same before the bark has fallen away. (Natural size.)


Figure 4. Under side of a piece of bark showing white stains of the fungus. (Natural size.)




1



3



2



4


PLATE III.

Figure 1. *Nummularia Bulliardi* on beech, showing a common form of stroma. (Natural size.)

Figure 2. A stroma of a fungus (*Diatrype*) very similar to *N. Bulliardi*. (Natural size.)

Figures 3 and 4. Orbicular stromata of *Nummularia Bulliardi* on beech. (Natural size.)


Figure 5. Stroma taken from a log which was falling to pieces. (Natural size.)




1




2



3



4



5

PLATE IV.

Figure 1. *Nummularia tinctor.* (Natural size.)


Figure 2. Same showing section of the wood beneath stroma. The dark colored parts are a beautiful orange.

Figure 3. Same showing furrowed stroma. (Natural size.)

Figure 4. *Nummularia microplaca* on *Sassafras*. Closely resembles *Hypoxylon Sassafras*. (Natural size.)



1



2



3



4

THE GENUS ROSELLINIA IN INDIANA.

GLEN B. RAMSEY.

In this paper it is the author's purpose to present the genus *Rosellinia* as found in Indiana. Limited means for collecting and the unfavorable weather conditions during the past year for the growth of this class of fungi render an exhaustive treatise on the genus *Rosellinia* impossible. Only those species that have been found in Indiana, together with their descriptions and an account of their habits and habitat have been included. A description of some of the more common parasitic species has also been appended.

There are now over one hundred seventy species described, the most of them being saprophytic. As in other genera of the Pyrenomycetes, *Rosellinia* has a vegetative phase which is found in the substratum or host. The white thread-like mycelium may readily be found in decaying logs and stumps. There are, however, some species that seem to flourish in wood that is quite firm. In most cases the actively parasitic stage is found on the roots and consists of a great abundance of white mycelium which does the greatest harm to the plant. This mycelium growing into the root system stops up the xylem cells, prevents the roots and rootlets from performing their functions, thus finally starving the plant to death.

The fruiting parts of *Rosellinia* do not develop until late in the season, the conidial stage being found in late summer or early autumn, with the perfect or ascigerous stage following and maturing in late fall or early winter. The perithecia with their abundant asci and filiform paraphyses are found in good condition for collecting from October to February. The spring rains and warm weather, together with the frost action during the winter, cause the perithecia to disintegrate rapidly when spring comes. Most of the specimens at hand were collected during the early winter months.

In *Rosellinia* the perithecia are more or less crowded or gregarious and superficial, but often having the base sunken in the matrix. Perithecia ovate to sub-globose, papilliate, sub-carbonaceous, black, bare

or bristly, with a distinct ostiolum. The asci are cylindrical, eight spored. Spores continuous, broadly ovate, elliptical, oblong or fusiform, brown or black, with or without hyaline appendages. Paraphyses filiform. Stroma tomentose, often wearing off with age and exposing the perithecia as round, brownish black heads with their papilla-like ostiola in the center.

In studying the species of *Rosellinia* we are confused at times when we find forms which closely resemble certain forms of the genus *Hypoxylon*. In such cases a clear distinction seems impossible, yet these two genera are clear cut in their separation by botanists, being separated on the ground of the presence of a stroma in *Hypoxylon*, and the absence of a distinct stroma in *Rosellinia*. To anyone that has made a study of either of these two genera, the superficiality of this basis of separation is quite evident. Students of the genus *Hypoxylon* know that the perithecia of certain forms become scattered, and especially with age, the stroma is wanting.

The genus *Rosellinia* is placed under the *Sphaeriaceae* by both Lindan and Ellis and Everhart. This separates it widely from *Hypoxylon*. Saccardo puts it under the brown spored one-celled forms of *Xylariaceae* along with *Hypoxylon*. The author likes this position on account of the great similarity of spores, asci and perithecia, as well as the above mentioned similarity of forms where the absence of stromata is noticeable.

Variation within a given species often makes it almost impossible to formulate a key that will hold in all cases. In order to eliminate this difficulty the species are made to run in two ways. The species *R. subiculata* for example has in the earlier stage a decided waxy sulphur-yellow subiculum and the perithecia are scattered, but as it grows older the subiculum finally disappears so that one might readily confuse it with other species that never have a subiculum. The ascis and spore measurements are probably the most constant and reliable in forming a basis for a key. The second key, it is hoped, will prove helpful in determining any doubtful species that do not run satisfactorily in the key of external characteristics.

The accompanying figures from photographs will assist in determining the species. In order to get the greatest contrast possible, time exposures were made in a subdued light and a special contrast developer was used.

The description of species have been adapted for the most part from

the original descriptions as given by Ellis and Everhart in "North American Pyrenomycetes", and Saccardo's "Sylloge Fungorum." Practically all of the descriptions have been rewritten and additional data added from specimens at hand. All measurements are original. Where asci and spore measurements by Ellis and Everhart differ, their figures are also given.

KEY TO SPECIES.

(Based on external characteristics.)

- I. Perithecia large ($\frac{3}{4}$ - $1\frac{1}{2}$ mm.), seated on a subiculum.
- A. Subiculum usually prominent.
1. Brown or purplish brown, persistent.....1. *R. aquila*
2. Sulphur-yellow, evanescent6. *R. subiculata*
- B. Subiculum scanty.
1. Dark brown, perithecia crowded.....2. *R. medullaris*
2. Black, perithecia confluent3. *R. mammiformis*
- C. Subiculum wanting, perithecia more or less scattered.
1. Base glandular-roughened4. *R. glandiformis*
2. Not glandular-roughened5. *R. mutans*
- II. Perithecia small ($\frac{1}{3}$ - $\frac{1}{2}$ mm.).
- A. Perithecia gregarious, often crustaceous, not bristly.7 *R. pulveraceæ*
- B. Perithecia usually scattered, bristly, dark brown....8. *R. ligniaria*

(Based largely on microscopic characteristics.)

- A. Perithecia large ($\frac{3}{4}$ - $1\frac{1}{2}$ mm).
- B. Spores more than 18 microns long.
- C. Asci more than 150 microns long.....1. *R. aquila*
- CC. Asci less than 150 microns.
- D. Asci 7-8 microns wide.....2 *R. medullaris*
- DD. Asci 8-10 microns wide.....3. *R. mammiformis*
- BB. Spores less than 18 microns long.
- C. Asci more than 95 microns long.....4. *R. glandiformis*
- CC. Asci less than 95 microns.
- D. Perithecia $\frac{1}{2}$ - $\frac{5}{8}$ mm.5. *R. mutans*
- DD. Perithecia $\frac{3}{4}$ -1 mm6. *R. subiculata*
- AA. Perithecia small ($\frac{1}{3}$ - $\frac{1}{2}$ mm).
- B. Asci more than 70 microns long.....7. *R. pulveraceæ*
- BB. Asci less than 70 microns long.....8. *R. ligniaria*

1. *R. aquila* (Fr.) De Not.

Sphaeria aquila Fr.

Sphaeria byssiseda Meckl.

Rosellinia aquila De Not.

Perithecia large, globose, 1-1.25 mm. in diameter, gregarious, crowded or sometimes confluent, with a distinct black, conic-papilliform ostium; dark brown at first with a thin tomentose coating, finally becoming bare. Subiculum rather thick and prominent, dark to purplish brown, nearly enveloping the perithecia at first but finally disappearing to a greater or less extent. Outer walls of the perithecia thick, brittle and carbonaceous. Inner wall coriaceous. Asci long, cylindrical (p sp.) 10-12.5 x 165-190 microns. Spores uniserrate, oblong, brown, 10-11 x 22.5-27.5 microns. (E & E) gives asci (p sp.) 8-10 x 100-130 microns. Sporidia 6-9 x 16-27 microns, with or without a short, obtuse, hyaline apiculus, 2-2.5 microns long at each end.

Common on decaying and fallen limbs, near Bloomington, Ind. Specimens at hand are on *Fagus*, *Acer*, *Quercus* and *Juglans*.

2. *R. medullaris* (Walls) Ces. & De Not.

Sphaeria medullaris Walls.

Rosellinia medullaris Ces. & De Not.

Rosellinia macouniana E. & E.

Perithecia more or less erumpent, large 1-1.5 mm. in diameter, ovate to sub-globose, covered at first with a pruinose-pubescent coat of a dull red or brick color, becoming black with age; loosely adnate, apex convex to conic-papilliform, surface dirty-roughened with a finely powdered sooty covering; very fragile. Wall double and intermediate in thickness between *R. aquila* and *R. thelena*. Subiculum slight.

Spores 7.5-12 x 20-25 microns. E. & E. give asci (p sp.) 7-8 x 100-120 microns. Sporidia 6-7 x 19-20 microns; ovoid, somewhat acute, brown, broader but not pointed as in *R. mammiformis*.

On *Cercis canadensis* and *Juglans*, Monroe County, Ind.

On examining a great number of perithecia the most of them were found to contain a white, granular mass such as described in *R. medullaris* by Saccardo, but close observation showed this material to be a fine powder of wood that had been brought into the perithecia from the bottom by a small larva that probably feeds upon the contents of the perithecia.

3. *R. mammiformis* (Pers.) Ces. & De Not.

Sphaeria mammiformis Pers.

Hypoxylon mammæformæ Berk.

Hypoxylon globulare (Bull.) Feikl.

Rosellinia mammiformis Sacc.

Perithecia gregarious, crowded or confluent, globose, large, 1-1.5 mm. in diameter, fragile, black and bare but not shining. Ostiolum abrupt, papilliform.

Asci (p sp.) 8-10 x 100-115 microns (E. & E).

Spores 10-12 x 20-25 microns, oblong, elliptical, sometimes slightly curved. E. & E. give sporidia 7-9 x 19-25 microns, without any distinct apiculus. It can be easily distinguished from *R. aquila* by the blacker, thinner walled and more fragile perithecia and the lack of a decided subiculum.

On bark of *Acer*, near Bloomington, Ind.

4. *R. glandiformis* E. & E.

Perithecia scattered, the base sunk in the wood about one-fourth. ovate-globose, roughened with glands, with a reinforcement around the lower half similar to the cup of an acorn. This thickening is, however, sometimes reduced to a mere granular coat. Ostiolum papilliform, small, sometimes obsolete, the apex being evenly rounded.

Asci not present in specimens at hand. E. & E. give (p sp.) 8-10 x 100-114 microns; paraphyses abundant. Spores 7-10 x 13.75-17.5 microns. E. & E. give 7-8 x 14-15 microns. Common on *Loriodendron* *Juglans* and *Fraxinus*, Monroe County, Indiana.

5. *R. mutans* (C. & P.) Sacc.

Sphaeria mutans C. & P.

Rosellinia mutans Sacc.

Perithecia more or less crowded or gregarious, rather small, about .5-.75mm. in diameter, at first clothed with a thin, tawny, evanescent tomentum, finally becoming smooth, black and shining; mostly globose with a papillate ostiolum. In the specimen at hand the region about the ostiola showed a distinct tendency to depress.

Asci subcylindrical (p sp.) 6.5-7.5 x 80-92.5 microns.

Spores uniserrate, elliptical, brown, 4.25-5.5 x 9-12.5.

Common on decaying *Juglans*, near Bloomington, Indiana.

6. *R. subiculata* (Schw.) Sacc.

Sphaeria subiculata Schw.

Hypoxyton subiculosum Berk.

Rosellinia subiculata Sacc.

Perithecia thin, usually gregarious or crowded but often more or less scattered in the early stages; globose, brownish-black and shining, mostly superficial, about .75-1 mm. in diameter. Ostiolum small and papilliform. Perithecia seated on a sulphur-yellow, waxy-pruinose subiculum which disappears with age leaving the black, shining perithecia closely resembling *R. mutans*.

Asci cylindrical (p sp.) 6.25 x 90 microns.

Spores inequilateral, elliptical, brown, 5-6.25 x 10-12.5.

Ellis and Everhart give asci 6-7 x 80-90. Spores 5-5.5 x 10-12 microns.

Common on *Quercus*, *Loriodendron*, and other rotten deciduous wood near Bloomington, Ind., and Jolietville, Ind.

7. *R. pulveracea* (Ehr.) Fekl.

Sphaeria pulveracea Ehr.

Sordaria Friesii Niessl.

Rosellinia pulveracea Fekl.

Rosellinia Friesii Niessl.

Sphaeria millegriaria Schw.

Sphaeria transversalis Schw.

Perithecia very small and minutely roughened, about $\frac{1}{4}$ mm. in diameter, densely gregarious, often forming a continuous crustaceous layer or scattered and tending to follow the check marks in the wood. Ostiolum papilliform, soon perforated.

Asci cylindrical (p sp.) 9-10 x 70-75 microns.

Spores elliptical, brown, 7.5-8.75 x 11-13.25 microns. E. & E. give asci 10-12 x 60-70 microns. Sporidia 6-9 x 8-15 microns, mostly 7-8 x 10-12.

Common on water beech and sycamore near Jolietville, Ind. Normally found on decorticated wood while it is yet more or less firm.

8. *R. ligniaria* (Grev.) Nke.

Sphaeria ligniaria Grev.

Rosellinia ligniaria Sacc.

Perithecia gregarious or crowded, sometimes forming a crust similar to *R. pulveracea* and in some cases tending to follow the check marks in

the wood. Perithecia ovate-conical, very black and superficial, about $\frac{1}{4}$ mm. in diameter, clothed with very minute black bristles about 20-30 microns in length.

Asci (p sp.) 8-10 x 65-70 microns.

Spores 7-8.75 x 10-14 microns.

Common on Fraxinus and Ostrya near Bloomington, Ind. Found on decorticated wood and underneath loosened bark.

ROSELLINIA AS A PARASITE.

Unlike Hypoxylon and Nummularia, Rosellinia is of great economic importance on account of several of its species being active parasites. Of the one hundred seventy species now described, at least eight are known to be injurious and destructive to living plants. No doubt many other species will be found to be parasitic when a more thorough study is made of them. The following is a brief account of some of the most destructive species:

Rosellinia quercina Hart. Perithecia scattered, seated on a black mycelium, black globose, about 1 mm. in diameter. Asci sub-cylindrical, eight spored, 8-10 x 160-170 microns. Spores brown, acute at both ends, 6-7 x 28 microns.

This species is called the oak root fungus, and attacks the roots of seedling oaks that are from one to three years old. The mycelium spreads rapidly through the ground from one plant to another and is especially destructive during warm, damp weather. This mycelial form was formerly referred to a special genus, Rhizoctonia. The effects of the fungus are first shown by the wilting and drying of the leaves near the top, the lower ones following in order until the whole plant is killed. If a seedling so affected is pulled up and the roots examined, a fine, thread-like mass of white mycelium will be found completely enveloping the roots. The tap root will have dark ovoid bodies about the size of a pin head where the lateral roots join. The tap root is often quite rotten where the mycelium has enveloped it, and especially in the neighborhood of the black tubers. Numerous black sclerotia are found on the surface of the dead roots. The strands of mycelium readily penetrate the young rootlets not yet protected by a layer of periderm and may kill the plant in from ten to fifteen days. Slender hyaline conidiospores are usually found near the base of the stem and on the adjoining soil. Later the perithecia are formed on the dense mass of mycelium covering the superficial roots.

Rosellinia aquila (Fr.) De Not. This species attacks many kinds of trees but is probably best known for its activity on mulberry roots. As a rule the trees are killed by the dense mycelium enveloping the roots. The mycelium penetrates every part of the root and is especially abundant in the medullary rays. When the host is quite dead the dark brown perithecia are found crowded together on the brownish velvety patches that previously bore the conidia. The conidial form of this species is known as *Trichosporium fuscum*. This is one of our most common species, and a description is found elsewhere in this paper.

Rosellinia necatrix (Hart.) Berl. This fungus produces a disease known as the "white root rot." This species is, however, rather rare in this country. One of the peculiar characteristics of the disease is its power of attacking practically every plant with which it comes in contact. It is especially disastrous in vineyards, orchards, etc. Some of its more common victims are vines, fruit trees, oaks, maples, beeches, pines, beans, potatoes and beets.

The mycelium of this fungus travels underground and attacks the rootlets, killing them and gradually working its way up to the larger roots and from them to the body of the plant proper. Here in some instances it breaks through the cortex as a white fluffy mass of mycelium. Sclerotia are formed on the exposed parts of the infected roots, which give rise to dark, bristle-like conidiophores which bear numerous conidia at their tips. Globose swellings on the exposed portions of the mycelium are sometimes formed, which, according to Viala, are capable of emitting mycelium which forms a new plant. The ascigerous stage has been discovered by Viala, appearing only on trees that are well decayed.

Rosellinia radiciperda Mass. This is the cause of the disease known as the "New Zealand white root rot" and is very closely allied to the fungus causing the "white root rot" of Europe, *R. necatrix*. It attacks the roots of the apple, peach, pear, etc., and also such plants as docks, ferns, sorrel, cabbage and potatoes. Sometimes the trees are killed here and there, but often whole areas are swept away.

The white filamentous mycelium attacks the roots and the bark just under the ground. This eventually gives rise to sclerotia which later produce the conidial stage. Next the mycelium becomes dark colored and gives origin to the black globose pycnidia containing stylophores. The ascigerous stage is found on dead tree trunks and stumps that have been dead for considerable length of time.

Rosellinia ligniaria (Nitschke) has been found to attack living ash trees by Mr. W. Carruthers.

Rosellinia massinkii (Sacc.) is reported by Halstead on hyacinth bulbs.

Rosellinia bothrina (B. & Br.) Sacc. is the cause of the tea root rot.

Rosellinia echinata (Mass.) is reported on almost all kinds of trees and shrubs.

The works most freely drawn upon regarding parasitic species are the following:

Hartig—Unters. Forsbot. Inst. Münschen.

Hartig—Diseases of trees.

Massee—Kew Bulletin (1816).

Massee—Diseases of Cultivated Plants and Trees.

Viala—Mon. du Pourridié des Vignes et des Arbres Fruitiers.

Prillieux—Malad. des Plantes.

Wright—Journal Mycol. 5, p. 199.

All species figured and described in this paper have been verified by Dr. Charles H. Peck, or compared with specimens in the New York State Museum by H. D. House, acting State Botanist of New York. I take this opportunity to express my thanks for their assistance.

For valuable material and aid in formulating this paper, I am greatly indebted to Prof. J. M. Van Hook of Indiana University.

Indiana University,

Bloomington, Ind.

EXPLANATION.


A Bausch and Lomb Microtessar 72 mm. lens was used in making all photographs. All figures are twice natural size.

PLATE I.


Figure 1. *R. aquila*.

Figure 2. *R. medullaris*.


Figure 3. *R. mammiformis*.



1



2



3

PLATE II.


Figure 4. *R. glandiformis*.

Figure 5. *R. mutans*.


Figure 6a. *R. subiculata*, old stage showing crowded condition of the perithecia and the absence of the subiculum.



4



5




6a

PLATE III.


Figure 6b. *R. subiculata*, young stage showing scattered perithecia seated in a sulphur yellow subiculum.

Figure 7. *R. pulveracea*.


Figure 8. *R. ligniaria*, showing crustaceous form resembling *R. pulveracea*.



6b



7



8

SOME LARGE BOTANICAL PROBLEMS.

J. C. ARTHUR.

Every farmer, without doubt, desires to produce bumper crops. As agriculture is one of the large factors in national prosperity, it is to the interest of every person in whatever walk of life that bumper crops should be produced. The Experiment Station in each State has been established to assist the cultivator, whether farmer, orchardist, gardener, or any other grower, to solve the problems that hinder maximum production. Some of the problems fall naturally to the botanist. It is well to review them, and not only recognize where the problems lie, but have some idea of their importance. As there are plenty of urgent botanical problems in the home State, it will not be necessary to go outside of Indiana to find illustrative material.

Probably the associated problems that give greatest concern to the cultivator, but which are obscure and little understood, and therefore much in need of study, are the plant diseases which most botanists believe to be connected with soil sanitation. Often they occasion great loss in a crop without the cause being apparent. The soil seems to be all right and proper cultivation has been given, but the plants fail to make their best growth or even dwindle and die. Reference is made to a variety of diseases caused by minute fungi or bacteria, and which attack various garden, truck and field crops. Some instances may be cited without pretending to give them in the order of their importance.

Soil Fungi Attacking Vegetables.—A conspicuous set of diseases, given the name of wilt, is due to certain fungi or bacteria, in which the plants develop normally and may even be bearing, when within a few days they wilt and die as completely as if the roots had been severed. The wilt of watermelons, usually due to a species of *Fusarium*, and of cantaloupes, more often due to bacteria, has often carried off a large part or even all of these crops, so extensively grown in the southern half of the State. The same or similar diseases extend to cucumbers, squashes, and other cucurbits. The partial remedies so far used are rotation of crops and disease-resistant varieties. The germs in certain cases are known to have

remained in the soil in viable condition for eight years without a suitable host in the meantime, and the extent to which rotation of crops serves to check the disease is yet uncertain. The disease-resistant varieties of melon so far provided do not equal the others in flavor, so say our Indiana growers, and are consequently not much used. It is, therefore, still a large problem for the plant pathologist to find means for protecting the melon crop and others of like nature from the attack of such destructive germs.

A similar disease of the tomato has been very injurious at times in Indiana. Mr. W. H. Dyer of Vincennes reports a loss during the one season of 1913 of 6,000 bushels of fruit, some thirty acres of his field being entirely destroyed by the *Fusarium* wilt. Other soil diseases, like leaf spot, fruit rot, etc., also prove destructive at times.

Some of the leaf diseases of potato may be checked by spraying with Bordeaux mixture, while for scab on the tubers a serviceable and practical remedy is known in formaldehyde, showing that some good work has been accomplished along this line. But there are a number of other diseases, often causing heavy loss, that are little understood, and whose method of control is yet to be discovered. Mr. W. A. Orton of the United States Department of Agriculture stated recently before the Wisconsin Potato Growers' Association that six million dollars were lost to the country during the present year of 1914 from potato diseases.

There are a number of damping-off diseases to be classed here. They attack the young plants and cause them to die before becoming established. In the field the disease will spread from plant to plant over large areas. One season a field of beets was reported as largely destroyed in this way. But it is in cutting beds and seed beds under glass where such ravages are most marked.

The growing of vegetables under glass has become a large industry in Indiana. It is a kind of intensive culture where every individual plant bears an important relation to the final profits. Complaints are frequently received of losses in the cucumber and lettuce crops, which prove to be due to the inroads of the fungus, *Sclerotinia libertiana*, sometimes called lettuce-drop.

These diseases, and many similar ones are believed to be harbored in the soil from year to year, and in some cases to be carried from crop to crop by the seed, and possibly by other yet undetected means. How many kinds of germs are concerned in such diseases is not yet known.

They are certainly bewilderingly numerous. The study of any one of them requires much research, and in no case has such a study been fully carried out. Two remedial methods are especially advocated, rotation of crops, and the breeding of disease-resistant varieties. Neither of these methods is yet much or substantially developed. The disinfection of seed and direct destruction of the germs in the soil have yet scarcely been tried out. Spraying of the crop may be found in some cases to serve a good purpose, but so far not much help has been secured in this direction.

Soil Fungi Attacking Forage Crops.—A soil disease, which may become a serious enemy of the alfalfa crop, has been reported from a number of places during the past season. In Clark County, in the southeastern part of the State, eighteen fields were examined during an alfalfa tour, and every one, as reported by Mr. Hutchins, the county agent, was found infested with *Sclerotinia trifoliorum*, a fungus much to be dreaded, that sometimes attacks clover as well. As alfalfa culture is just in its beginning in this State the disease presents an important problem. A banker from another part of the State in writing to the Experiment Station regarding his experience with this disease said in substance: I always thought that the three sure things in this world were death, taxes and alfalfa, but I am afraid that I must leave out alfalfa, possibly quit growing it, on account of the disease.

Other root diseases caused by *Fusarium*, *Rhizoctonia* and *Ozonium*, as well as less severe leaf diseases, are known to damage the alfalfa crop, and may be expected to increase, unless measures are taken to hold them in check. Outside a rotation of crops little has yet been suggested by way of control for any of these pests.

Soil Fungi Attacking Grain.—In seeking the cause why grain, especially wheat, does not yield as well as it once did, and as everyone believes it should continue to do, there is no longer any doubt that much stress should be laid upon soil fungi, especially species of *Fusarium*, *Macrosporium*, *Alternaria*, *Helminthosporium* and *Colletotrichum*. These fungi, sometimes one of them at a time, sometimes more than one, attack the roots of the grain, and work their way into the upper parts of the plant, the stem, leaves and heads. If the attack is light the grain is somewhat less vigorous than it otherwise would be, and the harvest correspondingly lighter, but there is no appearance of disease. *Fusarium* sometimes becomes so active that it causes what is called scab, or pink mold, and

the head of wheat is a total loss, but the soil is usually not suspected as the source of the trouble.

When grain is grown continuously on the same fields the soil tends to accumulate germs of this character. Such an infested soil may be likened to a house long used by sick people without cleaning. What is needed is a thorough disinfection. Unfortunately, at present we know no satisfactory way of disinfecting soils, and we do not even know the kinds and virulence of the diseases to any extent. Our chief reliance so far has been in crop rotation. Possibly something might be done by disinfecting the seed, which is known to harbor the germs. Mr. Hoffer found, as reported in last year's Proceedings of this Academy, that out of thirty-four varieties of wheat examined by him twenty showed germs of this sort, most of them being *Fusarium*.

The subject is a large one of the highest economic importance, and in very great need of study.

Diseases Not Associated With the Soil.—There is no need of specifying where work is needed in this class of troubles, the subject has been before the public too long. Many such diseases are now well understood and efficient remedies already put into use, as in the case of oat and wheat smut, corn smut, and some fruit rots. But there are many rusts, mildews and blights yet to be worked out, and their economic importance rated.

Weeds.—As a rule weeds have not been regarded with sufficient seriousness. But many farmers appreciate the advantages of clean culture, and would like to exterminate the more pernicious kinds. There is a group of weeds, including especially the bindweeds, wild sweet potato, horse nettle and trumpet creeper, that need careful investigation as to the best methods for their control. Field experiments along this line are needed. Horse nettle is now infesting thousands of acres of land in the southern and central parts of the State and is gradually gaining ground northward. Trumpet creeper is becoming so pestiferous on account of its heavy, woody roots, as to make the work of cultivating the infested land a difficult and tedious task. Bindweeds and wild sweet potato are proving to be a bane to corn culture, especially on bottom lands. Large fields can be found where nearly every corn plant is entwined by some one of these kinds of weeds. It is safe to say that in such cases the crop yield is reduced by at least one-third. Aside from reducing the yield such weeds also increase the cost of field operations and harvesting and in other ways depreciate the value of land.

It is quite possible that suitable investigation would produce as marked and practical results in the case of most of these weeds, as has been reached in the recent study of the wild garlic situation in southern Indiana. An intensive study of this weed has resulted in a special method of eradication that promises to be a boon to owners of land where garlic has been established, and to render its complete extermination entirely feasible.

Poisonous Plants and Molds.—The subject of plants and molds, which are or may be poisonous to stock needs careful investigation. Many plants and various molds have been suspected of poisonous action of which there is little definite knowledge available. In the majority of cases the evidence is based on reports obtained from farmers, with no scientific investigation or actual tests to support them.

The botanical department of the Experiment Station often receives letters, with specimens of plants, stating that those particular kinds of plants are believed to have been responsible for the death of several head of stock, usually cattle. It is known that some of the wild larkspurs of the State are poisonous, and that loss of stock has been due to them, but no very exact information is at hand.

Certain silage molds have been under suspicion, and moldy corn is quite definitely known to act in a poisonous way.

The importance of careful study of authenticated cases of poisoning from the botanical standpoint will not be gainsaid.

Conclusion.—Probably enough has been said by way of citing examples, and pointing out lines of work to show that there is much valuable investigation for the benefit of the cultivator that falls naturally to the trained botanist. Most such work must be done by the technical man who is supplied with suitable equipment. That the State should properly provide the means for carrying forward work of this character seems to require no argument.

*Purdue University,
Lafayette, Indiana.*

THE ALBA B. GHERE COLLECTION OF BIRDS' EGGS PRESENTED TO THE MUSEUM OF PURDUE UNIVERSITY.

HOWARD E. ENDERS.

Alba B. Ghere of Frankfort, Indiana, was born in 1870 and died March 25, 1904, at the age of thirty-four years.

It is said of him that he was a born naturalist, an enthusiastic student of birds, their eggs and their nests. To this line of endeavor he gave all of the spare time of his life from the fourteenth year to the end. A careful list of the spring and fall migrations was recorded for a period of years, and at the end of each season was forwarded to the U. S. Department of Agriculture for record. The incomplete list of the spring migration of 1904, in which he entered notes to a day or two before his death, accompanies the collection of eggs.

That Mr. Ghere was a careful observer is attested to by the fact that his notes, among those of many other persons, are quoted at five or more points by Dr. C. H. Merriam in the Bulletin No. 1, on The English Sparrow in North America (1889), Division of Economic Ornithology and Mammalogy, Washington, D. C.

Through the kindness of his mother, Mrs. India S. Cottingham, of Frankfort, Indiana, the complete collection of eggs, the oölogical instruments, and his oölogical books were presented to Purdue University in October, 1914.¹ The collection numbers somewhat more than four hundred specimens of eggs from about one hundred and thirty species of birds, some of which are western or marine species that were acquired by exchange or by purchase. It may be of special interest to know that one egg is said to be from the once abundant passenger pigeon.²

¹Among the books are government publications on birds; Davis and Baker's *Oologist's Directory*, 1885; Davies' *Nests and Eggs of North American Birds*, 1886; Capen's *Oology of New England*, 1886. The latter is an elaborate quarto in natural colors, and is one of a limited small edition.

²Its measurements and diagnostic characteristics seem to correspond with the descriptions given for this species.

199. Louisiana Heron.....Hydranassa tricolor ruficollis.
 200. Little Blue Heron.....Florida coerulea.
 201. Green Heron.....Butorides virescens.
 202. Black-crowned Night Heron.....Nycticorax nycticorax naevius.

Order VIII. PALUDICOLÆ.

Family *Rallidæ*.

212. Virginia Rail.....Rallus virginianus.
 214. Carolina Rail; Sora.....Porzana carolina.
 218. Purple Gallinule.....Ionornis martinica.
 221. American Coot.....Fulica americana.

Order IX. LIMICOLÆ.

Family *Phalaropodidæ*.

224. Wilson Phalarope.....Steganopus tricolor.

Family *Scolopacidæ*.

261. Bartramian Sandpiper.....Bartramia longicauda.
 263. Spotted Sandpiper.....Actitis macularia.

Family *Charadriidæ*.

269. Lapwing.....Vanellus vanellus.
 274. Semipalmated Plover.....Aegialitis semipalmata.
 280. Wilson Plover.....Oethodromus wilsonius.

Order X. GALLINÆ.

Family *Tetraonidæ*.

289. Bob-white.....Colinus virginianus.

Order XI. COLUMBÆ.

Family *Columbidæ*.

315. †Passenger Pigeon; Wild Pigeon...Ectopistes migratorius.
 319. White-winged Dove.....Melopelia leucoptera.

Order XII. RAPTORES.

Family *Cathartidæ*.

325. Turkey Vulture; Turkey Buzzard..Cathartes aura.

†Its measurements and diagnostic characteristics seem to correspond with the descriptions given for this species.

Family *Falconidæ*.

337. Red Tailed Hawk..... *Buteo borealis*.
 339. Red-shouldered Hawk..... *Buteo lineatus*.
 357. Pigeon Hawk..... *Falco columbarius*.
 360. Sparrow Hawk..... *Falco sparverius*.
 364. American Osprey; Fish Hawk..... *Pandion haliaetus carolinensis*.

Family *Strigidæ*.

365. Barn Owl..... *Strix patrincola*.

Family *Bubonidæ*.

366. American Long-eared Owl..... *Asio wilsonianus*.
 367. Short-eared Owl..... *Asio accipitrinus*.
 375. Great Horned Owl..... *Bubo virginianus*.

Order XIV. COCCYGES.

Family *Cuculidæ*.

387. Yellow-billed Cuckoo..... *Coccyzus americanus*.

Family *Alcedinidæ*.

390. Belted Kingfisher..... *Ceryle alcyon*.

Order XV. PICI.

Family *Picidæ*.

- ?? Undetermined species.
 412. Southern Flicker..... *Colaptes auratus*.
 412a. Northern Flicker..... *Colaptes auratus luteus*.

Order XVI. MACROCHIRES.

Family *Caprimulgidæ*.

420. Nighthawk..... *Chordeiles virginianus*.
 425. White-throated Swift..... *Aeronautes melanoleucus*.

Family *Trochilidæ*.

428. Ruby-throated Hummingbird..... *Trochilus colubris*.
 429. Black-chinned Hummingbird..... *Trochilus alexandri*.
 431. Anna Hummingbird..... *Calypte anna*.

Order XVII. PASSERES.

Family *Tyrannidæ*.

- ?? Flycatcher.
 461. Wood Pewee.....Contopus virens.
 466a. Alder Flycatcher.....Empidonax traillii alnorum.
 467. Least Flycatcher.....Empidonax minimus.

Family *Alaudidæ*.

- ?? Larks, undetermined.

Family *Corvidæ*.

488. American Crow.....Corvus americanus.

Family *Icteridæ*.

497. Yellow-headed Blackbird.....Xanthocephalus xanthocephalus.
 ??? Oriole.....Undetermined.
 506. Orchard Oriole.....Icterus spurius.
 507. Baltimore Oriole.....Icterus galbula.

Family *Fringillidæ*.

- 519a. Crimson House Finch.....Carpodacus mexicanus frontalis.
 546. Grasshopper Sparrow.....Coturniculus savannarum passerinus.
 552a. Western Lark Sparrow.....Chondestes grammacus strigatus.
 584. Swamp Sparrow.....Melospiza georgiana.
 585. Fox Sparrow.....Passerella iliaca.
 598. Indigo Bunting.....Cyanospiza cyanea.
 601. Painted Bunting; Nonpareil.....Cyanospiza ciris.

Family *Hirundinidæ*.

612. Cliff Swallow.....Petrochelidon lunifrons.
 613. Barn Swallow.....Hirundo erythrogastra.

Family *Ampelidæ*.

619. Cedar Waxwing.....Ampelis cedrorum.

Family *Vireonidæ*.

631. White-eyed Vireo.....Vireo noveboracensis.
 ?? Unidentified species.

Family *Mniotiltidæ*.

683. Yellow-breasted Chat.....Icteria virens.

Family *Troglodytidae*.

703. Mockingbird.....Mimus polyglottos.
 705. Brown Thrasher.....Toxostoma rufum.
 713. Cactus Wren.....Heleodytes brunneicapillus.
 718. Carolina Wren.....Thryothorus ludovicianus.
 721. House Wren.....Troglodytes aedon.
 722. Winter Wren.....Olbiorchilus hiemalis.
 724. Short-billed Marsh Wren.....Cistothorus stellaris.
 725. Long-billed Marsh Wren.....Telmatodytes palustris.

Family *Turdidae*.

758. Russet-backed Thrush.....Hylocichla ustulata.
 761. American Robin.....Merula migratoria.
 761a. Western Robin.....Merula migratoria propinqua.
 766. Bluebird.....Sialia sialis.
 767. Western Bluebird.....Sialia mexicana.
 768. Mountain Bluebird.....Sialia arctica.

Unidentified, about thirty species, chiefly of the common native birds.

Purdue University,

West Lafayette, Ind.

A NOTE ON A PECULIAR NESTING SITE OF THE CHIMNEY SWIFT.

GLENN CULBERTSON.

As an illustration of the ability of birds to adapt themselves to new conditions, the chimney swift is a striking example. Driven from the hollow tree as a nesting site by the woodman's axe and fires, the swift adapted itself to the broad open chimneys of the settlers' cabins, and later to the narrow flues of a later day. The projecting spines of the tail feathers fortunately answered the same purpose in a soot-lined chimney that they had done in the soft decayed wood of a hollow tree.

During the past summer the writer's attention was called to a still greater change in the nesting site of a pair of swifts. Near the residence of Mr. James Storie, one and one-half miles north of Moorefield, Switzerland County, a pair of swifts, being excluded from the chimneys by wire netting, have nested for two seasons in an old-fashioned dug well walled with stone.

The well is some twenty feet deep and three feet in diameter, and has over it a square curb about three feet high, one-half of which was permanently left open. The nests were built in each case at a distance of some seven or eight feet below the level of the ground, and at approximately the same distance above the water. The young were matured and brought forth in safety both seasons.

NOTES ON INDIANA EARTHWORMS.

H. V. HEIMBURGER.

The great group of animals known commonly as earthworms, comprises four families of the order Oligochaeta: the Moniligastridæ, the Megascolecidæ, the Glossoscolecidæ and the Lumbricidæ. Three of these families are represented in our fauna.

At the present time there are more than a thousand well recognized species of earthworms known. By far the larger number of these species have been described during the last twenty years. Most of the work on this group has been done by Europeans, chief among whom is Michaelsen of Hamburg, whose publications make up the bulk of the recent literature on the group. Professor Frank Smith of the University of Illinois has worked on the group in this country and Eisen, a Swede who lived for several years in San Francisco, has worked on West and Central American species.¹ Very little is known of the earthworm fauna of the Central States. Only about fifty species are known to occur in the United States, this scarcity of forms being due in part to lack of study and partly to the actual scarcity of species.²

It has been pointed out by Beddard³ and Michaelson⁴ that the earthworms are an important group for the Zoögeographer: Arldt,⁵ in a recent paper, shows the value of the group for the paleo-geographer and gives many of the facts of distribution now known as well as indicating the value of the group in theoretical considerations within the fields of geography and geology.

The Moniligastridæ is a small family limited to Borneo, Ceylon, Southern India and neighboring islands. This family is regarded as the most ancient of the group.

¹Eisen; American Oligochaetes with special reference to those of the Pacific Coast and Adjacent Islands. Proceedings Calif. Acad. Science Vol. II, No. 2, 1900.

²Frank Smith; Earthworms From Illinois. Trans. Ills. Acad. Science. 1912.

³Beddard; A Textbook of Zoögeography. Cambridge, 1895.

⁴Michaelsen; Die geographische Verbreitung der Oligochaeten. Berlin, 1903.

⁵Ardt; Die Ausbreitung der terricolen Oligochaeten in Laufe der erdgeschichtlichen Entwicklung des Erdreliefs. Zool. Jahrb. Abt. f. Syst. Geog. u. Biol. Bd. 26, pp. 285-318. 1908.

The Megascolecidae are supposed to have arisen from the same root as the Moniligastridae but at a later period, and dates back at least to the Triassic. This is the largest of the earthworm families and contains more than half the known species. The family is widely distributed, chiefly in tropical regions and the southern hemisphere. But one genus, *Diplocardia*, is represented in our fauna. According to Michaelsen, this genus probably appeared in Mexico or Central America during the Jurassic. Derivatives from this genus have spread into Africa, but the genus itself spread northward and is known from Mexico, Lower California, Texas, Florida, Nebraska and Illinois.

The Glossoscolecidae seem to have developed as early as the Jurassic in the northern continental area. The genus *Sparganophilus* of this family is found in Mexico and various parts of the United States. Related forms are known from Central and South America, where many species are recognized.

The Lumbricidae is recognized as the most recent family of the group and is derived from the Glossoscolecidae, probably in southwest Asia. The family is thought to have invaded Europe in the Eocene and North America in the Oligocene. But few endemic species are known from the United States.

In the glaciated regions of the world, it is probable that the endemic species have been destroyed during the Ice Age. These regions have been repopulated by species which have migrated from the south and the earthworm fauna in such places is largely composed of forms carried in by man. In the southern part of Europe are found many endemic species while northern Europe is occupied almost wholly by forms also found further south. The line separating the northern territory with peregrine forms from the southern territory with endemic forms, corresponds very closely to the line of the most southern extension of the glacial ice sheet. It would be interesting to know if a similar condition exists in America.

Last year, under the direction of Professor Frank Smith of the University of Illinois, I began a study of the earthworms of Illinois and Indiana. I received, last fall, some material from Mr. C. E. Allen, of Wabash College, and some from my brother at Kewanna. During the past summer I made some collections in several counties of the State but was unable to make as extensive collections as are desirable. The materials I have contain some forms that may have to be described as new species and I believe that careful collecting in the State will disclose

several species new to science. It is my intention to make several collecting trips through the State during June and July, 1915, but the task of making a complete collection for the State is not to be completed in a few weeks by one man.

Earthworms are easy to collect and no great difficulty is experienced in caring for the collections. The most interesting forms are to be expected in uncultivated areas such as woodlands, stream banks, and the margins of swamps and lakes. Many interesting forms may be taken under logs and under the bark of old logs. The worms are especially easy to collect in the spring when driven from their burrows by the heavy rains. At such times they may be picked up in large numbers from roads and sidewalks. I wish to gain a more extended knowledge of the distribution of *Diplocardia* in the State and to know what forms are found in the unglaciated areas of the State. If any members of the Academy will aid in securing materials, I shall be very glad to have the material and to return named specimens in exchange. I shall be glad to correspond with any one who may be interested in such work.

Following is a list of species I have taken in Indiana together with some notes as to the habitats in which the worms were found. The space of this paper is not sufficient for descriptions of the species. The monograph by Michaelsen is the most authentic single work on the group.¹ The nomenclature used in this list is that of Michaelsen's monograph, except where the nomenclature has been modified in his later papers.²

Family MEGASCOLECIDÆ.

Genus *Diplocardia* (Garman).

1. *Diplocardia communis* var. *typica* Garman.

This form is quite common about Urbana and in other parts of Illinois. I have not found it in Indiana.

D. communis var. *singularis* Ude.

Collected under logs in recently cleared land near Culver, Marshall County. I have other specimens collected in Putnam and Vigo counties, which are very similar to *singularis* but very much larger and differing in minor points.

¹Michaelsen; Oligochæta. Vol. 10 in Schulze's Thierreich. Berlin, 1900.

²See particularly.

Michaelsen; Zur Kenntnis der Lumbriciden und ihrer Verbreitung. Ann. Zool. Mus. Imp. Acad. Sciences, St. Petersburg. 1910.

2. *Diplocardia riparia* F. Smith.

Collected at Terre Haute in a wooded pasture land.

3. *Diplocardia udei* Eisen.

Collected at Terre Haute together with *riparia*. The specimens are somewhat larger than Eisen's species and differ in details of setal modifications. If not *udei* this form is probably to be regarded as a variety of *udei*. The species was described from North Carolina and has not been reported outside that State.

I have other specimens of *Diplocardia* from Indiana, of which the classification is still uncertain.

Family GLOSSOSCOLECIDÆ.

4. *Sparganophilus eiseni* F. Smith.

This species is amphibious, and is found in very wet mud or among the roots of aquatic plants. Collected at Culver, on roots of Eel grass and in mud at margin of Lake Maxinkuckee. Very abundant in mud of small stream near Kewanna, Fulton County.

Family LUMBRICIDÆ.

5. *Helodrilus tetradrus* var. *typicus* Sav.

Collected at Culver, in vegetable drift at edge of lake. An amphibious species found usually in wet soil along margins of streams or among vegetable detritus in very moist places.

6. *Helodrilus roseus* Sav.

Collected at Culver, Greencastle, Terre Haute. This species is probably to be found in all parts of the State. It is very commonly found in lawns and cultivated fields. My specimens were collected in woods and along stream banks.

7. *Helodrilus constrictus* Rosa.

Collected at Kewanna, under logs at handle factory; Culver under logs in woods.

8. *Helodrilus subrubicundus* Eisen.

Collected at Culver and Kewanna, under logs. The identification of this species is not absolutely certain as no sections have been made. However, this form is fairly well determined from external characters.

9. *Helodrilus chloroticus* Sav.

Collected at Culver, Lake Maxinkuckee, in moist soil at edge of lake. Greencastle, in moist clay soil in abandoned quarry hole. Crawfordsville, in moist clay soil, banks of small stream.

10. *Helodrilus foetidus* Sav.

Collected at Kewanna and Culver. This is the common evil smelling, barnyard or manure worm. Collected at Culver in decaying straw near ice houses. At Kewanna this worm was found in large numbers, in moist soil along a stream just below the outlet of a sewer. The whole locality was quite offensive because of the sewage.

11. *Helodrilus caliginosus* Sav.

Collected in Fulton, Marshall, Putnam, Madison and Vigo counties. This is perhaps our most cosmopolitan species. It is a European species that has spread wherever Europeans have settled. Perhaps 90% of any random collection in the State would consist of this species.

12. *Helodrilus longus* Ude.

Collected at Greencastle, in woods.

13. *Helodrilus zeteki* Smith and Gittins. (Mss.)

Collected at Kewanna, Culver, Summitville. All of my specimens were collected in loose sandy loam under logs, and under the bark of logs. In June I took about thirty-five specimens of adult worms in a woods near Kewanna. At this time I obtained many cocoons of the species, the worms being in the height of sexual maturity. This species was described by one of Professor Smith's students from specimens collected in a woods near Urbana, Ill. The description of the species is not yet in press.

14. *Octolasion lacteum* Oerley.

Collected at Crawfordsville, Culver, Greencastle, Summitville, Kewanna. My Crawfordsville specimens were sent me by C. E. Allen, and were collected in the banks of Sugar Creek. This species is very common under logs and in moist soil everywhere I have collected. It is very widely distributed throughout the world. I have never found it in cultivated fields.



NOTES ON ORTHOPTERA AND ORTHOPTERAN HABITATS IN THE VICINITY OF LAFAYETTE, INDIANA.

HENRY FOX.

(Assistant, Cereal and Forage Insect Investigations,
U. S. Bureau of Entomology.)

Between September 3, 1912, and November 30 of the following year I was stationed in pursuance of official duties at Lafayette, Tippecanoe County, Indiana. At intervals during my stay there I made a series of observations on the *Orthoptera* and Orthopteran habitats of the surrounding country which, incomplete as they are, nevertheless constitute a distinct positive contribution toward an accurate knowledge of the faunal features of the region. My earlier studies on the distribution of *Orthoptera* in Pennsylvania and New Jersey¹ had impressed me with the importance of detailed local lists of species in a scientific study of distribution. The usual distribution as given in most works of reference is entirely too general for accurate study, no regard being paid to local peculiarities of distribution or to the relative abundance of the species in different parts of its range. Take, for example, such a form as *Psinidia fenestralis*. Its range, as usually given, extends from Massachusetts to Florida, Texas, northern Indiana and southern Minnesota. Such a statement would incline one to think that the entire region south of say a line drawn from Cape Cod to the southern extremities of the Great Lakes and thence to the southern border of Minnesota would be characterized by the presence of this species. As a matter of fact such is very far from being the case. In the East, for instance, *Psinidia fenestralis* is regularly found only in the low sandy belt fringing the coast, while in the interior it is of extremely local occurrence, being met with only on widely scattered, isolated deposits of loose sand. All positive data on this species indicates that its distribution is conditioned by the presence of areas of loose sand.

¹Data on the Orthopteran Faunistics of Eastern Pennsylvania and Southern New Jersey. Proc. Acad. Nat. Sci., Phila., 1914, pp. 441-534.

Wherever these are lacking the species is absent. In southern New Jersey where sandy deposits are of practically universal occurrence *Psinidia* is common, while on the opposite side of the Delaware River in Pennsylvania it is unknown. A case like this shows us how essential it is that we should have accurate local data before we can be certain of the exact range of a species.

The studies of Morse, Hancock, Rehn, Hebard, Vestal and others have clearly shown the intimate relation between the distribution of numerous *Orthoptera* and certain features of the environment. As Shelford has so well pointed out the success or failure of a species in any place will depend on how closely the environmental complex approximates to that condition at which the normal physiological activities of the species can be carried on to the best advantage. Where this state of affairs obtains the species will attain its maximum abundance; if one or more of the factors of the environmental complex are less favorable it will be present in diminished numbers, while if any essential factor is prohibitive the species will be absent. It is the aim of biogeography to explain the facts of organic distribution in terms of physiology, as an expression of the reactions of organisms to the varying conditions of their surroundings. In the case of the *Orthoptera* this can be done only when we know much more than we do now about the intrinsic qualities of the species and their ability to accommodate their activities to varying intensities of environmental factors. To acquire such knowledge will require much experimental investigation. In the absence of such knowledge we must meanwhile be content to record the facts of distribution as actually observed and to point out any correlation which may exist between the range of species and the different types of environments. By the accumulation of data along these lines a good foundation will be laid for the ultimate causal interpretation of distribution and kindred biological problems.

In the present article I have endeavored not only to give a full list of the species observed about Lafayette, but in addition to point out the more evident physical and botanical features of the region with which the local distribution of the *Orthoptera* is correlated. Most of the facts here given were gathered by myself, but I am also indebted for some valuable additional data to Mr. P. W. Mason, Instructor in Entomology at Purdue University, whose kindness in placing his notes at my disposal I here take pleasure in acknowledging.

GENERAL DESCRIPTION OF THE REGION.

Most of my observations were made on the west side of the Wabash River extending from Battle Ground on the north to the mouth of Indian Creek on the south. (For these and other localities a good map to consult is the map accompanying the report of the soil survey of Tippecanoe County in the field operations of the U. S. Bureau of Soils for 1905.) On the east side of the river a few observations were made from the mouth of Wild Cat Creek, about two miles north of the city, to a spot east of Battle Ground, about a half mile south of the mouth of Buck Creek. Only one trip was taken on the east side south of Lafayette. It was limited to the line of the Wabash Railroad and extended about three miles below the city.

According to the Bureau of Soils' report on the soils of Tippecanoe County the general altitude of the country is about 750 feet above sea level. Back from the Wabash River and its tributaries the country forms a nearly level, or at most slightly undulating, plain. Near the river it is much more rugged, a relatively steep line of bluffs leading down to the valley of the Wabash which is about 100 feet below the general level of the upland. Similar conditions prevail along the main tributaries, such as Burnett, Wild Cat and Indian Creeks.

The valley of the Wabash forms a nearly level tract varying according to location from a half to two miles in width. It is formed of what are known as bottom lands, or more specifically "first bottoms" to distinguish them from the older bottoms which are no longer covered by the overflow from the river. The surface of these first bottom lands is according to the report already mentioned between 10 and 20 feet above low water mark. They are "subject to overflow during periods of high water." During the destructive floods of March, 1913, these bottom lands were completely submerged.

The margin of these bottom lands is formed by the line of steep bluffs already mentioned as forming the edge of the upland. Locally, as is the case in the vicinity of West Lafayette and of Battle Ground, these bluffs recede a mile or two back from the river and in the embayments thus formed "second bottoms" are developed, that is, "fossil" floodplains or terraces representing an earlier, prehistoric stage of deposition. The surface of these "second bottoms" is level or slightly rolling and on the side facing the river is marked by a gentle slope rising from forty to

fifty feet above the present bottoms. They are never covered by overflow from the river at the present time.

The whole region about Lafayette is deeply buried under glacial deposits, the depth of these deposits being usually very great (at least 150 feet), though in limited areas they may be quite thin or lacking. Only rarely, however, do the underlying Palæozoic limestones reach the surface. One such outcrop I have seen on the upland near Montmorenci where the Lake Erie and Western Railroad crosses Indian Creek. Outside of these rare and insignificant cases, the whole country is underlaid by a very coarse glacial gravel. Overlying this is usually a layer of loess varying in thickness from an inch to several feet. From this loess are derived the representative soil types of the region.

The drainage of the region is in general good. The streams are few and in periods of protracted drought frequently dry up entirely in their upper courses. Most of the rainfall, however, is carried off by underground drainage, the underlying gravel allowing the ready percolation of water. Locally, as in upland swales and depressions and at the base of the river bluffs, where the seepage of underground water takes place, the ground is, except in seasons of drought, more or less completely saturated with water resulting in the formation of swamps. At the present time most of these naturally wet areas, especially on the upland, have been artificially drained and the land utilized for growing crops. The bottom lands are at present well drained, the cultivation of the soil breaking it up into a loose condition which allows the water to flow off readily beneath the surface.

On the upland the dominant soil is a fine-grained, silty loam, varying in color from light brown to almost black, the color depending upon the amount of organic matter present, which is usually considerable. Of this soil the Bureau of Soils recognizes two categories which are termed respectively Marshall silt loam and Miami silt loam. Both are nearly alike in mineral content, being characterized by relatively high per centage of silt and clay and extremely low per centage of sandy constituents, but differ in their organic content, the Miami being as a rule much poorer in this respect than the Marshall. The table shows the mechanical composition of the soils, the data being taken from the Bureau of Soils report.

SOIL.	Fine Gravel.	Coarse Sand.	Medium Sand.	Fine Sand.	Very Fine Sand.	Silt.	Clay.
Marshall Silt Loam.....	0.2	1.1	1.3	3.9	5.7	67.2	20.0
Miami Silt Loam.....	0.4	1.8	1.0	2.2	7.3	68.4	18.9

On the "second bottoms" the soils contain a much greater percentage of sand and are correspondingly poor in silt and clay. The quantity of organic matter is variable depending upon location and drainage conditions. Typical examples of "second bottom" soils are the Sioux sandy loam and Miami sand the composition of which, as given by the Bureau of Soils, is shown in the following table:

SOIL.	Fine Gravel.	Coarse Sand.	Medium Sand.	Fine Sand.	Very Fine Sand.	Silt.	Clay.
Sioux Sandy Loam.....	2.2	9.8	10.2	28.5	10.3	27.1	11.2
Miami Fine Sand.....	.0	3.0	13.3	52.0	11.5	13.8a	6.4

The characteristic soil of the river bottoms is the Wabash silt loam. This is the material deposited during periods of high water. It resembles the upland soils in its high silt-clay content, but differs in having a considerable percentage of sand. The mechanical composition of this soil as given by the Bureau of Soils is as follows:

SOIL.	Fine Gravel.	Coarse Sand.	Medium Sand.	Fine Sand.	Very Fine Sand.	Silt.	Clay.
Wabash Silt Loam.....	0.0	Trace.	0.3	27.0	2.1	66.1	28.4

These, as well as the other soils of the region not herein specifically mentioned, are all characterized by their prevailing fine texture, a rule that holds even in the case of the sandy soils in which the major constituent is the fine sand, so that, in spite of the rapid drainage afforded

by the underlying boulder drift, the capacity of the soil to retain moisture is quite high. For this reason all the soils are very productive and are in consequence in a high state of cultivation. About Lafayette, with insignificant exceptions, practically all the land is under cultivation. On the upland the principal crops are corn, oats, clover and wheat, while the bottom lands form one unbroken stretch of corn. The only waste places—oases for the naturalist—are on the upland an occasional grove or more rarely a swampy depression, on the bottoms frequent, though small, bogs marking the places where the underground waters ooze out from the marginal bluffs. In such places the rarer and more interesting *Orthoptera* are to be found.

ORTHOPTERAN HABITATS.

No attempt at an exhaustive study of the various Orthopteran habitats was made owing to the limited time that could be spared for that purpose. Consequently in the following pages only the grosser features of the habitats are mentioned. About Lafayette, owing to the intense cultivation of the region, nearly all the country is open, in consequence of which the dominant *Orthoptera* are campestral types. Where the ground is unfilled it is usually covered with a close growth of blue grass (*Poa pratensis*), which in damper spots is replaced by foxtail (*Chatochloa viridis* and *glauca*). In such situations the grasshoppers usually encountered include the following species:

Syrbua admirabilis, *Arphia xanthoptera*, *Chortophaga viridifasciata*, *Encoptolophus sordidus*, *Dissosteira carolina*, *Melanoplus atlantis*, *Melanoplus femur-rubrum*, *Orchelimum vulgare*, *Conocephalus strictus* and *Nemobius fasciatus*.

In cultivated lands this assemblage is largely characteristic of the grassy borders of roads, paths and fence-rows. Most of the species named continue abundant in such places with the possible exception of *Arphia xanthoptera* and *Chortophaga viridifasciata*, both of which appeared to be rather scarce in the particular cultivated tracts examined by me.

A second group of *Orthoptera* is characteristic of dry upland woods. On the level uplands woodland is represented only by widely scattered groves, in most of which the trees have been thinned out. This allows a rich growth of blue grass which is largely utilized as pasturage for cattle. Such pastured woodlands are almost invariably very barren in *Orthoptera*, those that do occur being similar to those found in the open

country. The most nearly continuous and undisturbed areas of woodland are those which clothe the tops and sides of the bluffs which, as already mentioned, form the outer margins of the river-bottoms. These are exclusively hardwood formations, the dominant tree at higher levels being the white oak (*Quercus alba*), with which are commonly associated the sugar maple (*Acer saccharum*), pig-nut hickory (*Hicoria glabra*), red oak (*Quercus rubra*), shell-bark hickory (*Hicoria ovata*), bass-wood (*Tilia americana*), elm (*Ulmus* sp. not det.), beech (*Fagus ferruginea*), dogwood (*Cornus florida*) and aspen (*Populus tremuloides*). Wherever these woodlands are sufficiently open to admit sunlight blue grass usually springs up and forms a continuous cover to the ground or, if the soil is exceptionally dry, an aggregation of more or less scattered tufts with interspaces of bare earth. Where the grass is thick one usually finds *Melanoplus scudderi*, while in places where it is short and scattered *Spharagmon bolli* and *Melanoplus luridus* are usually encountered. Along the edges of the woods in undisturbed ground these more strictly sylvan types were observed to meet and to intermingle with a campestral assemblage which usually included *Syrbula admirabilis*, *Arphia xanthoptera*, *Chortophaga viridifasciata* and *Encoptolophus sordidus*. In scrubby areas and in tall herbaceous growths *Atlanticus testaceus* was fairly common.

In strong contrast to the foregoing group is an assemblage characteristics of moist areas. Such areas most frequently occur at the outer margin of the river bottoms where the seepage from the neighboring bluffs keeps the ground perpetually moist and soggy. The soil in such places is a typical muck, frequently intermixed with gravel and silt. In nearly all the swamps I have visited the vegetable content of the soil appeared to be thoroughly decomposed. At one place (1) in a wet depression in the midst of a fairly large woods on the upland about one and a half miles northwest of West Lafayette the substratum was a true peat. In the bottomland swamps, however, the soil appears in all cases where I have examined it to be a muck. Such a swamp harbors a rich vegetation of which the dominant member in wetter spots is rice cutgrass (*Homalocenchrus oryzoides*) with which are often associated cat-tails (*Typha latifolia*) and jewel-weeds (*Impatiens biflora*). Surrounding the cutgrass areas in slightly dryer ground is usually a dense thicket composed of tall herbaceous plants, especially composites, among which I noted the taller ragweed (*Ambrosia trifida*), ironweed (*Vernonia fasciculata*), joe-pye-

weed (*Eupatorium purpureum*), boneset (*Eupatorium perfoliatum*) and a bewildering variety of members of the sunflower tribe (*Helianthus*, *Bidens*, etc.). Where this thicket is sufficiently open there frequently occur patches of wild rye (*Elymus virginicus*, *E. canadensis*).

The central portion of these swamps dominated by *Homalocenchrus* appeared to be characterized by a rather different assemblage of Orthoptera than that typical of the surrounding thickets, though, owing to the usually restricted size of the swamps, it was not possible in all instances to clearly distinguished the two groups. In general, however, the *Homalocenchrus* areas appeared to be characterized by such Orthoptera as *Orchelimum nigripes*, *Neoconocephalus palustris*, *Stauroderus curtippennis*, *Conocephalus attenuatus* and *Paroxya hoosieri*. The surrounding thickets were especially characterized by the short-winged *Melanopli*, such as *Melanoplus obovati-pennis*, *M. scudderi*, *M. gracilis* and *M. viridipes*, together with numerous examples of *Melanoplus differentialis*, *Conocephalus nigropleurum* and *Conocephalus memoralis*. Two forms that appeared to occur indifferently in both zones were *Orchelimum vulgare* and *Conocephalus brevippennis* (incl. *ensifformis*).

RELATIVE FREQUENCY OF THE SPECIES.

As regards numbers the most abundant grasshopper in this region is *Melanoplus femur-rubrum* which appears to swarm everywhere on both upland and lowland, though it appeared to be less frequent in wooded areas than in more open situations. Next to it in point of numbers I would place *Melanoplus atlantis* which is common, but more local than *femur-rubrum*. Other species which appeared to be present in what may be regarded as abundance were *Encoptolophus sordidus*, *Dissosteira carolina*, *Melanoplus differentialis*, *Orchelimum vulgare*, *Conocephalus strictus* and *Nemobius fasciatus*. Much less frequent, but on the whole rather common were such species as *Syrbula admirabilis*, *Arphia xanthoptera*, *Chortophaga viridifasciata*, and *Melanoplus femoratus*. Some species appeared to be of frequent or regular occurrence locally wherever the special conditions making up their normal environment prevailed. Thus *Spharagemon bolli* and *Melanoplus scudderi* and *luridus* occurred, usually in considerable numbers, wherever there were dry open woodlands, while in the swamps, or their borders, three species, *Melanoplus differentialis*, *Orchelimum nigripes* and *Conocephalus brevippennis* were in all but one or two instances abundant. Associated with the last three were frequently

considerable numbers of *Dichromorpha viridis*, *Sturoderus* (*Stenobothrus*) *curtipennis*, *Melanoplus scudderi*, *Melanoplus obovatipennis*, *Melanoplus femoratus*, *Conocephalus fasciatus*, and *Conocephalus nigropleurum*. Certain species were scarce in most places, but were found to be common or even abundant in one or two restricted areas. Thus *Hippiscus rugosus* was found in only one place, but was there quite common. *Paroxya hoosieri* was taken in numbers in a swamp (16) in the Wabash bottoms opposite Battle Ground but was not observed elsewhere. A peculiar variety of *Orchelimum nigripis* and *Conocephalus attenuatus* literally swarmed in a boggy depression (14) on the upland about 2 miles northwest of West Lafayette. The former variety I did not find in any other place, while of the latter I noted elsewhere only a single individual which I captured in a bog in the Wabash bottoms (6) about half a mile south of Lafayette.

Certain species were observed to be of rather infrequent occurrence but could hardly be called rare. Among these were *Schistocerca americana*, *Melanoplus viridipes*, *Melanoplus gracilis*, *Scudderia texensis*, *Scudderia furcata*, *Neoconocephalus palustris*, *Conocephalus nemoralis* and *Atlanticus testaceus*. The following species appeared to be quite scarce: *Truxalis brevicornis*, *Orphulella speciosa*, *Chlœaltis conspersa*, *Schistocerca alutacea*, *Melanoplus walshii*, *Neoconocephalus robustus crepitans* and *Conocephalus saltans*.

DESCRIPTION OF LOCALITIES WHERE COLLECTIONS WERE MADE.

1. A fairly extensive bit of woodland on the edge of the upland about a mile northwest of West Lafayette. The timber was in part rather dense, but there were a number of open spots well fitted for sylvan Orthoptera. There had been no grazing in the portion of the woods where the collecting was chiefly done, so there was considerable undergrowth. Most of the land which these woods covered was dry or only moderately humid, but it included one or two depressions where the ground was either soggy or covered with standing water. One of these, a very limited tract, was included in the northwestern edge of the wood and was occupied by an almost pure growth of button-ball bush (*Cephalanthus occidentalis*); the other was slightly larger and occupied by a mixed growth of sapling silver maples (*Acer saccharinum*) and red-berried elder bushes (*Sambucus racemosa*) together with a variety of other plants. Both of these swampy areas proved to be quite barren in Orthoptera. The best collecting was done along a path entering the woods at its northwest corner and in the

neglected clearings adjoining it. This path was nearly overgrown with grassy and sedgy thickets in which were numerous tall composites. Among the grasses I recognized *Brachyelytrum erectum*, *Panicularia nervata*, *Bromus purgens* and *Hystrix hystrix*; the sedges were species of *Carex*, one of which appeared to be *C. lupulina*. In these grassy areas and the rank herbage bordering it I found on July 27 a considerable number of nymphs of *Melanoplus scudderi*, also smaller numbers of adults of *Melanoplus gracilis*, *Dichromorpha viridis* and *Chlœaltis conspersa*. Near the edge of the wood, in a grassy opening not far from the button-bush bog, I found a single female nymph of *Truxalis brevicornis*. At various points along the edges of the woods and in cut-over areas *Dissosteira carolina*, *Melanoplus atlantis* and *Spharagemon bolli* were of frequent occurrence.

2. This was on the west bank of Burnett Creek in the stream bottoms about two and one-half miles southwest of Battle Ground. The surface is elevated only a few feet above the level of the stream and forms a nearly flat tract between the stream and the neighboring terrace. It is well wooded, the larger trees being chiefly cottonwood (*Populus deltoides*) and buttonwood (*Platanus occidentalis*). The larger trees were much scattered and beneath them the marshy ground supported a rich undergrowth of small trees, shrubs and tall herbage. The principal shrubs were hazel (*Corylus americana*) and Pussy-willows (*Salix discolor*). In the more open bogs the vegetation consisted of a reedy herbaceous growth in which I noted such plants as *Typha latifolia*, *Homalocenchrus oryzoides*, *Cinna arundinacea*, *Panicularia nervata*, *Scirpus atrovirens*, *Ambrosia trifida*, *Sagittaria latifolia*, *Vernonia fasciculata*, *Eupatorium purpureum*, *Eupatorium perfoliatum* and the usual host of sunflower-like composites (species of *Helianthus*, *Bidens* and allies). The soil at this place is mapped by the Bureau of Soils as Wabash fine sandy loam, but in these bogs it was almost a true peat. This place was visited twice, on August 9th and September 13. On the former date thirteen species were taken. Of these the most common in or about the bogs were *Conocephalus brevipennis* and *Melanoplus differentialis*. With them were smaller, but not inconsiderable numbers of *Melanoplus obovatipennis* and *Conocephalus nigropleurum*, while only a few examples of each of the following species were taken in similar haunts: *Melanoplus scudderi*, *Melanoplus gracilis*, *Melanoplus femoratus*, *Scudderia furcata* and *Orchelimum vulgare*. One individual of *Truxalis brevicornis* was observed and captured along the edge of a rather extensive growth of cat-tail (*Typha latifolia*). *Melanoplus femur-rubrum*,

which abounded in the surrounding country, appeared to be scarce in this place as only a single individual was observed. In a relatively dry part of the woods, where the ground was slightly damp, but by no means wet, were observed in a few specimens of *Dichromorpha viridis* and a single male *Spharagemon bolli*, the latter doubtless a stray individual from the dryer groves of the adjoining upland. On September 13 the fauna had much the same character, but was evidently poorer in both individuals and species. Of the latter only nine were recorded and of these only two, *Orchelimum nigripes* and *Chlœaltis conspersa* had not been taken on the earlier date. The former species is usually the most abundant of the bog "long-horned" grasshoppers, but at this place it was exceptionally scarce. Of *Chlœaltis conspersa* only a single male was taken along the edge of the cat-tail bog close to the spot where the *Truxalis* was taken on the earlier date. Besides these other species taken or observed on September 9 were *Dichromorpha viridis* (10), *Melanoplus obovatipennis*, *M. differentialis*, *Scudderia furcata* (10), *Orchelimum vulgare*, *Conocephalus brevipennis* and *C. nigropleurum*.

3. The Purdue Experimental Farm in West Lafayette is located on "second bottom" land. The soil is the Sioux loam. Nearly all the land is under cultivation, the principal crops being corn, wheat, rye, oats, clover, cow-peas, alfalfa and soy beans. Where the land is untilled, as along fences and the borders of paths, there is a firm blue-grass sod in which scattered patches of clover (*T. pratense*) are frequent; also the usual weeds, such as witch-grass (*Panicum capillare*), spreading panic-grass (*P. dichotomiflorum*), crab-grass (*Echinochloa crus-galli*), foxtail (*Chaetochloa viridis*, *C. glauca*), Orchard grass (*Dactylis glomeratus*) and *Eragrostis major* and *purshii*. In the more fully cultivated portion the "home" of the Orthoptera was in this relatively undisturbed grassy sod, although they spread from this in large numbers into the neighboring plats. The most abundant species here was naturally *Melanoplus femur-rubrum*; other common forms were *Encoptolophus sordidus*, *Dissosteira carolina*, *Melanoplus atlantis*, *Orchelimum vulgare* (specially in the taller grasses, such as fox-tail, etc.) and *Conocephalus strictus*, the latter very common in the denser areas of blue grass. Other species of frequent occurrence, but not so abundant as those just mentioned, were *Syrbula admirabilis*, *Chortophaga viridifasciata*, and *Melanoplus differentialis*. Occasionally a specimen of *Schistocerca americana* would be taken or observed in the rank weedy growth bordering the experimental plats and in the more thickly

planted plats themselves. Once two individuals of *Neonocephalus robustus crepitans* were taken and another heard in the corn fields; both of those captured were taken on corn in the early evening.

The most interesting collecting on the Purdue grounds, however, was done in a small waste lot not far from the Lake Erie and Western Railroad. About half of this lot was occupied by a nearly pure growth of timothy (*Phleum pratense*), while the remaining half had at some time or other been used as a dumping place for manure or other refuse and was now occupied by a rich growth of *Elymus virginicus*, with which were intermixed some areas of *Bromus (ciliatus?)* and a few clumps of a taller species of *Elymus*, probably *canadensis*. On one side near a fence row was a rank growth of sumac (species not determined). In another part of the field at one end of the *Elymus* formation in a shallow gully was a rank growth of green foxtail (*Chaetochloa viridis*). Collections were made here at intervals throughout the summer. The species were much the same as those occurring in the cultivated areas, but in addition a number of species were taken which were absent or very rare in the latter. In this waste land most of the collecting was done in the timothy, which had recently been cut, a circumstance which made it relatively easy to capture the grasshoppers. *Melanoplus femur-rubrum*, *Melanoplus atlantis*, *Encoplophus sordidus* and *Conocephalus strictus* were here abundant, while both *Syrbula admirabilis* and *Arphia xanthoptera* were of frequent occurrence. Early in July *Melanoplus femoratus* was fairly common in this tract, but it soon ceased to be an evident component of the fauna. Two ♀♀ of *Orphulella speciosa* were taken on July 22; repeated search failed to reveal any additional specimens of this apparently very rare species. A single male *Scudderia texensis* was also captured here the same date. In the *Elymus* patch a solitary male *Conocephalus fasciatus* was taken also on the same date; while much later in the season—September 13—a small colony of *Conocephalus nemoralis* was found in a place where the *Elymus* was encroached upon by the sumac thickets. *Melanoplus differentialis* was also frequent here. Outside of these three species, the forms in the *Elymus* area were the same as those in the timothy with the exception of *Arphia xanthoptera* which appeared to be limited to the latter. The foxtail growth formed the favorite habitat of *Orchelimum vulgare*. The same grass also yielded a female of *Stenobothrus curtispennis*.

A short distance west of this lot in the adjoining field, which had been

planted in clover, I captured a female specimen of *Schistocerca alutacca*. The capture was made close to the railroad, along which there was a mixed growth of elder (*Sambucus*) and white melilot (*Melilotus alba*). The latter formed a very rank growth in some abandoned gravel pits on the opposite side of the railroad. The color of this specimen was much duller than that of examples from the New Jersey sphagnum bogs, being an olive brown or pale leather color with hardly a trace of green, and with the dorsal stripe, although easily recognizable, by no means conspicuous.

4. At this point some roadside collecting was done. The place is on the slope leading from the "second bottom" at West Lafayette to the upland immediately north of the town. The roadside vegetation consisted in the dryer parts of a mixture of blue grass and timothy and in the gullies of a rank growth of *Milolus alba*. The Orthoptera were all of common types. *Melanoplus femur-rubrum* swarmed everywhere, while its congener, *M. differentialis*, was almost entirely limited to the thickets. In the blue-grass-timothy areas *Conocephalus strictus* was common, while *Syrbula admirabilis* was of frequent occurrence.

5. This place, locally known as "the tank" from the presence of the storage tank of the West Lafayette water company, is on the edge of the upland at the head of a deep ravine known as Happy Hollow. It overlooks the Wabash bottoms, "second bottoms" being absent from this point north. The soil is Miami silt loam. The land was untilled the past season and had evidently not been in cultivation for a long period. It was open, but at its southern edge where it meets the steep slopes leading down into Happy Hollow was bordered by the relatively dense woods which clothe these slopes. The open areas were closely covered with blue grass with which were locally intermixed small areas or scattered clumps of wiregrass, *Poa compressa*, and foxtail, *Chactochloa glauca*. There were also considerable clover and some low trailing briers. Close to the woods the blue grass became rather sparse and grew only in short scattered clumps with open places between where the bare soil was exposed or where certain hardy herbs, mostly composites, grew. In one or two places on the higher land where the blue grass was very thin, were formations of *Andropogon scoparius* with *A. furcatus* as a minor constituent. At one place immediately adjoining the woods was an extensive patch of *Tridens flava*. Within the outer edge of the woods on some level stretches where the less eroded parts of the bluff project out into the ravine, were a few scrubby

areas containing only scattered grasses, but with many low saplings and some herbaceous undergrowth. The woods on the upper portions of the ravine slopes adjoining the upland were of the mixed hardwood type. The dominant tree was the white oak, but with it were many hickories, elms, sugar maples, lindens, red oaks, beeches and dogwoods.

Collections were made in the open fields above the woods, along the borders of the woods and in the woodland scrub areas. In the more open areas, farthest from the woods, wherever the blue grass or its congener, *Poa compressa*, was thick and luxuriant, common species were *Melanoplus femur-rubrum*, *Encoptolophus sordidus* and *Conocephalus strictus*; *Syrbula admirabilis* was of frequent occurrence. Where the grass was shorter and coarser with some interspaces a number of additional species were common such as *Melanoplus allanis*, *Arphia xanthoptera*, *Dissosteira carolina* and *Hippiscus rugosus*. Of *Arphia xanthoptera* and *Hippiscus rugosus* both the yellow-winged and the red-winged types appeared to be about equally frequent. Both of these species were common in the more barren areas along the very edge of the woods, where they were associated with *Spharagemon bolli* and *Melanoplus luridus*, each of which was of frequent occurrence, but did not appear to spread any appreciable distance from the immediate vicinity of the trees. Within the woods in the scrub areas previously referred to the two last-mentioned species were the only ones found. Other species occurring at this locality were *Chortophaga viridifasciata* and *Orchelimum vulgare*, long-winged phase. Nymphs of the former were frequent in some areas of dwarfed blue grass in spring and again in the fall, while a smaller number of the latter were found in a scrub area along the borders of the woods.

6. This includes the outer edge of the Wabash bottoms a short distance south of West Lafayette. The outer edge of the bottoms at this point is marked by a gently sloping bluff which leads up to the second bottoms of West Lafayette. Near the base of the bluff is a road and below the road, between it and the level surface of the present bottom, is a short slope which was partially wooded, the common trees being cottonwoods, honey-locust, hackberry, elm and shingle oak. The woodland here formed a narrow fringe and beyond it, occupying all the level areas, were the usual corn fields of the bottoms. Beneath the trees was a fairly dense undergrowth of shrubs and tall grasses of which species of *Elymus* were most frequent, especially *E. virginicus*. The soil was a mixture of the gravel derived from the material of the bluff itself and alluvium depos-

ited by the river during periods of overflow. Owing to its position, the presence of the rank vegetation and of the resulting humus the soil was in most places moderately damp, but not actually wet. This, however, was not the case in one spot where the ground was perpetually moist on account of the constant seepage from the bluff. The substratum at this spot was a black or dark grey muck with much gravel in its deeper levels. Trees were absent from these wetter areas and they were accordingly occupied by a rank growth of the usual herbaceous swamp plants the more conspicuous of which in this swamp were *Typha latifolia*, *Homalocenchrus oryzoides* and *Ambrosia trifida*. South of the swamp was a small bit of open woodland in which there was a rich undergrowth of grasses. Of these the species of *Elymus*, chiefly *E. virginicus* with some *canadensis*, occupied the better lighted areas while in the more shaded spots such forms as *Homalocenchrus virginicus*, *Muhlenbergia* apparently *M. tenuiflora*, *Korycarpus arundinaceus* and *Hystrix hystrix* were common. Adjoining this woodland on the south was an open pasture in which there was a good stand of *Tridens flava*.

Quite a number of interesting Orthoptera were taken in this locality. In the drier situations the patches of *Elymus canadensis* yielded such species as *Dichromorpha viridis*, *Melanoplus viridipes*, *Melanoplus atlantis*—an unusually humid environment for this form—*Melanoplus femur-rubrum*, *Melanoplus femoratus*, *Amblycorypha rotundifolia* and *Conocephalus nemoralis*. With the exception of *Melanoplus femur-rubrum* none of these were common or widespread, being in most cases represented only by scattered individuals or an occasional colony. Other grasses besides the *Elymus* were searched for Orthoptera, but, excepting *Tridens*, proved to be barren. In the dense thickets of ragweed, *Ambrosia trifida*, surrounding the more boggy spots *Melanoplus femur-rubrum* and *Melanoplus differentialis* were abundant, while in the same tracts a few examples of *Melanoplus scudderi* were also taken. In the swamp the Orthoptera were most numerous in the *Homalocenchrus oryzoides* and the immediately adjoining thickets; they were apparently quite infrequent in the cat-tails. The most abundant swamp species were in order of relative numbers *Melanoplus femur-rubrum*, *Conocephalus bevipennis*, *Melanoplus differentialis*, *Orchelimum nigripes*, *Orchelimum vulgare* and *Conocephalus nigropleurum*; in much smaller numbers were found such species as *Oecanthus fasciatus*, *Oecanthus quadripunctatus*, *Scudderia furcata*, *Neoconocephalus palustris*, *Orchelimum gladiator*, *Conocephalus fasciatus*, *Conocephalus nemoralis*

and *Conocephalus attenuatus*. In the pasture in which *Tridens flava* was a common plant *Syrbula admirabilis* and *Conocephalus strictus* were of frequent occurrence along with larger numbers of the ubiquitous *Melanoplus femur-rubrum*. Close to the border of the same field, where there were some extensive patches of *Elymus virginicus*, several examples of *Dichromorpha viridis* were observed.

7. This was a level tract of very open woodland located on top of the bluff overlooking the bottom lands included in locality 6. The ground here had been used for pasturing cattle and the herbaceous vegetation was accordingly quite short and scanty. Orthoptera were scarce. Each *Melanoplus atlantis* and *Melanoplus femur-rubrum* were frequent, while in one place where there was considerable slope and a fair amount of scrub growth a few examples of *Spharagemon bolli* were seen. Late in June *Atlantius testaceus* occurred in small numbers, several being captured one night on low shrubs and tall weeds.

8. This locality was a small open grove at the top of the highest line of bluffs at the north end of a ravine situated nearly half way between West Lafayette and the mouth of Indian Creek. The soil was the Miami silt loam which in this exposed situation was quite dry and barren and had a decided sandy appearance. Along a recently cut roadside I found at this point a young specimen of the black-jack oak, the presence of which naturally indicates the barren character of the location. The soil here at the time of my visit—August 24th—was formed of blue-grass sod with occasional patches where the ground was bare or but sparsely covered with vegetation. In such places the common woodland Panicum, *P. huachuca*, was frequent. In two or three places erosion had worn slight gulleys from which most of the finer soil particles had been washed away leaving a very hard and stony soil on which very little vegetation had as yet obtained a foothold. These gulleys were the favorite habitats of the more geophilous Orthoptera such as *Arphia xanthoptera*, *Spharagemon bolli* and *Dissosteira carolina*.

Only about a half hour was spent in collecting at this spot, during which examples of the following species were taken or identified: *Syrbula admirabilis*, *Arphia xanthoptera*, both yellow-winged and orange-winged types, *Encoptolophus sordidus*, *Sphragemon bolli*, *Dissosteira carolina*, *Melanoplus femur-rubrum* and *Melanoplus luridus*.

9. This locality is about half a mile southeast of Battle Ground in a region covered by Miami fine sand. Practically the whole country is under

cultivation. The only collecting was done in a limited bit of roadside where the banks were occupied by a mixed growth of two tall bunch-grasses, *Tridens flava* and *Andropogon furcatus*. My visit to this spot was made August 30. At that time the following species were taken or observed, all being fairly frequent: *Melanoplus femur-rubrum*, *Melanoplus atlantis*, *Conocephalus strictus*, *Dissosteira carolina*, *Syrbula admirabilis* and *Arphia xanthoptera*.

10. This locality is on the east side of the Wabash about three miles north of Lafayette and a mile southwest of Wild Cat Creek. At this point there is a well-marked bluff marking the dividing line between the upland, here formed by Sioux sandy loam, and the Wabash bottoms. At the base of the bluff is an extensive marsh, shown on the Bureau of Soils map as a crescent-shaped patch of muck. The upland immediately bordering the bluff is occupied by a cemetery in which there are many large trees, the whole forming an open grove with no undergrowth except the ordinary blue-grass sod. In this cemetery, frequenting the relatively dry blue grass were numerous examples of *Melanoplus scudderi* and *Melanoplus luridus* along with the usual *Melanoplus femur-rubrum* and *Encoptolophus sordidus*. On the steeper slopes, where there was a considerable amount of herbaceous undergrowth and some patches of *Andropogon furcatus*, a few examples of *Spharagemon bolli* were seen and, near the base of the slope, in a shallow depression, where there was a thick growth of a bright green, succulent grass, a small number of *Stauroderus curtippennis* were found. *Dissosteira carolina* was as usual common on paths and driveways both on the upland and in the bottom. The swamp at the base of the bluff was quite open and was of the type usual to the bottoms with rice cut-grass, *Homalocenchrus oryzoides*, forming the dominant vegetation of the wetter areas. On the side toward the bluff this growth was bordered by a thicket formed mostly by tall herbaceous plants among which sunflowers and goldenrods were conspicuous; while on the opposite side toward the open bottom lands it was bordered by a weed vegetation in which a tangled growth of smart-weed (*Polygonum*) predominated. In the rice cut-grass the common Orthoptera were *Melanoplus femur-rubrum*, *Conocephalus brevipennis*, *Orchelimum nigripes*, *Orchelimum vulgare* and *Melanoplus differentialis*. Both of the last-named species and also *Conocephalus nigropleurum* were frequent in the surrounding thickets, while *Orchelimum vulgare* and *Melanoplus femur-rubrum* swarmed in the *Polygonum* areas. In addition to the two forms last mentioned other species tak-

en in the bordering thicket in fair numbers were *Melanoplus obovatipennis* and *Conocephalus nemoralis*. With them were found occasional examples of *Dichromorpha viridis*, and *Melanoplus femoratus*. In the *Homalocenchrus oryzoides* two specimens of *Neoconocephalus patustris* and one each of *Scudderia texensis* (a ♂ apparently this species) and *Orchelimum vulgare* long-winged type were taken. Collecting was done at this place on September 6th.

11. At this place collections were made on July 19 and October 3. The locality was the low alluvial tract along the Wabash at the mouth of Wild Cat Creek. Most of the land is under cultivation, but there is some open woodland on the adjoining bluffs. Along the roadside were the usual weedy tracts inhabited by *Melanoplus atlantis* and *Melanoplus femur-rubrum*, the latter being by far the most abundant. In the ranker herbage and weedy tracts *Melanoplus differentialis* was of frequent occurrence. On the bare paths and in the plowed fields *Dissosteira carolina* was common. The remaining species were few in number and were found only in grassy depressions close to the river. In one of these which contained an almost pure stand of *Elymus virginicus* a few examples of *Stauroderus curtipennis* were observed on July 19; in the same place a single specimen of each of the following was taken: *Melanoplus walshii*, *Orchelimum gladiator*, *Conocephalus fasciatus* and *Conocephalus nigropleurum*. In another depression, examined on October 3, the dominant growth was a species of *Muhlenbergia*; in this *Orchelimum vulgare* and *Conocephalus brevipennis* were common, a single specimen of *Conocephalus nigropleurum* was also taken here.

12. While on an inspection trip on the upland between West Lafayette and Montmorenci on August 12 I made a rapid examination of several small areas in which the ground was more or less damp and covered either with thick succulent blue grass or species of *Carex*. Orthoptera did not appear to be very common in such places, except such ubiquitous forms as *Melanoplus femur-rubrum* and *Orchelimum vulgare*. In one rather wet depression, where there was a nearly pure growth of *Carex*, *Conocephalus fasciatus* was rather common: it also occurred, though in small numbers, in blue-grass depressions. In one of the latter bordering a small grove a small number of *Stauroderus curtipennis* were observed. A male *Scudderia furcata* was taken near here in some thick grass at the side of a small stream.

13. This was a very limited tract on the edge of the bluff overlook-

ing the Wabash valley about three miles southwest of Lafayette. Collecting was done only along the right of way of the Wabash Railroad. This locality was visited only once, and that on October 12, when many species had died out or become very scarce. Only five species were noted, four of which were common in the waste lots adjoining the railroad. They were *Melanoplus femur-rubrum*, *Encop^tolophus sordidus*, *Dissosteira carolina* and *Melanoplus atlantis*, the last-named being the least frequent. The only other species observed on this trip was a male *Schistocerca americana* which was found in a local growth of *Andropogon furcatus*.

14. This was a very interesting undrained depression of considerable size situated in an open field on the upland about two miles northwest of West Lafayette. The substratum in the depression was a dark muck. At the time of my visit, October 13-14, it was quite dry and crisp at the surface, but within a fraction of an inch below was still quite moist and sticky. The centre of the swamp was nearly devoid of vegetation; doubtless in times of normal rainfall it is submerged. Surrounding this is a wide fringe of reedy vegetation formed of cat-tails, *Typha latifolia* and a tall species of rush, which was similar in general aspect to *Juncus effusus*, though owing to the lateness of the season I was unable to certainly identify it. Intermixed with both of these was a luxuriant growth of rice cut-grass, *Homalocenchrus oryzoides*. Surrounding these again was an outer thicket of tall herbaceous plants, such as asters, goldenrods, iron-weeds, sunflowers and their associates.

The Orthoptera of this swamp were unlike any found elsewhere in the extreme abundance of two Tettigoniids, a peculiar color-phase of *Orchelimum nigripes* and *Conocephalus attenuatus*, both of which simply swarmed throughout the *Typha-Homalocenchrus* areas although they largely avoided the rush and were entirely lacking in the herbaceous marginal thicket. The large numbers of *Conocephalus attenuatum* in this place was surprising, for, although it has been known for a long time to be native to the state, I had, previous to my discovery of this marsh, been able to procure only a single example in the region about Lafayette and was accordingly inclined to look upon it as a very rare species in this particular part of the State.

Other species associated with the two species just mentioned in the cut-grass-cat-tail formation were *Conocephalus nigropleurum*, *Conocephalus saltans* and a small *Orchelimum* which Mr. Rehn has assigned to *O. agile*. All of these were quite scarce at the time I examined the place, only a

single example of each of the last two species being seen and only a single pair of *Conocephalus nigropleurum*. Noteworthy was the entire absence in this marsh of the two most frequent marsh "long-horned" grasshoppers of this region, *Conocephalus brevipennis* and the typical phase of *Orchelimum nigripes*.

In the herbaceous thickets forming the marginal vegetation of the marsh Orthoptera were not very common, the only species taken there being *Melanoplus differentialis* and *Melanoplus obovatipennis*, both of which were only moderately common. In the open clover field surrounding the marsh the only species observed was *Melanoplus femur-rubrum*.

15. This was a small lateral ravine which opened into the valley of Indian Creek close to where it empties into the Wabash. It was visited June 28. Orthoptera were very scarce at this time, but on a steep wooded slope where there was much bare ground with scattered growth of the woodland *Panicum*, *P. huachuca*, I captured a male *Melanoplus fasciatus*. The woods here were denser than usual and were cool and shady with only few scattered openings where the direct rays of the sun reached the ground. The soil at this spot is mapped as Miami silt loam.

16. This includes the east bank of the Wabash and the adjoining bottoms about 2½ miles southeast of Battle Ground examined August 30th. The river bank here slopes very gently and at the time of my visit was covered next the river with a growth of sedge, apparently *Scirpus americanus*, and landward of this by *Homalocenchrus oryzoides*. Above this on higher ground was a fringe of woodland with a dense undergrowth of *Muhlenbergia*. Beyond this were the flat cultivated lands of the bottoms. At this point the bottoms are about a quarter of a mile wide. At their outer edge—the edge away from the river—they are characterized by the usual line of high bluffs forming the edge of the neighboring upland. At the base of the bluffs was the usual seepage zone, which at this place was represented by an extensive marsh in which *Homalocenchrus oryzoides* formed the bulk of the vegetation. Bordering it were the accompanying thickets of tall composites.

The Orthoptera of the river bank at this place were disappointingly scarce. The only species at all common was *Orchelimum nigripes* which was observed only in the cut-grass. A single specimen of *Neoconocephalus palustris* was taken in the sedge, but it had apparently strayed there from the cut-grass areas. No other species were noted on the river margins. On the cultivated parts of the flood plain there were in several places rank

growths of common weeds and in these *Melanoplus femur-rubrum* and *Melanoplus differentialis* were abundant. The best collecting from the standpoint of variety was afforded by the marsh at the base of the bluffs. Here in the cut-grass I found considerable numbers of *Paioxya hoosieri*, the only place where I obtained this interesting species. With it were large numbers of *Orchelimum nigripes*, *Conocephalus brevipennis* and *Conocephalus nigropleurum*. In the marginal thickets were observed such forms as *Melanoplus differentialis*, *Conocephalus fasciatus*, *Melanoplus scudderi* and *Melanoplus obovatipennis*.

17. This was a small open groove on rather dry barren soil. It is located on a gentle slope just above the Wabash bottoms on the west side of the river about three miles southwest of West Lafayette. The soil is Sioux sandy loam. In the groove at this point it supports a rather weak growth of blue grass. In the driest parts the blue grass is sparse and in such places *Panicum huachuca* becomes a noticeable constituent of the herbaeous flora. The Orthoptera taken here were the usual species of dry open woodland. In July and early August *Spharagemon bolli* was quite frequent while later in the season *Melanoplus scudderi* and *Melanoplus luridus* were common.

ANNOTATED LIST OF SPECIES.¹

Diaperomera femorata Say. A single specimen, a male, taken in low woods on Burnett Creek near Battle Ground (2), August 9.

Acrydium (Tettix) ornatus Say. Moderately frequent in spring on dry hillsides and in stubble fields on the upland near West Lafayette.

Truxalis brevicornis (Linnaeus). Two specimens; a female nymph taken July 27 in upland deciduous woodland about one mile northwest of West Lafayette (1) in a grassy tract a short distance from a bog dominated by buttonbush, *Cephalanthus occidentalis*; a mature male taken August 9 in low woods along Burnett Creek (2) near Battle Ground at the edge of a bog containing cat-tail (*Typha latifolia*), sedges (species of *Carex*, *Scirpus atrovirens*), *Sagittaria*, *Panicularia nervata*, *Helianthus* spp. Species apparently quite scarce as no other examples were seen.

Syrbula admirabilis (Uhler). Of frequent occurrence in all relatively dry grassy areas at higher levels and locally at least, where conditions are suitable, not uncommon in bottom lands. The species is prevailingly campestrial in its habitat, being especially fond of open grass lands; less fre-

¹The nomenclature used here is that given in an unpublished list of unsynonymized terms compiled by Mr. Morgan Hebard.

quently it may occur along the grassy borders of open woodland. I have taken it in areas occupied by blue grass (*Poa pratense*), timothy (*Phleum pratense*), red-top (*Tridens flava*) and bunch-grass (*Andropogon scoparius* and *furcatus*). I have the following records: July 23, 2 nymphs in dense patch of *Bromus* on Purdue Experimental Grounds (3); July 26, nymphs fairly frequent in timothy stubble on Purdue Experimental grounds (3); August 1, adults, especially males, frequent and nymphs common in grassy roadside patch (timothy-blue grass) on upland slope (4) north of Lafayette; also in dry blue grass and *Poa compressa* in field at "the tank" (5); August 20, several males in open field dominated by *Tridens flava* along outer margin of the Wabash bottoms below West Lafayette (6); August 24, frequently within the borders of dry open woodland on the top of the bluffs at the head of a ravine (8) about half-way between West Lafayette and the mouth of Indian Creek, chiefly in dry blue-grass, and associated with *Arphia xanthoptera*, *Spharagemon bolli*, *Encoptolophus sordidus* and *Melanoplus luridus*; August 30, frequent in roadside and fence-row grasses, especially in a patch of *Tridens flava* and *Andropogon furcatus* about a half mile southeast of Battle Ground (9); September 1, of common occurrence in grassy uplands at head of "Happy Hollow" (5) occurring in blue grass, wire-grass (*Poa compressa*) and *Andropogon furcatus* and *scoparius*; October 4, appears to be getting scarce now on the Station grounds (3); October 26, a single female seen in dense patch of *Poa compressa* in locality 5; October 31, a dead female found on cement sidewalk on Experimental Station grounds (3).

Orphulella speciosa (Scudder). Apparently very rare, only four specimens having been seen or taken throughout the entire season. These occurred in dry, open, grassy tracts on untilled land.

July 22, two females taken in timothy stubble in waste lot on Purdue Experimental Grounds (3); in both of these the tegmina are longer than the abdomen and their tips reach the tips of the hind femora. One female has the discoidal area of the tegmina occupied in part by a double row of cells, a character of its congener *pelidna*; the specimen, however, is unquestionably *speciosa*. August 1, a male taken in *Poa compressa* in a field at "the tank" (5). September 1, a male taken in patch of *Andropogon* on upland at the head of Happy Hollow (5).

Dichromorpha viridis (Scudder). Appears to be only moderately frequent and is largely restricted to damp situations within or along the

edges of open woodlands. It is more frequent in bottom lands than in more elevated tracts.

July 27, frequent in grassy and sedgey spots in humid upland woods (1); August 9, in small numbers in low woods and thickets along Burnett Creek (2); August 20, occasional in a low field along the outer edge of the Wabash bottom near West Lafayette (6) occurring in *Tridens flava* and *Elymus virginicus*; September 6, occasional in the undergrowth on a wooded slope near a *Homalocenchrus oryzoides* marsh (10); September 13, a female taken in low woods along Burnett Creek (2).

Chlaenius conspersa Harris. Occasional in grassy spots in damp woodlands; very local.

July 27, several males and one female observed in humid upland woods northwest of West Lafayette (1), in a grassy clearing where the prevailing herbaceous vegetation consisted of *Carex*, *Elymus* and *Hystrix*; September 13, a single male taken in low woods along Burnett Creek (2), at the edge of a cat-tail bog.

Stenoderus (Stenobothrus) curtipennis (Harris). Apparently only moderately frequent and rather local, occurring in humid tracts well covered with succulent grasses.

July 19, in small numbers in the bottoms near the mouth of Wild Cat Creek (11), in dense growth of *Elymus virginicus*; August 12, a female taken in patch of fox-tail (*Chatochloa viridis*) in a waste lot on the Purdue Experimental Grounds (3); July 12, a small colony in a moist grassy depression along the edges of woodland on the upland between Lafayette and Montmorenci (12); September 6, quite scarce in grassy areas on a wooded slope south of Wild Cat Creek (10).

Arphia sulphurea (Fabricius). Found only once in late April in a sparse growth of blue grass (*Poa pratense*) at the top of a high bluff at Happy Hollow (5). It was at this time in the nymph stage. No others were observed during the season, but it is doubtless more frequent in the spring months than my very meagre field observations made at that season would indicate.

Arphia xanthoptera (Burmeister). Frequent in untilled areas in numerous dry situations, chiefly in upland localities. Both yellow-winged and orange-winged examples occur in nearly equal numbers. The species appears to occur only occasionally on fully cultivated land.

August 1, frequent on the bluffs at the head of Happy Hollow (5), occurring in dry grassy areas and on bare ground on the gentle inclines

adjoining the wooded ravine slopes. Both sexes were represented, also yellow-winged and orange-winged examples in approximately equal numbers; August 24, several examples of both color types observed in open woodland at the head of the ravine between West Lafayette and the mouth of Indian Creek (8), occurring in blue grass areas and in a dry gully; August 28, several of the orange-winged type observed in timothy stubble on a waste lot of the Purdue Experimental Grounds (3); August 30, several observed in an open clover field on dry sandy ground about a half mile east of Battle Ground (9); September 1, both yellow-winged and orange-winged individuals nearly equally common on the bluffs at the head of Happy Hollow (5); October 4, scarce in cultivated ground on Purdue Experimental Grounds (3).

Chortophaga viridifasciata (DeGeer). Only moderately frequent, chiefly in dry upland grass lands. Nymphs were observed in late April, adults from early May to late June and nymphs from early October to the end of November. The species appeared to be most frequent in sparse blue grass areas on the barren slopes at the top of the bluffs.

Encoptolophus sordidus (Burmeister). Abundant in all open dry areas or in quite open woodland.

July 22, nymphs common in timothy stubble in a waste lot on the Purdue Experimental Farm (3); August 1, nymphs common in dry blue grass and *Poa pratensis* areas on the bluffs at the head of Happy Hollow (5); August 19, adult males observed today for the first time on the Purdue University Farm (3) in blue grass areas; August 24, occasional in open woodland on the bluffs at the head of a ravine between West Lafayette and the mouth of Indian Creek (8) in blue grass; September 1, both sexes common in open grassy fields on the bluffs at the head of Happy Hollow (5); September 6, occasional in an open grove on bluffs (10) near Wild Cat Creek; September 13, common in blue grass borders of paths and fences on Purdue Experimental Farm (3); October 4, common on Purdue Experimental Farm (3); October 12, common in waste ground along Wabash Railroad south of Lafayette (13); October 25, occasional on roadside vegetation at the outer edge of the Wabash bottoms near West Lafayette (6); October 26, November 2, small numbers in grassy fields at head of Happy Hollow (5).

Hippiscus rugosus (Scudder). Common in one locality, but not observed elsewhere. It was found August 1 and again on September 1 on the tall bluffs at the head of Happy Hollow (5) where it occurred on untilled

ground in short blue grass and *Poa compressa* areas in dry fields and along the edges of woodlands. It was represented by both yellow-winged and vermilion-winged individuals, the two forms being present in apparently equal frequency.

Spharagmon bolli (Scudder). Frequent in dry open woodland in scrubby and grassy clearings; also along woodland borders, but never in open country.

July 12, a few observed on a hillside covered with open scrub near the borders of woods south of West Lafayette (7); July 23, moderately frequent in an open oak woods south of West Lafayette (17); August 1, frequent along the borders of woods on the bluffs at the head of Happy Hollow (5) in sparse grass and scrub areas; August 9, a male taken in low, humid woods on Burnett Creek (2), probably a stray example from the neighboring upland; August 24, several, in open woodland on the bluffs at the head of the ravine between West Lafayette and the mouth of Indian Creek (8); September 1, several observed in clearings in the woods on top the bluffs at the head of Happy Hollow (5); September 6, few seen in a dry grassy area, largely occupied by *Andropogon furcatus*, on a wooded slope (10) near Wild Cat Creek.

Dissosteira carolina (Linnaeus). Common everywhere on bare ground and in dry grassy areas, where the grass is short, with patches of bare earth intervening. Appeared as adults about July 7 and persisted until the end of October.

Schistocerca americana (Drury). Of sporadic occurrence from late March until at least the middle of October, apparently most frequent in early fall.

March (late), a male taken on a building lot at West Lafayette, in blue grass (3); July 22, a female taken in a field of soybeans on Purdue Experimental Farm (3); September 10, observed a female on Purdue Experimental Farm (3) in blue grass; September 30, a male observed on Purdue Experimental Farm (3); October 4, a male observed on roadside in West Lafayette (3); October 12, a male observed in bunch grass, *Andropogon furcatus*, on bluff along Wabash bottoms south of Lafayette (13).

Schistocerca alutacea (Harris). Evidently very rare and sporadic. I captured a female on August 5 in a field on the Purdue Experimental Farm (3) near the Lake Erie and Western R. R., at a point where there was a fence border growth of elder (*Sambucus*) and melilotus (*M. alba*).

Subsequently, September 24, another specimen, also female, was taken near the same spot by Mr. P. W. Mason.

The specimens were of a much duller tint than those which I have taken in the New Jersey cedar bogs. The latter are typically a bright greenish-olive with a very conspicuous bright yellow mid-dorsal stripe and purplish tegmina. The Lafayette specimens were of a dull olive-brown or leather color with a distinct, but not especially conspicuous, mid-dorsal stripe of pale yellow. The place where the specimens were taken was relatively quite dry.

Melanoplus scudderi (Uhler). Moderately common, at least, locally, in grassy tangles and herbaceous undergrowth in or near woodland.

July 27, nymphs common in grassy clearings in upland woods (1), northwest of West Lafayette; August 9, an adult male taken in low woods in a thicket at the edge of a bog on Burnett Creek (2); August 20, a male taken in tall herbaceous thicket near a bog at the outer margin of the Wabash bottoms near West Lafayette (6); August 30, a male taken in open thicket at the edge of a bog at the base of a bluff on the outer margin of the Wabash bottoms opposite Battle Ground (16); September 6, frequent in blue grass in an open grove on the bluff near Wild Cat Creek (10), associated with *M. luridus*; October 4, several observed in the grassy thickets of roadside adjoining an open patch of woodland (17), south of West Lafayette; October 26, two females observed in grassy fields on the bluff at the head of Happy Hollow (5).

Melanoplus viridipes Scudder. Apparently very local, only a single specimen, a male, having been taken on June 24 in a patch of *Elymus virginicus* in the fringe of trees marking the outer limits of the Wabash bottoms near West Lafayette (6).

Melanoplus obovatipennis (Blatchley). Frequent locally in the herbaceous thickets surrounding marshes or damp spots generally.

August 9, an adult male and female and four nymphs taken in the thickets surrounding a small bog in low woods on Burnett Creek (2); August 30, a male taken at the edge of a marsh characterized by *Homalocenchrus oryzoides*, *Impatiens* and *Ambrosia trifida* at the base of a bluff at the outer edge of the Wabash bottoms opposite Battle Ground (16); September 6, fairly common in herbaceous thickets (goldenrod, sunflowers, etc.) along the edge of a *Homalocenchrus oryzoides* marsh at the base of a wooded bluff near Wild Cat Creek (10); September 13, a pair taken in swamp border thicket. (*Eupatorium purpureum*, *Solidago* spp., etc.) in

low woods on Burnett Creek (2); October 13-14, frequent in herbaceous thickets (asters, goldenrod, ragweed, etc.) surrounding a cat-tail marsh on the upland northwest of West Lafayette (14).

Melanoplus gracilis (Bruner). Apparently moderately frequent locally in moist or slightly humid woodland locations, frequenting grassy and sedgey tangles and herbaceous thickets in the vicinity of bogs.

July 27, males moderately frequent in grassy and sedgey areas and surrounding thickets in humid upland woods northwest of West Lafayette (1); August 9, adults of both sexes found in small numbers in a bog occupied by *Homalocenchrus oryzoides*, *Carex* spp., *Scirpus atrovirens*, *Sagittaria* sp., *Salix* thickets, etc., in low woods on Burnett Creek (2); September 13, a female taken in a bog border thicket in the same locality (2), associated with *M. obovatipennis*.

Melanoplus fasciatus (Walker). Probably quite rare. A single male specimen was taken June 28 in an exceptionally dense bit of woodland near the base of a steep bluff not far from the mouth of Indian Creek (15). The ground where it was taken was quite bare, except for a few scattered plants of *Panicum huachucae* and a few other forms not determined. My determination of this specimen was kindly verified by Prof. Blatchley.

Melanoplus walshii Scudder (*M. Blatchleyi* Scud.). Only a single specimen, a female, was taken July 19 in a dense growth of *Elymus virginicus* on the flood plain of the Wabash near the mouth of Wild Cat Creek (11).

Melanoplus atlantis (Riley). Abundant, though somewhat local, in open grassland in relatively dry situations. Most frequent in upland localities, but it also occurs in small numbers in the bottoms wherever the conditions allow the formation of dry grassland. The species reaches maturity the latter part of June and persists through the summer and well into the fall. The adults appeared to be most abundant about July 20; they apparently decreased in numbers in late summer and early September, but in some places they seemed to increase again in early October. At the latter period a number of copulating pairs were taken and the individuals were found in localized groups, facts which would perhaps indicate the recent maturing of the specimens and the possibility of a second or fall brood of adults. It is conceivable at least that some of the earlier laid eggs might under favorable conditions hatch out in the fall and thus pro-

duce the apparent increase of adults at this time. Mature examples of this species were seen as early as June 16 and as late as October 25.

Melanoplus femur-rubrum (DeGeer). The most abundant grasshopper, swarming everywhere, except in woodland locations and on very dry and barren ground. Its predilections are for relatively humid areas, and it is in consequence especially abundant in the bottom lands, and about ditches and other moist spots. It avoids dense herbaceous thickets and favors open grasslands and clover fields. It reached maturity by the last of July and was found continuously from then until frost. The last record I have is November 2.

Melanoplus luridus (Dodge). Of regular occurrence, though not always common on grassy spots in dry woods or in their immediate vicinity. Usually associated with *Spharagemon bolli*.

August 1, a male taken in blue grass close to the edge of the woods on the bluffs at the head of Happy Hollow (5), nymphs also found here; August 24, several of both sexes found in mixed blue grass and *Panicum huachucae* in open woods on a bluff at the head of the ravine (8) between West Lafayette and the mouth and Indian Creek; September 1, a small number in a clearing in the woods on the bluff at the head of Happy Hollow (5); September 6, frequent in blue grass in a dry open grove on the bluffs near Wild Cat Creek (10), associated here with *Melanoplus scudderi*.

Melanoplus bivittatus (Say). All specimens seen were of the red-legged or *femoratus* type. The species is only moderately frequent and more or less local. It was found in fair numbers about the middle of July on the grounds of the Purdue Experiment Station, but it soon became quite scarce and after the early part of August only occasional individuals were noted and that only in the more or less rank vegetation that flourishes in neglected spots along the stream bottoms.

July 22, moderately frequent in timothy in a waste lot and in the nearby corn and soybean patches on the Purdue Experimental farm (3); August 9, fairly common in thickets in or near low woods on Burnett Creek (2); August 20, occasional in marshes and surrounding thickets at outer margin of Wabash bottoms below West Lafayette (6); September 6, a female observed in tall herbaceous thickets at the base of a bluff near Wild Cat Creek (10).

Melanoplus differentialis (Thomas). Abundant in sheltered situations in all humid situations; less frequent, but not uncommon in upland

localities. The species is especially characteristic of the dense thickets of tall ragweeds, *Ambrosia trifida*, which abound in all moist areas and are especially frequent about the boggy spots at the foot of the bluffs along the outer edge of the stream bottoms. The earliest nymph stages appear to be limited entirely to bogs and the surrounding thickets, but as the grasshoppers increase in size they wander from these haunts into the adjoining fields and uplands. The adults appeared early in August and persisted to about the middle of October.

August 9, adults noted in thickets bordering a bog along the outer edge of the Wabash bottoms near West Lafayette (6); also in thickets of *Melilotus alba* on the upland (3); August 9, common in bog border vegetation in or near low woods on Burnett Creek (2); August 20, common in *Homalocenchrus oryzoides* bog and adjoining thickets of *Ambrosia trifida* and associated plants at the base of the bluff at outer margin of the Wabash bottoms near West Lafayette (6); August 30, abundant in similar situations and corn fields on the Wabash bottoms near Battle Ground (16); September 6, abundant in the same kind of environment at the base of the bluffs near Wild Cat Creek (10); September 13, common in thickets and grassy tangles along Burnett Creek (2); October 13, common in herbaceous thickets surrounding a cat-tail swamp on the upland north-west of West Lafayette (14).

Paroxya hoosieri (Blatchley). Found August 30 in considerable numbers in a bog dominated by *Homalocenchrus oryzoides* and *Impatiens* at the base of a bluff along the outer margin of the Wabash bottoms opposite Battle Ground (16); not found elsewhere.

Scudderia texensis Saussure-Pictet. Apparently scarce as only a few specimens were secured.

July 22, a male taken in timothy patch on waste lot on Purdue Experimental Farm (3); July 26, a female taken, location not recorded; September 6, a female, apparently this species, taken in high thicket at the base of wooded bluff near Wild Cat Creek (10).

Scudderia furcata Brunner. Only a few examples found. August 9, two males taken in tall herbaceous thickets about the edge of a small bog in low woods along Burnett Creek (2); August 12, a male taken in a grassy roadside ditch near a small stream between Lafayette and Mountmorenci (12); August 20, two males and an equal number of females taken in a *Homalocenchrus oryzoides* bog at outer edge of Wabash bot-

toms near West Lafayette (6); September 13, a male taken in a swamp border thicket in low woods on Burnett Creek (2).

Amblycorypha oblongifolia (DeGeer). Only a single specimen, a male, was taken on the night of July 23 in a grove of young silver maples at a nursery two miles southeast of Lafayette. The species is, however, much more frequent than the single capture would indicate since its notes were frequently heard at night throughout midsummer.

Amblycorypha rotundifolia (Scudder). A female specimen was taken July 12 in a patch of *Elymus virginicus* in a narrow fringe of woodland at the base of the bluff on the outer edge of the Wabash bottoms below West Lafayette (6).

Microcentrum laurifolium (Linnaeus). This, or the related species, *retinerve*, appears to be common in trees at Lafayette since its notes were heard continuously throughout late July and August. Only one specimen was actually taken and identified as belonging to *laurifolium*.* It flew into the office at the Experiment Station.

Neoconocephalus robustus crepitans (Scudder). Late in August three males were heard stridulating in the corn plats on the Purdue University Farm (3) and on the evening of August 26 two of these were captured, one being taken in some crab grass (*Syntherisma sanguinalis*), the other on a corn tassel. According to Blatchley this species has hitherto been noted in Indiana only in Laporte County along the shore of Lake Michigan.

Neoconocephalus palustris (Blatchley). Of regular occurrence in open *Homalocenchrus oryzoides* bogs, but not especially frequent.

August 20, one male, two females, in *Homalocenchrus oryzoides* bog at base of bluff on Wabash bottoms near West Lafayette (6); August 30, a female taken in mixed *Scirpus americanus* and *Homalocenchrus oryzoides* on low banks of the Wabash River opposite Battle Ground (16); September 6, one specimen of each sex taken in *Homalocenchrus oryzoides* bog at base of bluff near Wild Cat Creek (10).

Orchelimum vulgare Harris. Abundant everywhere in tender succulent grasses; uncommon in woodland situations.

July 22, males and nymphs abundant in growth of *Chatochloa viridis* on a waste lot of the Purdue Experiment Farm (3); August 9, one male and a female taken in woodland bog in low woods on Burnett Creek (2), far from common here; August 20, common in *Homalocenchrus oryzoides* at base of bluffs along margin of the Wabash bottoms near West Lafayette

*Based on description in Blatchley, Orthoptera of Indiana.

(6); August 28, common in tall roadside vegetation, about ditches, etc., at West Lafayette; September 4, abundant in mixed growth of *Muhlenbergia* sp. and *Chatochloa viridis* in a neglected field at West Lafayette; September 6, common in *Homalocenchrus oryzoides* bog and in adjoining weed areas at base of the bluffs near Wild Cat Creek (10); September 13, small numbers observed in swamp border thickets in low woods on Burnett Creek (2); October 13, rather scarce at West Lafayette (3).

Orchelimum vulgare, long-winged phase. This is the form which has commonly been called *glaberrimum* by Blatchley and the majority of recent writers. Rehn and Hebard, however, have recently reached the conclusion that this term correctly applies to the entirely different red-faced *Orchelimum* of the Middle and South Atlantic States which Davis has called *erythrocephalum* and which I have so designated in my paper on New Jersey Orthoptera. In the last-mentioned work the form termed *glaberrimum* has since been recognized to be a distinct species which Rehn and Hebard are about to describe. Occurs in the same situations as the preceding species, but is much less frequent though by no means uncommon.

July 22, a male taken in patch of *Chatochloa viridis* in a waste lot on the Purdue Experimental Farm (3); August 20, a male and female taken in a *Homalocenchrus oryzoides* bog at the base of the bluffs on the edge of the Wabash bottoms near West Lafayette (6); August 24, a male taken in corn plat at Purdue Experimental Farm (3); August 28, several males observed at night while stridulating on young trees and tall herbs on the bluff at the head of Happy Hollow (5); September 4, one female taken in thick grass on a neglected lot at West Lafayette; September 6, a male taken in *Homalocenchrus oryzoides* at base of bluff near Wild Cat Creek (10); October 4, several seen on Purdue University Farm (3).

Orchelimum gladiator (Bruner). Only two specimens, both males, taken during the season. Both were found in bottom lands in thick grass.

July 12, a male taken in a *Homalocenchrus oryzoides* bog at the base of the bluff on the outer edge of the Wabash bottoms below West Lafayette (6); July 19, a male taken in a thick growth of *Elymus virginicus* on the east bank of the Wabash at the mouth of Wild Cat (11). Professor Blatchley kindly verified my determination of these specimens.

Orchelimum agile (DeGeer). A single individual (female) was taken October 14 in the cat-tail marsh on the upland northwest of West Lafay-

ette. It was in the company of large numbers of *O. nigripes*. Professor Blatchley, to whom the specimen was submitted, assigned it to his *O. campestre*. Mr. Rehn, to whom the same specimen was also sent and who with Mr. Hebard has recently revised the entire genus, informs me it is *O. agile*.

Orchelimum nigripes Scudder. An abundant and characteristic species of open grassy bogs and damp situations generally, being especially abundant in rice cut-grass, *Homalocenchrus oryzoides*.

August 20, moderately frequent in a *Homalocenchrus oryzoides* bog at the base of a bluff along the Wabash bottoms near West Lafayette (6); August 30, abundant in wet places covered with *Homalocenchrus oryzoides* on river bank and bottoms on the east side of the Wabash opposite Battle Ground; September 6, common in *Homalocenchrus oryzoides* in a marsh at the foot of the bluff near Wild Cat Creek (10); September 13, a few specimens observed in a sedgey bog in low woods along Burnett Creek (2); October 3, a few observed in a humid depression covered with *Muhlenbergia* near mouth of Wild Cat Creek (11).

Orchelimum nigripes Scudder (variety). On October 13 and 14 I found a form of this genus in the cat-tail marsh on the upland northwest of Lafayette which I was unable to determine, but which Mr. Rehn to whom I submitted specimens informs me is a race of *O. nigripes* from the typical form of which it differs in the absence of black from the tibiæ and, so far as my Lafayette material is concerned, in its somewhat greater size. On the dates mentioned it literally swarmed in the mixed cat-tail and rice cut-grass areas of the marsh, but was entirely lacking in the marginal thickets.

Conocephalus (Xiphidium) fasciatus (DeGeer). Local and, as a rule, not very common; found typically in open wet or damp locations thickly covered with succulent grasses and sedges.

July 19, a male taken in a thick growth of *Elymus virginicus* on the east bank of the Wabash near mouth of Wild Cat Creek (11); July 22, a male taken in a patch of *Elymus virginicus* on a waste lot on the Purdue Experimental Farm (3); August 12, both sexes moderately common in roadside gulleys and in wet depressions covered with low sedges (*Carex* spp.) on the upland between Lafayette and Montmorenci (12); August 20, a female taken on *Homalocenchrus oryzoides* in a bog at the foot of the bluffs along the margin of the Wabash bottoms below West Lafayette (6); August 30, several examples observed along the margin of a *Homalocen-*

chrus oryzoides bog at the base of the bluffs on the Wabash opposite Battle Ground (16).

Conocephalus (Xiphidium) brevipennis (Scudder) (incl. *ensiferum* Scudder). Abundant in the bottom land marshes, in both open and wooded situations.

August 9, adult males and nymphs frequent in grassy and sedgey swamps in low woods along Burnett Creek (2); August 20, abundant in open *Homalocenchrus oryzoides* bog at foot of the bluffs along edge of Wabash bottoms near West Lafayette (6); August 24, common in a similar type of bog in a ravine near the mouth of Indian Creek (15); August 30, common along the margin of a *Homalocenchrus oryzoides* bog at the base of bluffs on the edge of the Wabash bottoms opposite Battle Ground; September 6, common in a *Homalocenchrus oryzoides* bog at the foot of the bluffs near Wild Cat Creek (10); September 13, frequent in bogs and bog border thickets in low woods along Burnett Creek (2); October 3, common in swamps and humid situations generally on the bottoms near the mouth of the Wild Cat (11), occurring in thick growths of *Muhlenbergia* and *Elymus*.

Conocephalus (Xiphidium) nemoralis (Scudder). Locally present in moderate frequency, occurring in grassy and herbaceous tangles usually in the vicinity of woodlands.

Aug. 20, a male taken in dense growth of *Homalocenchrus oryzoides* in a bog at the foot of the bluffs along the margin of the Wabash bottoms near West Lafayette (6); also nine other individuals composed of both sexes taken in a small patch of *Elymus virginicus* along the border of an adjoining woods; September 6, moderately frequent in thicket undergrowth at the foot of a wooded bluff near Wild Cat Creek (10); September 13, several observed in a thick growth of sumac and *Elymus virginicus* on a waste lot on Purdue University Farm (3).

Conocephalus (Xiphidium) nigropleurum (Bruner). Frequent in herbaceous thickets, especially those forming the margins of bogs dominated by *Homalocenchrus oryzoides*, in both open and woodland situations and usually associated with *Orchelimum nigripes*.

July 19, an immature male taken in a thick growth of *Elymus virginicus* on the east bank of the Wabash at mouth of Wild Cat Creek (11); also an adult male at the edge of a *Homalocenchrus oryzoides* bog at the foot of the bluffs along the east side of the Wabash about half way between localities 11 and 16; August 9, both sexes moderately frequent in the herb-

aceous thickets bordering a small bog in low woods along Burnett Creek (2); August 20, in small numbers in a *Homalocenchrus oryzoides* bog along the outer edge of the Wabash bottoms below West Lafayette (6); August 30, frequent in a *Homalocenchrus oryzoides* marsh at the base of the bluffs on the east side of the Wabash opposite Battle Ground (16); September 6, several observed in border thickets surrounding a *Homalocenchrus oryzoides* bog at base of the bluff near Wild Cat Creek (10); September 13, frequent in border thickets (joe-pye weed, boneset, etc.) surrounding a grassy bog in low woods on Burnett Creek (2); October 2, a male taken in *Muhlenbergia* patch on east bank of Wabash near mouth of Wild Cat (11); October 14, a male and a female taken along the edge of a mixed *Typha latifolia* and *Homalocenchrus oryzoides* marsh on the upland northwest of West Lafayette (14).

Conocephalus (Xiphidium) strictus (Scudder). Abundant in dry, open grass land; most frequent at higher elevations, but occasionally in suitable locations in the bottoms. A common associate of *Syrbula admirabilis*.

July 22, nymphs abundant in timothy and *Elymus virginicus* in waste lot on the Purdue Experimental Farm (3); July 31, adults and nymphs common in tangles of blue grass and bindweed along a fence on the Purdue Farm (3); August 1, nymphs and adults common in blue grass on roadside north of West Lafayette (4); August 5, common in thick blue grass and foxtail areas on Purdue Farm (3); August 20, frequent in an open pasture, dominated by *Tridens flava*, on the outer edge of the Wabash bottoms near West Lafayette (6); August 30, common in a roadside growth of *Tridens flava* and *Andropogon furcatus* near Battle Ground (9); September 1, frequent in grassy fields on bluffs at the head of Happy Hollow (5); October 3, now scarce on Purdue University Farm.

Conocephalus (Xiphidium) saltans (Scudder). Only a single specimen, a male, taken October 14 in a cat-tail cut-grass marsh on the upland (14) northwest of West Lafayette. It was associated with *Xiphidium attenuatum*.

Conocephalus (Xiphidium) attenuatus Scudder. Abundant October 13 and 14 in the upland marsh (14) just referred to, where it swarmed in the mixed cat-tail and *Homalocenchrus oryzoides* formation, but appeared to be entirely lacking in the surrounding herbaceous thickets of asters, goldenrods and associated plants. Only a single specimen was seen elsewhere, a female taken August 20 in a rice cut-grass bog at the foot

of the bluffs on the outer edge of the Wabash bottoms below West Lafayette (6).

Atlanticus testaceus (Scudder) [*pachymerus* Burm.]. Moderately frequent locally in open undergrowth of dry upland woods.

June 17, a male taken by J. J. Davis in a scrub area on the bluff at the head of Happy Hollow (5); June 26, three males taken on tall weeds in open woods on "second bottom" south of West Lafayette (7).

Oecanthus fasciatus Fitch. August 20, several taken in grassy bog at base of the bluffs bordering the Wabash bottoms near West Lafayette (6).

Oecanthus quadripunctatus Beutenmüller. Several taken in same locality and on the same date as the preceding.

Nemobius fasciatus DeGeer. Abundant in grassy places in both dry and moist areas.

Nemobius carolinus Scudder. A female collected October 9 by Mr. P. W. Mason on a road near West Lafayette was identified as this species by Mr. Morgan Hebard.

SOME INSECTS OF THE BETWEEN TIDES ZONE.

CHARLES H. ARNDT.

All insects, except a few species which live entirely in the water and have functional gills, are air breathing animals, breathing by means of tracheæ. Thus we may desire to know how air breathing insects living in a zone which is submerged twice a day, prevent themselves from being drowned; or, if breathing by means of gills, they protect themselves from superoxygenation during the low tide. We may further be interested in any adaptations, or unique instincts, which make the inhabitants of such a locality especially adapted to their environment.

The following observations on the habits of the insects of the between tides zone, were made in the region directly north of Jones' bath house at the head of Cold Spring Harbor Bay; which is included in the lines drawn from 600 E., 200 S., to 200 N., 400 E; and 200 N., 800 E., on map of Inner Harbor made by Johnson and York. Many observations were made on the extreme outer limit of the *Spartina cynosuroides* 275 N., 400 E. The slope of the shore to the west of the boat landing from the outer limit of the *Spartina* to within two feet of the inner limits of the *Spartina* is about 6%. From the latter point to the inner limit of the *Spartina* the slope is more abrupt. The *Spartina* is here replaced by a short (about six inches high), densely matted grass, *Juncus Gerardi*. This covers the entire region around the bath house with the exception of a few pebble-covered areas on which there is a sparse growth of *Spergularia Marina*. (Map 2.)

The highest tides of the summer, July 8, 9.2 feet; August 3, 8.8 feet; submerged the region as far as the bath house. From July 11 to July 29 the *Spartina* area was never entirely submerged, due to the low tides and the absence of any strong easterly winds. The observations extended from July 1, to August 5, 1913.

INSECTS OF THE SPARTINA REGION.

MEGAMELUS MARGINATUS.

This yellowish brown, 2.5 cm. long, leaf-hopper is a common inhabitant of the salt marshes from Connecticut to Florida (Van Duzee). This insect I found only in the *Spartina* area, never more than a few feet from the inner limit. Due to the alertness of this leaf-hopper on sunny days and its inconspicuousness, many hours of observation failed to reveal its whereabouts during high tide, until one rainy morning, while watching their actions from a boat, I found them resting head downwards on the inner part of the shallow "U's" formed by the grass blades. They remained in this position as they were slowly covered by the incoming tide. Their position on the blade is especially advantageous as it encloses large bubbles of air under their wings which serve the double purpose of supplying them with air and of making them inconspicuous by giving them a silvery appearance which makes them resemble closely the stem of the *Spartina* on which they rest, which also has a silvery appearance due to bubbles of air on its surface. On a cloudy day they cling so tenaciously to the blades of the *Spartina* that the blades may be cut off and placed in a jar of water. On July 18, 9 a. m., I put eight of them under sea water by this method. I kept them submerged for twenty-seven hours. On lowering the water in the jar they were still able to fly. I kept them submerged for two days on several occasions, with apparently no ill effect.

Besides this reaction to the tides, which prevents them from being washed away, and their inconspicuousness, which makes them invisible to their natural enemies, the inhabitants of the sea, they have a spur on the third pair of legs which is peculiarly modified to secure their survival in a region at times covered with water. To secure a larger contact area with water to allow them to remain at rest on its surface as well as to hop upon it, this spur has been developed until it is as long as the proximal tarsal segment. The prominent hoods at the distal end of the tibia and also on each tarsal segment, are other modifications for the same purpose. They never walk on the surface of the water, but can hop on it with great ease. The two hooks on the terminal segment of the leg enable it to secure a firm hold on the *Spartina* blade at the time of submergence.

This leaf-hopper was never found further inward than the inner limit of the *Spartina* area. It can be readily distinguished from the leaf-hoppers inhabiting the higher regions of the salt marsh by the prominent

hoods on its legs, by the greatly developed spur, and by the great length of the proximal tarsal segment in comparison to the tibia. In the other species the tibia is at least four times the length of the proximal tarsal segment, while in this species it is only twice as long.

See Plate I, figures 1, 2, and 3. 1, *Megamelus*; 2, Leaf-hopper from the *Juncus* area; 3, Leaf-hopper from the area never covered by ordinary tides. The species whose hind leg is figured in 2 retreated before the tide, but on no occasion during the summer was the region it inhabited completely submerged. These figures show a peculiar development of structures of advantage in each particular environment. No. 3 has no hoods, No. 2 has them somewhat developed, in No. 1 we find the greatest development. If, as I have suggested, these hoods have been developed to aid in hopping on the surface of water, No. 3 would have no use for them and they would necessarily be useless structures.

The chief enemy of the *Megamelus marginatus*, is the only other permanent resident of the *Spartina* area, a small spider, *Grammonata trivittata*. Its principal source of food is this leaf-hopper.

The hoods on its feet, the greatly developed proximal tarsal segment, and the spur, are the peculiar modifications which determined that this leaf-hopper should inhabit this particular region. But why it is only found in the *Spartina* region, is not as easily answered. It may have been the severer competition in the other regions of the marsh, or perhaps the *Spartina* grass is its favorite food, the one on which it is especially adapted to live; or, again, the habits necessary for its continued existence in the tidal zone may make it the easy prey of its natural enemies living in the other areas.

I could not compare the resistivity of this leaf-hopper to drowning with that of those farther back on the marsh, because I could not get any of them to remain under water without placing them in vials covered with cheesecloth. Two to four hours submergence usually killed them. Even the leaf-hopper of the *Spartina* could only survive for several hours when submerged in this manner. This may suggest that the leaf-hopper may secure its air supply from the *Spartina* by piercing the blade to the air channels. I have no experimental evidence to prove this.

GRAMMONATA TRIVITTATA.

This spider inhabits the salt marshes from Long Island to Maine (Emerton). The females are about 3.5 mm. long. Their color is a dark

reddish brown. These spiders are very abundant in the *Spartina* region. A short search will reveal a number of them running up and down the *Spartina* blades or resting head downward on them.

As the tide comes in, they retreat up the *Spartina* grass. When the tide has once chased them out on the isolated blades of *Spartina* (6 to 8 inches from the tip), they retreat to within about two inches of the tip where they remain head downward until the water almost touches them. Then they begin to run wildly up and down the blade, from tip to water, from water to tip, as if they were very much afraid of the water. After doing this a number of times they will calmly walk down the blade under the surface of the water until they come to the pit formed by the union of the stem and the blade. Here they remain until the tide retreats. The pit furnishes them protection from aquatic enemies and supplies them with air, for there is always a considerable amount of air left in the pit. That this reaction is due to an effort to secure protection and air, was shown in several experiments. When they were placed on blades of grass weighted to the bottom of the jar which was slowly filled with water, they went down the blades and attempted to crawl under objects at the bottom of the jar. However, to get under some object did not satisfy them, they kept on moving until they came to a bubble of air. Their actions were especially interesting when safety pins were used to weight down the blades. They would walk entirely around the wire part of the pin until they came to the sheath where there was a quantity of air. Into this they crawled. They always cling tightly to the wire, never attempting to leave it. It is however not essential for them to find a bubble of air. I kept a number of them submerged for three days without any noticeable bubbles being present. When they go below the surface, they always entangle numerous small air bubbles in their short, dense hairs, which are curved backward seemingly for this special purpose. I performed many experiments to compare the resistance of this spider to drowning with that of other species by submerging them in water. In my experiments I used practically all species found farther back on the marsh; *Tetragnatha*, *Lycosa*, *Epeira*, *Attidæ*, *Themsidæ*. This was the only species, with the exception of the young *Clubonia*, which could be submerged in water when resting on a grass blade; i. e., the other species did not hold fast, but tried to escape on the surface of the water. The most striking difference between these spiders when they were placed in small vials whose mouths were covered with cheesecloth and then submerged in a large glass jar of sea

water, was the difference in the size of the air bubbles clinging to the different species. No prominent bubbles were noticed on the Grammonata, while the other species entangled large bubbles in their comparatively long hair which made them almost entirely helpless. Regardless of the disparity in the size of the air bubbles, the Grammonata could withstand a much longer submergence without causing death. The limit for the hardiest of the other species was twelve hours. When placed in vials in a similar manner and the air bubbles withdrawn by means of a pipette, the *Lycosa communis*, the hardiest of all other species, did not survive more than three hours submergence, while the Grammonata could remain for twenty-four hours in the same jar with no ill effect. This would seem to indicate that this species must have some special modification, as perhaps greatly developed air sacs, "lungs", to enable them to resist drowning so successfully. None of the other marsh spiders, with the exception of the crab spiders, are as free from long flimsy hair, which greatly impedes an animal's movements in water, as the Grammonata. Even the *Lycosa Communis*, whose habitat borders on the between tides zone, has comparatively few long hairs, and these are comparatively stiff.

The food of this spider, Grammonata, consists chiefly of leaf-hoppers, but it also captures the small flies which frequent its habitat during the low tide. It is always attached to the tip of the grass blade by a thread which enables it to run on the surface of the water after food without being washed away.

This spider, as the leaf-hopper, *Megamelus marginatus*, is found only in the *Spartina* area. Its great resistance to drowning, its unique covering of hair, and the instinct which causes it to seek safety by clinging to the object on which it rests, instead of seeking safety in flight as other species, all have determined that it could survive in the *Spartina* area; but as to the reason it is only found there, I cannot explain.

CLUBONIA Sp.

This was an immature spider, light tan in color, largest specimen 7 mm. The first specimen was found August 1 after a period of very low tides, at which time I found them in practically the entire *Spartina* area (rare in other areas). The frequent use of a sweep net as well as searching on the ground, failed to reveal an adult spider of this species. The probable solution is that they had hatched, during the low tide period, from eggs laid the previous year. I was very unfortunate in that they

first appeared during my last week at the laboratory, so that my observations did not extend over a period long enough to determine if they were permanent inhabitants of the area, or whether they had just migrated there as several species of Tetragnatha had. This last supposition I consider very improbable, as I found no Clubonias in any other part of the marsh.

Their reactions to the tides were such as would prevent them from drowning or being washed away, but they afforded little protection from their aquatic enemies save that secured from their inconspicuousness. When the tide came in, they crawled to the under side of the *Spartina* blades. When resting in this position with their long legs stretched out along the edges of the blade, they are difficult to see. That they are not destroyed by the tide was shown by the fact that on August 5, after the *Spartina* area had been submerged by the tides of three successive days, I still found them in abundance. I placed several of them, in the position I found them on the grass when covered by the tide, in a jar of sea water and kept them submerged for thirty hours. They were apparently unaffected, showing that this spider, as well as the *Grammonata trivittata*, must have some unique modification to prevent drowning. This spider, however, entangled more air bubbles in its longer, flimsier hair. Its feet are provided with toothed claws, perhaps developed to secure a firm hold on the grass blade.

I am anxious to secure more data concerning this spider. Is it a permanent resident of the *Spartina* area? What accounts for its sudden appearance? Where were the adults, or when? Its resistance to drowning is great enough to allow it to be a permanent inhabitant, but I doubt if it is sufficiently well protected from the inhabitants of the sea. However, the pads on its feet, which resemble those of a water strider and the one large toothed claw as well as several smaller ones on each foot may be structures of particular value to a resident of a between tides zone. The pads would enable it to move on the surface of the water, while the claws would enable it to secure such a firm hold on the grass blade that it could not be washed away.

INSECTS OF THE OUTER JUNCUS AREA.

BEMBIDIUM CONSTRUCTUM.

Of all the insects whose habits I observed, this brownish-black, 5 mm. long, beetle had the widest range. It was the only one to live in both

places that are daily covered by the tides and in places that are not. Its outer limit is 14 N., 4 E. (map 2). They are very abundant in the gravel which covers the wooden platform and also in the region of the tidal drift, especially under the newly washed up sea-lettuce (*Ulva*). In regions daily covered by the tide it is rarely found except in gravel covered areas. By crawling under the gravel it prevents itself from being washed away and protects itself from aquatic enemies. Its actions may be readily observed by hunting for them as the tide rises and the digging a channel around the place where they are found. As the water rises, they will run from place to place, darting under one stone only to leave it for another. They may approach the channel, but will rarely attempt to cross it. Before the tide has completely submerged the place, the bembida will crawl under the gravel to remain there until the tide retreats. When the place is covered by the tide, they may be found at a depth of from three or four inches. The water, when it covers them, encloses a large bubble of air under their wings.

Their action may be easily determined under artificial conditions by placing them in jars containing moist sand and gravel and slowly filling them with water. Several times, when filling the jars, I poured the water in so rapidly that some were swept to the surface. They would then swim wildly on the surface; but when a grass stem was thrust into the sand so as to extend above the surface of the water, they would swim to it and crawl down into the sand. The bubble of air under their wings is so large that should they lose their footing, they will immediately rise to the surface. Their ability to withstand submergence is wonderful. At the end of one day the bubble of air will have almost disappeared and they will have become stupid. By the above gravel jar I kept eight out of ten specimens alive for three days in sea water. Several recovered after five days' submergence, showing a great resistivity to drowning.

This beetle, as well as the *Heterocerous undatus* and *Salda* sp(?) lives on the carrion of the small animals so abundant in the between tides zone.

HETERO CERUS UN DATUS.

This beetle is very rare in this area. Several were found between 4 and 6 N., to 12 and 16 E. (Map 2.) They live in burrows in the ground, only venturing out for food at low tide. Their resistance to drowning is equal to that of the *Bembidium*.

SALDA Sp(?).

This shore bug is very active and very difficult to catch and hard to find except on sunny days, when their shiny black wings make them rather conspicuous. They are found chiefly in the area covered by the *Spergularia Marina*. They live in burrows in the ground, and only venture forth in search of food on sunny days at low tide.

Their resistance to drowning is not nearly as great as that of the beetles; twenty-four hours submergence usually being sufficient to kill them. Twelve hours had very little effect. Like the beetles, they enclose large bubbles of air under their wings. On several occasions I placed several of them on the surface of the water when the area was covered by the tide. They swam on the surface until they came to floating fucus thallus (other floating materials being rejected). They would then crawl to one of the deepest submerged branches, where they would remain. They were almost invisible on the thallus, due to the resemblance of the fucus to their wings with the enclosed air bubble. This, undoubtedly, is a protective instinct.

BDELIDÆ Sp(?).

This reddish-brown plant louse (2 to 3 mm. long) is widely distributed throughout the salt marsh. Their outer limit borders on the inner limit of the prevailing tides. They are never found beyond the inner *Spartina* limit, 6½-foot level. Their mouth parts are especially adapted to sucking plant juices and they live on decaying plants. They are especially abundant on fresh drift weed.

As stated, they are seldom found in the areas covered by the prevailing tides, yet on occasions of a sudden rise in tide levels, they may be found walking on the ground when it is covered by the tide. They do not seem to have any objections to such unusual conditions as they apparently make no effort to escape. They are easily washed away by the tide when once lifted from their feet, as their long slender legs are to their disadvantage. After the retreat of such an unusually high tide, they are not as numerous as they were previous to it, due perhaps to the fact that they have no protection from aquatic enemies, and also because many of them are washed away. There is little danger of drowning. One morning, at 6:45, I placed six of them in vials and submerged them in sea water. At 5 p.m. they were all lying on their backs, apparently dead, but on exposure to the air they all revived. I also placed several in an un-

covered glass cylinder and placed it where it would be covered by the tide. As the tide filled the jar they made no effort to change their positions. Those on the grass blades remained in the same position, as did also those on the ground. The Bdeliidae were still alive in the cylinder after it had been submerged by two tides; i. e., two periods of submergence of five hours each, one period of terrestrial conditions of four hours.

LYCOSA COMMUNIS.

Although the inner limit of the tidal drift, with its myriads of flies should furnish abundant food for spiders, this is the only species prevalent in the drift covered areas. This is a greyish spider from 4 mm. to 5 mm. in length. They venture out beyond the high tide limit, but will always retreat before the incoming tide. Their long, strong legs, which enable them to run rapidly, make them especially adapted to a region where safety lies in retreat. I often found them running inward on the *Juncus* when the ground was already covered by the tide. On several I found them isolated on the blades of *Spartina* grass. When this position was no longer conducive to dryness, they would run rapidly inward over the surface of the water. This is the only insect without wings, frequenting the between tides zone, which retreats before the tide. The *Lycosæ*. are not only the most rapid runners among the spiders, but can also withstand several hours more submergence than the other spiders of the salt marsh, *Grammonata* and *Clubonia*, of course, excepted.

Many winged insects, beetles, flies, etc., are found on the *Spartina* during the low tide; but, although I made no especial study of them, they seem to be only temporary residents. They are never abundant, or even present, immediately upon the retreat of the tide. I never found any submerged on the grass when covered by the tide. They are all good fliers and most undoubtedly retreat before the tide.

On the morning of August 5, when we had the first high tide which covered the outer *Spartina* area entirely for several weeks, I noticed many *Tetragnatha* spiders retreating up the *Spartina* blades because of the rising tide. As the water chased them to the tip of the blade they spun out a long thread which the north wind carried to the higher areas. They ran inward on the thread. When they accidentally became wetted by the water they became helpless. Should the wind have come from the south, they would have been destroyed. No *Tetragnatha* were found in areas that had once

been submerged by the tide. They had migrated to the outer *Spartina* areas during the low tide period. They are very abundant in the higher marsh areas.

It would be extremely interesting to know something of the life histories of the insects of the between tides zone, especially as to where they spend the winter, as to the methods of egg laying, and as to the types of larvæ.

CONCLUSION.

In almost every insect of the between tides zone there appears to be some peculiar protective feature; as, unique instincts, especially adapted external parts, or a greater resistivity to drowning.

Unique instincts, to prevent themselves from being washed away by the tides, are well shown in the tenacious clinging to the blades of *Spartina* by the *Grammonata trivittata*, *Megamelus marginatus*, and *Clubonia* sp (?). Another feature of the same type is the crawling of the *Bembidium constructum* under the gravel. That the environment has undoubtedly led to the formation of these instincts is illustrated by the comparison of the habits, when submerged, of the spiders of the *Spartina* area with those found higher on the marsh. Another instinct which serves the same purpose is the venturing forth for food by the *Salda* only on sunny days at low tide. The *Grammonata trivittata* and *Megamelus marginatus* undoubtedly rest head downward on the *Spartina* grass to prevent themselves from being caught unawares by the rising tide.

Inconspicuousness as a means of protection from aquatic enemies is shown by the swimming of the *Salda* to the *Fucus* thallus when disturbed during high tide. Other reactions serving the same purpose, are illustrated by the resemblance of *Megamelus* to the *Spartina* blade, the crawling the *Grammonata* into the pits at the junction of blade and stem, and the living of the beetles in burrows.


That one of the factors which determines the surviving species in a between tides zone is an ability to resist drowning, is shown by a comparison of the resistivity of *Grammonata* and *Clubonia* with that of other spiders.

A modification of the external features as an adaptation, is shown in the greatly modified legs of the *Megamelus marginatus* and in the short, stiff hair of the *Grammonata trivittata*. The legs of *Lycosa communis* are not essentially different from those of other *Lycosæ*; yet their long,


strong legs, which are conducive to swift movements and which give them the designation of running spiders, make them especially fitted to inhabit a zone in which safety may lie in retreat.

The terrestrial conditions which prevail for one-half of the time in the between the tides zone, make it necessary that the permanent inhabitants of such a zone be insects which breathe not by means of gills, but that they be air breathing insects. Terrestrial conditions are the normal ones for their activities. They are inactive during the high tide, or the period of submergence. The most striking phenomena is the strictly zonal distribution of the insects of the between the tides zone. Each species is undoubtedly specialized in some way to fit it to inhabit this area, and this specialization has rendered it incapable of surviving in other zones. Even the *Megamelus marginatus*, which travels with comparative rapidity, never migrates to the *Juncus* areas. To determine the exact reasons why the insects of this zone cannot survive in the higher areas of the salt marsh, more extensive studies of their life histories and habits must be made. Further observations concerning these features, I believe, will add much to the already extensive and fascinating data relating to adaptation among insects.

*Purdue University,
Lafayette, Ind.*



MAP I.



Map II. Region between 650-750 E, and 20 S-123 N. Scale 1:173.

- ~~~~~ Spartina limits.
- xxxxx Tide drift.
- Boundary gravel covered area.

Map 11
 Bath
 House





PLATE I

THE SNAKES OF THE LAKE MAXINKUCKEE REGION.*

BARTON WARREN EVERMANN AND HOWARD WALTON CLARK.

The total number of species of snakes known from the vicinity of Lake Maxinkuckee is ten. This number is not large; doubtless more thorough field work would increase the number slightly. While the species are not numerous, several of them are fairly abundant in individuals. This is particularly true of the common garter snake and the water snake. The former of these may be seen in suitable situations on almost any warm day from early spring until late in the fall, while the latter is almost equally frequent from the middle of summer to early fall about the borders of Lost Lake and along the Outlet.

Nearly all, perhaps all, of the species bear some relation to the life of the lake, some of them feeding on fishes when opportunity offers, and all feeding upon frogs. Only one of the species of snakes known from the Lake Maxinkuckee region is poisonous; that is the little prairie rattlesnake which, fortunately, is not abundant.

SPECIES OF SNAKES.

1. *Storeria dekayi* (Holbrook).

DE KAY'S SNAKE.

This pretty little snake occurs sparingly throughout the eastern United States and westward to Colorado and Wyoming. At Lake Maxinkuckee it is one of the rarest species. Our collection from about the lake contains only three examples, viz., one, No. 33529, U. S. National Museum, obtained October 8, 1900, and two others taken on October 17, 1907, one near the Outlet, the other on the east side of the lake.

This species is known also as Brown Snake and Ground Snake, the former because of the color, the latter because it is so frequently found burrowed in the ground.

It is not only a harmless little snake, but it is useful, its diet con-

*Published by permission of Hon. Hugh M. Smith, U. S. Commissioner of Fish and Fisheries.

sisting chiefly of crickets, grasshoppers, and other noxious insects. It is also fond of earthworms and slugs.

It reaches a length of only a foot or less. Its color is a grayish brown, with a lighter band or line along the back, on each side of which is a dotted line; there is also a dark patch on each side of the occiput, and the under parts are grayish. Scales in 17 rows; ventral plates 120-138.

2. *Thamnophis proxima* (Say).

RIBBON SNAKE.

This species is found from Wisconsin to Mexico. At Maxinkuckee it is one of the rarer snakes. The only example (No. 02779, or 33545, U. S. Nat. Mus.) in our collection was secured September 21, 1900, near Lost Lake, southwest of Mr. Green's house. Another was seen nine days later south of the lake.

It is a very slender, graceful snake. It is probably not rare in the weedy patches west of Culver, particularly about old, drained lake-beds where the ground is still wet and where there are occasional pools. In the spring of 1901, four were seen, two on April 9 at the drained lake west of Culver, one April 30 at Culver Creek, and one May 20 in Hawk's marsh.

In habits this species does not differ greatly from other garter snakes. It delights in marshy situations and is not averse to an occasional short stay in the water. Its food consists chiefly of small frogs, toads and insects, with an occasional small fish.

This snake may be known by the following characters:

Lateral stripe on third and fourth rows of scales; scales in 19 rows, little or not at all spotted; color chocolate brown, with three yellow stripes; light brown below lateral stripes; ventral plates 150 to 160; tail about one-third the total length which rarely exceeds thirty-six inches.

3. *Thamnophis sirtalis* (Linnaeus).

COMMON GARTER SNAKE.

This is one of the most variable as well as one of the most widely distributed of all snakes. It or its subspecies may be found in nearly all parts of the United States, and it is by far the most abundant snake about Lake Maxinkuckee; it is probably more numerous than all other species combined.

It may be found in all sorts of situations; in cultivated fields and

gardens, about yards and barn lots, in grassy meadows and in open woodland, in marshy ground along streams and about lakes, and particularly along paths and public highways. It perhaps most delights in reedy, boggy places and lake margins. It is the first snake to be abroad in the spring and one of the last to go into hibernation in the fall. The first warm days of spring will rouse them from their winter's sleep and bring them forth to bask in the sun. Then they may be found usually lying at full length on a mass of dead grass along a fence-row or in some such situation, well exposed so as to get the full effect of the sun's warmest rays. Here they will lie quietly through the middle of the day soaking out the accumulated chill of the long winter. Thus they will pass several days before they begin to move about or to seek food.

In the fall they appear to be active to the last, continuing to eat until they go into winter quarters. At this season they seem to move about more than usual, perhaps because searching for suitable hybernacula. It is at this season that one so frequently observes their tracks across the dusty highway and when so many are run over and crushed on the public roads and by railroad trains.

Numerous examples were noted about the lake and in the surrounding country, and many specimens were collected. Examples were noted in April, July, August, September, October and November, the earliest date being April 9 and the latest November 22.

A female 3 feet long was killed July 26 and 40 young, each 6 to 7 inches long, were taken from her body. This and all other species of the garter snakes are viviparous, bringing forth their young alive. Dr. J. Schenk, of Mt. Carmel, Ill., reports that 78 young, 3 to 7 inches long, were taken from a female 35 inches long.

The garter snake has quite a varied menu. They are known to feed upon insects, insect larvæ, small rodents, young birds and bird eggs, toads, frogs, angleworms, small mollusks, tadpoles, salamanders, and small fish. Frogs, toads, fish, shrews and field mice doubtless constitute the major portion of their diet. One found dead on the railroad tracks near the elevator in the late autumn of 1906 was examined. It was quite fat, as snakes are likely to be at that time of year. The stomach was empty of food, but contained a few ascaris-like parasites.

One the whole, however, this snake is beneficial to the farm and should be protected. The disposition which most people have to kill every snake on sight is entirely irrational and wholly unjustifiable.

This creature, like many other snakes, is protected by an abominably sickening odor, not noticeable at a distance, but as disagreeable a smell as one is apt to encounter. This odor, however, is noticeable only when the snake has been annoyed and has become angered. When angry it sometimes flattens out after the fashion of the blowing adder.

There is a great variation among the individuals of this species found about the lake, and two or more subspecies should probably be recognized. We, however, have grouped them all under the species.

From all other snakes found about the lake, particularly the ribbon snake which it most resembles, this species may be readily known by its having the lateral stripe on the second and third instead of the third and fourth rows of scales. This species is also stouter, the tail being one-fourth the entire length. Color olivaceous, dorsal stripe narrow, obscure; three series of small dark spots on each side, about 70 between head and vent; side and belly greenish; lateral stripe rather broad, but not conspicuous; colors generally duller than in other species; ventral plates 130 to 160. Length 2 to 4 feet.

4. *Thamnophis butleri* Cope.

BUTLER'S GARTER SNAKE.

This is the rarest of the species of garter snakes which occur at the lake. The only example we have seen was found freshly killed just south of the Indiana boathouse on the east side of the lake July 23, 1900. It is No. 33544 (02716), National Museum.

It may be known from other garter snakes of the region by the location of the side stripes which are on the second, third and fourth rows of scales, which is not the case with either of the other species.

5. *Natrix sipedon* (Linnaeus).

WATER-SNAKE.

The water-snake is a common and well-known snake throughout the whole eastern United States as far westward as Kansas, and is tolerably abundant throughout its range in wet places, such as streams, ponds and lakes. About Lake Maxinkuckee it is to be found along low bits of shore such as that about Norris Inlet and the various other inlets of the lake, and near the Outlet. One of its favorite haunts is that portion of the Outlet between the two lakes. Next to the common garter snake this water snake or "moccasin" is the snake most frequently seen about the lake.

We have records of numerous examples seen, the earliest date being May 3 and the latest August 29. It is probably most abundant in June. Definite dates are as follows: In 1899, one seen July 11 and another August 29. In 1900, one seen July 13, 17 and 20, all on the west side; one seen on east side of Lost Lake August 1, one at Fish Commission Station August 7, and one near the Inlet August 16. In 1901, one in Culver Bay May 3; a large one on west side May 6, one near Farrar's May 23; a large one on Long Point June 2; another on Long Point June 16; one at Outlet June 19; one 3 feet 9 inches long on west side June 22; and a small one on Long Point June 24. In 1906 a large one found dead on Long Point August 15, a small one in Green's marsh, one at the Outlet and one on Yellow River August 16. During the summer of 1906, after the dam was thrown across the Outlet at the railroad bridge the water in the Outlet below the dam became very low, and water snakes could be found along the edge of the water almost any time a visit was made to that place.

This is the species more often seen in the water than any other. It delights to lie coiled on some old log or root in or at the edge of the stream, or on the timbers at the dam or the logs of drift material. It inhabits rather open woodland ponds in great abundance, and in such places they often collect several together on projecting logs. In such situations it lies in wait, basking in the sun, making short excursions now and then into the water after fish or frog, or dropping quietly into the stream when disturbed by the near approach of any one. Then it hides under the bank, only its head being out of the water, or else swims swiftly away and out of reach. While swimming it usually keeps its head above water, but when closely pressed or annoyed it will go entirely under and swim along on or near the bottom.

The water-snake is frequently called "moccasin" or "redbelly" and is by many believed to be deadly poisonous. Its bite is, however, entirely harmless, and it is very different from the venomous "water-moccasin" or cotton-mouth of the south.

Although the water-snake is non-venomous, it has very little to commend it. It is repulsive in appearance and spiteful in temper. It is more destructive to fishes than any other of our snakes; indeed, it seems to subsist chiefly on fish. It will eat any kind of fish it can catch, though it doubtless prefers the soft-rayed species, such as the minnows, suckers and the like; it surely finds them easier to handle than the spiny-rayed species such as the bass and perch. We have found many different fishes

in the stomach of the water-snake; among them we may mention suckers of various species, various minnows, bass, rock-bass, sunfish, eel, carp and catfish. One large water-snake was found that had attempted to swallow a large catfish but the catfish straightened out and set its pectoral spines, and the snake, being unable to get the fish either up or down, perished, a victim of his own greed.

Besides fish the water-snake feeds also on frogs, crawfish and young birds.

The water-snakes mate early in spring, soon after coming out of their winter quarters, and then sometimes congregate in numbers of four or five together. The species is viviparous. In August, 1899, an old snake was found on the railroad track near the ice-houses. It had been run over by a train and ten young, which it contained, were prematurely liberated.

The water-snake probably comes out and basks on bright days in autumn after it has ceased taking food. One found dead near Farrar's in the autumn of 1906, October 20, was cut open and the stomach found to be empty, except for some ascarid-like parasites. The mesenteries were well loaded with a supply of fat, probably for the subsistence of the snake during its winter hibernation. It contained 30 ova, 15 on each side.

Color, brownish; back and sides each with a series of large, square, dark blotches alternating with each other, about 80 in each series; belly with brown blotches; rows of scales 23; ventral plates 130 to 150. Length 2 to 4 feet.

6. *Collopettis vulpinus* (Baird & Girard).

FOX SNAKE.

This large and beautiful snake ranges from New England westward to Kansas and northward. It does not appear to be common about Lake Maxinkuckee, as our notes record but eight examples, as follows: A fine example on the west shore of Lost Lake early in July, 1900, and another large one near the same place July 8; one seen near Lost Lake, September 3, and a large one gotten on Long Point September 25; one in Walley's woods August 25; another on Long Point September 25; one about 6 feet long August 14, 1906, west of Culver near the beaver-dam prairie on the road to Bass Lake; and a large one near the Gravel Pit early in June, 1907. Individuals seem most frequent in late summer or early fall.

The fox snake, often called the pine snake, frequents the dry, open woods and the neighborhood of briar patches and copses. We have never

observed it in the water or on the immediate lake shore. It is often called the pilot snake and is supposed to have some mysterious connection with the rattlesnake. Though entirely harmless, it is one of the most viciously disposed snakes. When provoked, as Dr. Hay observes, it shows its irritation by vibrating the tip of its slender tail, which, when striking a crumpled leaf or any other small object, may produce a rattling noise very much like that made by a rattlesnake under similar circumstances. A large example caught near Bass Lake August 14, bit Professor Wilson on the hand, causing blood to flow freely but producing no serious effect.

While entirely harmless, its habits are not unlike those of the black-snake and it doubtless destroys many eggs and young of ground-nesting birds. Besides these, its food consists of mice and other small rodents, the larger insects and their larvæ. It probably feeds to some extent on frogs and toads, but we have no evidence that it ever catches fish.

This is a large, light brown snake, with squarish, chocolate-colored blotches about 60 in number; scales in 25 rows; ventral plates 200 to 210; vertical plate broader than long.

7. *Bascanion constrictor flaviventris* (Linnæus).

BLUE RACER.

This common and familiar reptile, also known as the black snake or black racer, is found pretty generally distributed throughout the eastern United States and southward. It frequents open woodland, old fence rows and all places where dead leaves are common. It is the largest of the snakes of this region. It is an active, vigorous snake, moving over the ground with great rapidity. It is not a coward, as are most snakes, but will, on occasion, attack a person when disturbed, coming toward one rapidly and with head raised one or two feet. Cope says "the constricting power of the black snake is not sufficient to cause inconvenience to a man, but might seriously oppress a child. The pressure exercised by a strong individual wound round the arm is sufficient to compress and close the superficial veins, and cause the muscles to ache, but it is easy to unwind the snake with the free hand and arm." The black snake is harmless, and its bite, which it rarely inflicts, only amounts to a serious scratch.

The black snake is, in some respects, a useful species. Its food consists chiefly of field mice, white-footed mice, and other noxious animals. It also feeds upon frogs, toads, birds' eggs and young birds, and probably does more harm than good. The greatest objection to it is its disposi-

tion to rob birds' nests of their eggs and young. Ground-nesting birds are particularly apt to suffer from the depredations of the black snake; and those species such as the song sparrow, catbird, thrasher, robin, dove and redwing, which place their nests not far above the ground, and the bluebird, chickadee, and downy woodpecker, which deposit their eggs in holes in trees or snags not many feet up, are often despoiled of their eggs or young by this snake.

We have often seen black snakes coiled up on limbs of trees or crawling about among limbs several feet above the ground evidently searching for birds' nests. One of us remembers seeing a bluebird greatly disturbed by a large black snake which was apparently about to climb to the bluebird's nest which was in a hole only 3 or 4 feet up in an old elm snag. Coiled up at the foot of the snag, its head elevated perhaps a foot or 18 inches, the snake watched the bird intently, its head moving this way and that and following closely the movements of the bird which fluttered incessantly about the snake and was probably as completely "charmed" or under the power of the snake as birds ever get. When approached the snake became frightened and crawled away among the bushes; and then the bird flew to a limb near by.

A friend who is a close observer of animals tells us that he once saw a ruffed grouse fighting a black snake which was endeavoring to rob the grouse's nest. He shot the snake, and the grouse, after showing some astonishment, feigned lameness to lead him away from the nest.

Another friend says that he once saw a chipmunk "charmed" by a large black snake. The chipmunk was on a log about 12 feet long, the snake at one side near the middle of the log and with head elevated somewhat more than the height of the top of the log. The chipmunk when first seen was uttering the well-known chirping note so expressive of solicitude and running back and forth on the log, at first the full length of the log, then less and less until it ran but a few inches each way from the snake whose head all the time moved to the right and to the left, following closely the movements of the little rodent. At the same time the snake's tail, elevated and rigid, was rapidly vibrating and making a noise not unlike that made by a rattlesnake. Unfortunately the observer shot the snake without waiting to learn if the chipmunk were really in any manner under the control of the reptile.

The black snake is not rare about Lake Maxinkuckee. Our notes record seven or eight individuals seen at different times. The earliest rec-

ord is the last week in May and the latest October 14. A large example seen east of Lost Lake on the latter date was quite stupid and declined to move. A 4-foot individual seen in Walley's woods was evidently blind, due to shedding its skin which was so loose that it slipped off when the snake was handled. The eyes were white, and the snake instead of seeing, apparently listened. Another was seen in Walley's woods September 21, 1900. On August 13, 1906, a very large one was seen half-concealed in the briars near the ice-houses. When approached it made its tail rattle among the dry leaves precisely like a rattlesnake. On August 14, 1906, a large one was caught near Bass Lake. Another, 5 to 6 feet long, was seen in Walley's cornfield September 20, 1907. It was coiled loosely at the base of a corn-stalk and seemed disinclined to move, though it stuck out its tongue repeatedly.

This snake is usually lustrous blue-black or pitch-black above and greenish below; chin and throat white. Young olive, with rhomboid black blotches. Body very slender; eye large, scales in 17 or 19 rows; ventral plates 170 to 190. Length 4 to 5 feet.

8. *Lampropeltis doliiatus* (Linnaeus).

HOUSE SNAKE.

This is the common house snake or milk snake so abundant in most of the upper Mississippi Valley States. It does not appear to be very common, however, about Maxinkuckee. The only example seen by us was obtained July 28, 1899, at our station near the Arlington Hotel. It is one of the mildest and most useful of snakes and feeds largely upon the various species of small noxious mammals. Its habits, however, are not entirely beneficial, as it will, on occasion, eat such hens' eggs and birds' eggs as it may find.

We have never seen it swimming in the water and do not know whether it ever feeds on fishes or other aquatic animals.

Color, grayish, with 3 series of brown, rounded blotches bordered with black, about 50 in the dorsal row; an arrow-shaped occipital spot; belly yellowish-white, with square black blotches; dorsal scales in 21 rows. In the young the dorsal blotches are bright chestnut-red inside of the black margins, and the spaces between are sometimes white or clear ash.

9. *Heterodon platyrhinus* Latreille.

HOG-NOSED SNAKE.

This interesting reptile, also known as spreading adder and blowing viper, is found throughout the eastern United States. It is a common and well-known species in most parts of Indiana.

It frequents dry situations such as cultivated fields, old fence-rows, open pastures and roadsides; also dry hillsides and the banks of streams. At times it may be seen along water-courses and the shores of ponds and lakes. We have rarely observed it in meadows or on wet or marshy ground; nor have we noted it often about human habitations.

Although not often seen in the immediate vicinity of this lake, it is probably not uncommon in suitable situations, especially in dry sandy regions. It appears to be very well known among the inhabitants of the region, and is held in great dread by most of them; even its breath is supposed to be fatal. From its method of defending itself by appearing very terrible, a habit which has perhaps given its evil repute, it is one of the most interesting snakes in the region.

One was taken in Walley's woods on a bright day in the spring of 1901. When first approached it assumed a threatening attitude and gave vent to loud hisses; it then broadly flattened out the neck, and the bright colors and color-pattern, which had been more or less concealed by the scales, now stood out vividly, the color markings on the back of the neck standing out with especial clearness. When the snake found that none of those tactics availed, it stiffened out and appeared to be dead, and was easily picked up and placed in the collecting can.

During the summer of 1906 a large example of this species was seen on the shore of Lost Lake, but it escaped into a hole in the bank. In the autumn of the same year a young example about 5 inches long was captured near the ice office; and frequent reports of the species having been seen, were heard.

The bite of this snake is entirely harmless—even if it could be induced to bite. From the nature of its food, it is one of our beneficial snakes; it eats very few fishes, but subsists on frogs, mice, and insects, and their larvæ, or grubs. Instead, therefore, of meriting the persecution which it meets almost everywhere, it is well worthy of protection.

From all other snakes of this part of the State, this species may be known by its habit of flattening out both its head and body marvelously.

In color, it is brownish or reddish, with about 28 dark dorsal blotches, besides lateral ones and half-rings on the tail; sometimes the color is nearly uniform black. Vertical plate longer than broad, about equal to the occipitals; ventral plates 120 to 150; scales in 23 or 25 rows. Maximum length about 2 feet.

10. *Sistrurus catenatus* (Rafinesque).

PRAIRIE RATTLESNAKE.

This species, known also as the Massasauga, is likely to occur in all prairie regions from Ohio to Minnesota and southward. In Indiana it is known only from the northern portions of the State. It is the only poisonous snake occurring about Lake Maxinkuckee. All the other species found in that region or elsewhere in northern Indiana are entirely harmless. Formerly the Massasauga was abundant throughout this part of the State, but with the settling up of the country and the draining of the prairie grass-land and the marshes, it has become wholly exterminated in many places and practically so in many others. About Maxinkuckee, however, and elsewhere in Marshall County, it is far from extinct. It is apt to be found in any and all suitable places such as prairie meadows, about the borders of vanishing lakes, and in prairie marsh-ground anywhere.

In May, 1891, when the spring meeting of the Indiana Academy of Science was being held at Lake Maxinkuckee, several specimens were caught by members in attendance, chiefly in marshy ground about the lake. About 1896 a young man on the eastern side of the lake was bitten on the leg by one. The leg remained swollen for some time and complete recovery was very slow. On August 6, 1899, one was caught on Long Point between the Scovell and Walter Knapp cottages. It was 23 inches long and had five rattles. On August 3, 1900, one was killed two and one-fourth miles south of Arlington station. It was 18 inches long and had two rattles and a button. Several weeks earlier, near the same place, a dog was bitten by one, without fatal results. On August 26 a small one was killed on the east side of the lake near the T. W. Wilson cottage. On the same day one was killed in a field on the Hawk farm south of Culver. It was about 2 feet long and had nine rattles. Another young individual was killed September 3 on the east side, two and one-half miles southeast of the Maxwell cottage, and one with nine rattles was killed September 26, 1907, in a meadow on the Newman farm, four miles southeast of Culver.

These are all the records we have of the occurrence of the prairie

rattlesnake in the immediate vicinity of Lake Maxinkuckee. We have heard, however, of numerous examples being killed in marshy meadows northwest, west and south of the lake. In those regions there are numerous and considerable meadows of the wild grass or sedge, *Carex stricta*, which are cut in the early fall by farmers and others for hay or for use in the ice-houses, and other purposes. It is then that this venomous snake is met with most frequently.

Though habitually dwelling in marshy situations it is sometimes seen on higher, open ground. It is rarely seen in open woods or dry thickets.

We know but little about the habits or food of this snake. It apparently does not wander far but remains close about the particular marsh in which it makes its home. They are quiet and not easily disturbed or angered. When observed they will be still or quietly glide away unless interfered with. Then they will usually coil, assume a threatening attitude and rattle more or less. The rattling, however, soon ceases, to be renewed only when again provoked.

The Massasauga is known to feed on frogs, crawfish, meadow mice and shrews. We do not know that it ever feeds on fishes, but it is more than probable that it would not disdain to eat mud minnows or any other small fishes it might find in its swampy habitation.

The one fact that this is a venomous snake is sufficient reason for its extermination.

The species is viviparous, the young being brought forth alive. There are usually about six in a brood, each 4 to 6 inches in length when born. The birth of the young generally takes place about the first of September.

The prairie rattlesnake may be known from others of this region by the large, flat, triangular head on a slender neck, the presence of a deep pit between the eye and the nostril, the long, erectile, perforated poison-fang on each side of the upper jaw, and, usually, the presence of a rattle on the tail.

Color, brown or blackish, with about 7 series each of about 34 deep chestnut blotches, blackish exteriorly and edged with yellowish; a yellowish streak from pit to neck; body sometimes all black; scales in 23 or 25 rows; ventral plates 135 to 150. Length $2\frac{1}{2}$ to 3 feet.

STIRRING AS A TIME SAVER IN GRAVIMETRIC ANALYSIS.

W. M. BLANCHARD.

While making experiments recently on the deposition of metals with a rotating anode, it occurred to me that various gravimetric analyses might be greatly facilitated by rapid stirring of the precipitate. That stirring greatly facilitates precipitation is known to every one, but I do not know that any data have been recorded to show just how efficient the stirrer may be, hence these experiments. These results are very surprising.

A stirrer was made from a small glass rod about two millimeters in diameter. It had the shape of the letter T, the arms, each about eighteen millimeters long, being flattened at the ends and turned so as to resemble an ordinary propeller. The stem was about ten centimeters long and was attached to a wheel run by a small electric motor, which in turn was connected with an ordinary 110 volt lighting circuit. There was thrown in the circuit a small lamp bank so that the speed of the motor might be varied. The following analyses were carried out with the stirrer making an average of 900 revolutions per minute.

1. *Estimation of barium in crystallized barium chloride.* The sample of the pure salt, 0.2330 gram, was weighed out in a 150 c. c. beaker, diluted to about 50 c. c., acidified with hydrochloric acid, heated to the boiling point, treated with slight excess of dilute sulphuric acid, *stirred four minutes*, immediately filtered, washed, ignited, cooled, and weighed. Found for barium 56.25%; calculated, 56.23.

2. *Estimation of calcium in pure calcium carbonate.* A small sample of the purest calcium carbonate, 0.2225 gram, was transferred to a 150 c. c. beaker, converted into the chloride, diluted to about 50 c. c., heated to boiling, treated with slight excess of ammonium oxalate, made alkaline with ammonia, *stirred four minutes*, then filtered, washed, ignited, heated over the blast lamp, cooled and weighed. Found for calcium oxide 56.04%; calculated 56.03.

3. *Estimation of copper in cupric sulphate.* A small sample of recrystallized pure sulphate, 0.2000 gram, was weighed in a 150 c. c. beaker,

diluted to about 50 c. c., heated to boiling, treated with a few drops of sodium bisulfite solution and then with slight excess of ammonium sulphocyanate, *stirred three minutes*, filtered (Gooch), washed, dried at 130°, cooled and weighed. Found for copper, 25.42%; calculated, 25.46.

4. *Estimation of chlorine in sodium chloride.* A small sample of Kahlbaum's purest sodium chloride, 0.2162 gram, was dissolved in about 50 c. c. of water in a 150 c. c. beaker, acidified with a few drops of nitric acid, treated with slight excess of silver nitrate solution, *stirred two minutes*, filtered (Gooch), washed, dried at 130°, cooled and weighed. Found for chlorine 60.64; calculated 60.63.

DePauw University.

THE ALUNDUM CRUCIBLE AS A SUBSTITUTE FOR THE GOOCH CRUCIBLE.

GEORGE L. CLARK.

In order to test the efficiency of the recently introduced unglazed Alundum crucible, when used for the purposes in quantitative analytical chemistry generally assigned to the ordinary Gooch crucible, four different series of analyses were undertaken. These involved the determination of silver by precipitation as silver chloride, copper as cuprous sulpho-cyanate, aluminium as aluminium oxide from ignited aluminium hydroxide, and barium as barium sulphate. Such a selection was made in order to obtain as widely different precipitates as possible, as regards size of particles, ease of filtration and media in which produced, and at the same time be in general usage.

Both the Alundum and Gooch crucibles permit filtration, drying and weighing without disturbing the precipitate, but the porous nature of the former of course does away with the preparation of an asbestos mat. To discover whether or not such an advantage as well as others claimed for the Alundum crucible by its manufacturers, such as capability of withstanding very high temperatures, is sufficient to warrant its wide adoption for use in accurate quantitative analysis, was therefore the object in view in this study.

One crucible only was used for the determinations in one series, in order to discover what would be the effect of continuous usage and how thoroughly it might be cleaned in preparation for the next analysis. In each case the empty crucible was heated thoroughly for one hour in the drying oven at the temperature at which the precipitate was later to be dried. The apparatus for the filtration was that used with the Gooch crucible, since any more complicated or expensive apparatus would *per se* be a distinct disadvantage.

I. The determination of AgNO_3 as AgCl .

In these analyses 50 c. c. portions of a solution, each containing .2432 grams of pure AgNO_3 were used. In the first trials a solution of Kahlbaum's chemically pure NaCl was used to precipitate the AgCl from the hot solution of AgNO_3 while rapidly stirred. Stirring was continued for two minutes, the precipitate allowed to settle, filtered through the crucible, washed with water and then dried at 140° .

First analysis: Percent Ag in AgNO_3 calc. 63.50, found 65.24. No. of washings, 8.

The crucible was then washed thoroughly with pure water until all apparent traces of AgCl had been removed, dried at 140° and again weighed. Gain in weight, .0162 gr.

Second analysis: Percent Ag calc. 63.50, found 64.04. The crucible was now thoroughly washed by suction with NH_4OH to remove all AgCl and water to remove all NaCl from the pores. Weighing gave the original value for the crucible.

Pure dilute HCl was substituted for NaCl so that the error might not be due to the precipitating agent.

First analysis: Percent Ag calc. 63.50, found 63.52.

Second analysis found 63.60.

Finally a solution of purified KCl , a slightly more soluble salt than NaCl was tried. The empty crucible was in all cases washed with NH_4OH and water and it varied in weight by only one-tenth milligram.

Percent Ag calc. 63.50, found 63.98.

In all the analyses the AgCl precipitate was easily handled, the difficulty arising in the seeming impossibility of washing out of it and the pores of the crucible the NaCl and the KCl .

II. Determination of Cu in $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ as CuSCN .

This analysis seemed particularly adaptable because of the extensive use of the Gooch crucible necessitated in it. A solution of crystals of very pure $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ was made such that each 50 c. c. contained .2136 grams. Precipitation of CuSCN was affected from the hot solution in the presence of an excess of H_2SO_4 by means of $(\text{NH}_4)_2\text{SCN}$. Drying was at 140° .

Percent Cu in CuSO_4 calc. 25.46 found (1) 26.80 (Stirred 2 minutes and not permitted to settle.)

(2) 25.24 (Stirred 4 minutes rapidly and digested 15 minutes.)

(3) and (4) 25.39 (Stirred 4 minutes rapidly and digested 30 minutes.)

By thoroughly washing with water, drawn both ways through by suction and drying at 140° , the crucible maintained a weight varying only by

one-tenth milligram. Although the particles of CuSCN are much smaller than those of AgCl , they were retained by the pores of the crucible.

III. Determination of Al in $\text{Al}_2(\text{SO}_4)_3$ as Al_2O_3 .

The effect of high temperature upon the Alundum crucible as well as its applicability for filtration of gelatinous precipitates, were discovered by this analysis. Chemically pure $\text{Al}_2(\text{SO}_4)_3 \cdot 18\text{H}_2\text{O}$ was dehydrated by gentle dessication and final heating for $4\frac{1}{2}$ hours at 140° . A solution was prepared such that each 50 c. c. contained .2154 grams of $\text{Al}_2(\text{SO}_4)_3$. $\text{Al}(\text{OH})_3$ was precipitated from the hot solution in the presence of NH_4Cl by just a sufficient amount of NH_4OH . After boiling, stirring and allowing to stand, the precipitate was filtered through the crucible previously heated in the flame of a blast lamp for several minutes, and washed with water containing small amounts of NH_4Cl and NH_4OH . In seven attempts however in spite of all precautions and lengthened time of treating the precipitate, part was drawn through the crucible with the filtrate. The last three determinations were made with an entirely new solution of $\text{Al}_2(\text{SO}_4)_3$, each 50 c. c. containing .2001 grams.

EFFECT OF HEATING ON THE CRUCIBLE.

Except in one case where the change was negligible, the crucible lost in weight consistently from one analysis to the next. The first crucible was pulled apart by suction while cleaning with water following the third analysis, and the second in exactly similar fashion after the fourth analysis in which it was employed, showing that the material becomes extremely brittle after heating several times in the blast lamp.

IV. The determination of Ba in BaCl_2 as BaSO_4 .

The extremely small size of the particles of BaSO_4 under ordinary conditions recommended a further test of the crucible. The pure BaCl_2 was thoroughly dried by heating three hours at 140° and a solution made such that 50 c. c. contained .2007 grams. The hot solution slightly acidified with HCl was treated with pure dilute H_2SO_4 while vigorous stirring was maintained for five minutes, after which the BaSO_4 was allowed to settle for five minutes.

Percent Ba in BaCl_2 calc. 65.96 found (1) 65.86
 (2) 65.98
 (3) 65.79

The crucible was washed with water under suction and maintained

practically a constant weight. A final trial of the effect upon the weight of rubbing simply with the finger was made. The crucible lost .0039 grams.

SUMMARY OF ADVANTAGES.

- (1) The crucible was used to advantage in determining Cu as CuSCN , Ba as BaSO_4 and Ag in AgCl when HCl was used as the precipitating agent.
- (2) Manipulating was found to be surprisingly simple and rapid when an analysis was once under way.
- (3) Nearly a constant weight was maintained by the crucible if not heated over 140° when washed with water.

SUMMARY OF DISADVANTAGES.

- (1) It was found difficult at best to wash precipitates free from precipitating agent, especially if the latter should be left as a solid residue upon evaporation of the solution.
- (2) Suction was necessary, and, in the case of $\text{AgCl}, \text{NH}_4\text{OH}$, to thoroughly cleanse the crucible.
- (3) It was found unadapted to filtration of such gelatinous precipitates as $\text{Al}(\text{OH})_3$.
- (4) Heating six or eight times in the blast lamp was sufficient to render the crucible so brittle as to be broken even by suction, and a consistent loss in weight was observed.
- (5) In three cases the first use of crucibles led to results greatly in error.
- (6) Abrasion or friction was found to have a marked effect upon the weight of the crucible.
- (7) The ordinary digestion of fine precipitates to be filtered was not avoided when it was used.

Like the Gooch crucible therefore, the Alundum crucible apparently has only a limited field of usage but within that field it should be of considerable worth to the analyst.

THE CORRELATION OF HIGH SCHOOL AND COLLEGE CHEMISTRY.

JAMES BROWN.

This subject I submit for consideration, not as one who has anything final to offer, but as a teacher who has considered several different systems and has tried some of them.

Inasmuch as the objects sought in the various high schools and college courses differ, it is difficult or impossible to devise any system of correlation which will suit all cases with the maximum of efficiency. Local conditions and previous training of students, as well as the future plans of the students, so far as these are definite, must be determining factors. In any case, efficiency rather than convenience should be our guide.

In considering this question I have found it convenient to propose three alternatives for students who have completed a high school course in chemistry and elect to continue the subject in college. The alternatives are as follows: First, to admit the student at once to second year chemistry, usually qualitative analysis; second, to give the student the same course as those who have had no previous work in chemistry; third, to give to such students a special course in general chemistry.

The first alternative—to admit the student at once to second year chemistry—I do not favor for theoretical reasons and because my experience has found it unsatisfactory. In this case you have high school students, the nature of whose courses in chemistry has differed widely, subjected to the same prescription as college students whose courses have usually been more uniform and deeper. This is apt to be especially true because the college recitations and laboratory periods are usually longer and because, in a great many colleges courses in general chemistry more or less qualitative analysis is introduced. This enables the college student to start qualitative analysis at a somewhat advanced point.

On the theoretical side we find similar differences. The time is past, if it ever did really exist, when a course in qualitative analysis conducted in a mechanical way may be considered properly taught. The theory of the

subject is presented in our best text-books from the point of view of ionic equilibrium, the periodic system, and the electro-chemical series. Our best college text-books and laboratory manuals in general chemistry emphasize these same subjects. This, it seems to me, gives the correlation between general chemistry and qualitative analysis which is not secured by courses which do not place emphasis on these three subjects. Equations also must be well learned throughout all chemistry courses. We must not, to be sure, give too much time to equations to the exclusion of other parts of the science. But have you ever known a good chemistry student who could not write equations? I often wonder if equations are being neglected.

The second alternative—to put all students into the same course in general chemistry—admits of several interpretations. Shall we give full credit for the course to the student who has received an entrance credit in chemistry? This may mean duplication of credit. Such duplication exists in one form or another in some subjects. Shall we do the same in chemistry? This question is variously answered by different institutions.

Duplication of credit may be avoided by requiring different laboratory experiments and different written work in the laboratory and in connection with the text-book, from the two classes of students. This is rendered difficult by the different contents of the high school courses. Or we may avoid this duplication by giving only part credit for the college work to those who have entrance credit in chemistry. This may appear to the student to be work without credit, and is often opposed on those grounds.

The third alternative—to give a different course to the two classes of students—may be accepted in different forms. In some cases students have totally omitted the first part of the course, and taken the latter part entire. This I think is objectionable because of sins of omission and commission. The student should have much of what he omits in the first part, and duplicates much that is familiar to him in the second part. We may on the other hand give a shorter course covering the whole subject to our students with entrance credit, avoiding duplication of work which may be supposed to be familiar, and giving only what we think will impart the advanced point of view which we consider advisable.

This accomplishes in another way much the same end as the plan of assigning different work under the second alternative. These two plans are subject to the same difficulty. The students have had quite different courses in high school and do not well admit of the same diagnosis.

Will not a satisfactory solution of our problem be accomplished by the introduction into our high schools of the new courses in general science now being advocated? This would leave the specialization along different branches of science in the hands of the colleges and would enable us to treat all classes of students alike without fear of duplicating credit, or of omitting anything essential. Probably our high school science should be conducted with the purpose of enabling the student to interpret his daily environment. In college, however, while considering fully the interest of the student whose object in chemistry is cultural, we must be guided mainly by the professional student and by those who, for various reasons, wish to specialize in chemistry.

THE CHEMICAL COMPOSITION OF VIRGIN AND CROPPED INDIANA SOILS.

S. D. CONNER.

In November, 1913, the Soils and Crops Department of the Purdue Agricultural Experiment Station collected samples of a large number of typical virgin and cropped soils, with a view to determine the chemical composition, to see if there was any appreciable difference in them. The samples were taken with an auger and each sample represented not less than five borings to a depth of six and one-half inches. Subsoil samples were taken at the same time and represented the layer from a depth of 12 to 18 inches. Great care was taken to select fields where uniform and typical samples could be secured. The samples in each case represent a heavily cropped soil and an adjacent virgin soil of the same type which had never been cropped. The virgin soil samples were taken from a line fence row, or from a woodlot which had never been cultivated. The separate samples were properly prepared and analyzed for various elements. Also composite samples were prepared from the virgin soils, the virgin subsoils, the cropped soils and the cropped subsoils. The composites were made by taking equal weights of the separate samples and thoroughly mixing them. The analyses of the separate samples not being completed up to the present time, the analyses of the composite samples only are given in this paper.

There is a rather widespread idea that the chemist can take a sample of soil and by making a complete analysis, determine without any other information just what element is deficient in the soil and needed as a fertilizer. This is not true, and it is very seldom that an analysis alone will indicate the needs of a soil. The chemist can tell with a fair degree of accuracy just how much of each element is present in the soil, but he is not able from a chemical analysis alone, to say what various crops are able to extract from the soil. The ability to determine the fertilizer needs of various crops on different types of soil is more or less a matter of

experience and is based largely upon the results of field fertilizer tests. However, a chemical analysis of a soil is often of great benefit in studying problems of soil fertility.

It is generally recognized that the removal of plant food by crops is not the only factor which may change the composition of a cultivated soil. The agencies of wind and water play a very great part in effecting changes. Insects, worms and animals often work through the soil and subsoil, causing variations and intermixtures. The crops themselves bring up from below quite a little plant food and deposit it near the surface in the decaying roots and stubble. In spite of the fact that the tendency of nature is to build up and replenish the fertility of soil, there is no question but that the destructive system of cultivation that has been followed by the farmers of this country has more than counter-balanced nature's tendency to upbuild, and as a consequence the soil has become more or less depleted. It is believed that the analyses presented in this paper show what chemical changes have been effected in the average soil of Indiana by cropping it for from sixty to eighty years.

TABLE I.

Analysis of Composite (31) Virgin and Cropped Soils.

MATERIAL.	Virgin Soil.	Virgin Subsoil.	Cropped Soil.	Cropped Subsoil.
Insoluble silica, etc.	88.49%	87.30%	89.59%	86.37%
Potash (K ₂ O) (Acid soluble).....	.26	.36	.23	.34
Soda (Na ₂ O) (Acid soluble).....	.20	.20	.17	.21
Lime (CaO) (Acid soluble).....	.43	.34	.43	.51
Magnesia (MgO) (Acid soluble).....	.41	.60	.40	.60
Manganese oxid (Mn ₃ O ₄) (Acid soluble).....	.14	.08	.07	.09
Ferric oxid (Fe ₂ O ₃) (Acid soluble). Alumina (Al ₂ O ₃) (Acid soluble).....	5.03	8.01	5.31	8.60
Phosphoric acid (P ₂ O ₅) (Acid soluble).....	.12	.07	.11	.07
Sulphur trioxid (SO ₃) (Acid soluble).....	.06	.05	.06	.04
Volatile matter.....	5.28	3.20	3.88	3.25
Total.....	100.42	100.21	100.22	100.08
Total nitrogen.....	.18	.07	.13	.06
Total potash (K ₂ O).....	1.83	1.88	1.94	1.92
Total humus.....	1.98	.60	1.04	.40
Acid humus.....	1.16	.44	.84	.48

A glance at the analyses in Table I will show that although most of the soil ingredients have not changed enough to make any great difference in the chemical composition of the virgin and cropped soils, there are some notable exceptions.

The most serious losses from the standpoint of soil fertility are those of nitrogen, which shows a loss of 28%, and the organic matter, which shows a loss in the volatile matter of 26%, and in the humus of 47%. These losses are without doubt the main reason why our cropped soils are no longer as fertile as they formerly were. Fortunately the remedy for replacing nitrogen and organic matter is not beyond the means of the average farmer. Greater care in utilizing crop residues and barnyard manure, also the growing of legumes in a good crop rotation are necessary steps in replacing these vital losses. The purchase of organic matter, other than farmyard manure, is out of the question, while the use of nitrogenous fertilizers which often give profitable returns, can only be recommended as a temporary resort.

While the phosphoric acid and potash show only about 10% loss, it should be remembered that this 10% was the most available portion of these important elements. Fertilizer practice in the older and more worn lands of Indiana shows that there has been a loss in these elements and that in a great many cases their use as fertilizers is very profitable. Due to the fact already mentioned that the soil through natural agencies is constantly in motion, it should be pointed out that the addition of one or two tons of rock phosphate per acre to the land for the purpose of increasing the phosphorus content of it for a long time to come, is a practice of doubtful efficiency. There is a strong probability of loss of such fertilizer, due to removal by wind or water, or to being buried out of reach of the plants by these or other natural agencies. Smaller amounts of more available phosphorus or potash fertilizers, on the other hand, will be quickly utilized by the crops and hence not so liable to be lost. Experiments which have been conducted by the Purdue Experiment Station show greater profit from the use of acid phosphate than from raw rock phosphate. (Bul. 155, Purdue Experiment Station and the 27th Annual Report, Purdue Experiment Station, 1914.)

The analyses of calcium and magnesium in the virgin and cropped soils show no apparent change. This is rather surprising as we have been led to think of these elements, especially calcium, as being very soluble. The loss of lime, as reported by the Rothamsted Station, has been shown

to be from 500 to 1,000 pounds per acre per year. There is one important difference in the Rothamsted soil and the average Indiana soil, and that is in the fact that the Rothamsted soils in the experiments reported have from two to four per cent. calcium carbonate. The Indiana soils shown in these analyses, on the other hand, have no calcium carbonate. The calcium and magnesium in these Indiana soils are in the form of more or less insoluble silicates. The inference to be drawn, therefore, would be that there is no great loss of calcium or magnesium in acid soils in which these elements are in the form of silicates. This does not mean that these soils do not need lime for, as a matter of fact, they respond readily to the application of lime, which is needed for the proper growth of clover. The need for lime is greater now than it was in the virgin soils because the organic matter has been burned out of the cropped soil. Given two soils with the same calcium and magnesium content and the same degree of acidity but with different amounts of organic matter, the one with the greater organic matter content will grow better crops of clover and will not be in so great a need of lime as the other.

The virgin and cropped soils show no great difference in the content of sulphur. Experiments in Wisconsin and Kentucky have shown that in a number of instances sulphur has been reduced in soils by cropping.

Manganese shows quite a loss in the cropped soil. The effect of manganese on soil fertility is attracting more or less attention among soil investigators, and although nothing definite seems to be known about its action, it is possible that it does play an important part in agriculture.

The changes in the content of silica, iron and aluminum are believed to be of no importance as plant foods. They do, no doubt, have a very important bearing upon the physical constitution of the soil. The writer believes that the constitution of the silicates of iron, and especially of aluminum, has more to do with injurious soil acidity than any other factor.

The method of determining soil acidity (limestone required) in this work is that given in Bulletin 107 (Revised edition), Bureau of Chemistry, U. S. Department of Agriculture. This method shows a relative acidity in soils that is believed to more nearly represent toxic acidity than any other method, especially in soils containing much organic matter. It is interesting to note that while the acidity of the cropped soil has increased, the acidity of the cropped subsoil has decreased.

TABLE II.

FERTILITY IN VIRGIN AND CROPPED SOILS.

Pounds Per Acre in Two Million Pounds of Surface Soil and Four Million Pounds of Subsoil.

MATERIAL.	Virgin Soil.	Virgin Subsoil.	Cropped Soil.	Cropped Subsoil.
Volatile matter.....	105,600	128,000	77,600	130,000
Humus.....	39,600	24,000	20,800	16,000
Nitrogen.....	3,600	2,800	2,600	2,400
Potassium (Total).....	32,464	66,702	34,416	68,121
Potassium (Acid soluble).....	4,615	12,773	4,080	12,063
Phosphorus (Acid soluble).....	1,046	1,221	959	1,221
Calcium (Acid soluble).....	6,864	9,724	6,864	14,586
Magnesium (Acid soluble).....	4,953	14,496	4,832	14,496
Manganese (Acid soluble).....	2,016	2,304	1,008	2,592
Sulphur (Acid soluble).....	480	800	480	640
Limestone required (Acidity).....	60	2,600	100	1,120

Table II gives the pounds of the different elements in the plowed soil of 2,000,000 pounds per acre, and in the subsoil of 4,000,000 pounds per acre. It shows the same relative differences as given in Table I, but in different terms.

The writer wishes to acknowledge the assistance of Mr. J. C. Beavers of the Soils and Crops Department, also of Mr. J. B. Abbott, formerly of the same department, who collected most of the soil samples analyzed.

SEWAGE DISPOSAL.

CHARLES BROSSMANN, Consulting Engineer, Indianapolis.

Civilization and education has been accompanied by a wonderful growth of cities and has made the problem of sewage disposal one of civic, state and national importance.

Sanitation becomes of greater importance as communities become more congested.

It is only of late years that this question has received proper attention, the greatest progress having been made in the last few decades. The combined efforts of the scientist, chemist and engineer have been called upon to help solve this problem of ever-increasing importance.

Improper disposal of sewage has caused directly or indirectly a large percentage in the typhoid mortality rate.

The gathering of large numbers of people calls for additional safeguards and means of sanitation. In some instances sewage can be disposed of by dilution, discharging direct into large bodies of running water; but most streams are as a rule not of sufficient size, or are already so polluted that additional sewage would increase the burden already too large.

Generally sewage is diluted with the entire water supply of a city and is a dirty appearing water, containing a greater or less percentage of organic matter. There is usually enough organic matter present to make it disagreeable and to cause odors. The presence of various disease germs also make it a source of pollution to water bodies.

In general all methods of sewage treatment employ the principal of reduction through microscopic organisms. Bacteria of various kinds attack the organic compounds reducing them to simpler forms, doing so through successive stages. Reduction takes place through two classes of bacteria, namely aerobic (thriving in the presence of oxygen), and anaerobic (thriving in the absence of oxygen). These two processes occur in septic, Imhoff or other tanks and in various forms of filters.

The most prevalent form of getting rid of sewage is by dilution. Where the stream is sufficient in size to allow proper oxidation the sewage

will be properly taken care of without objectionable odors. Such a stream however should have a flow of about 300 cubic feet per minute for each 1,000 inhabitants. Instances where disposal by dilution alone is sufficient are not many and usually some additional treatment is necessary, suitable to local conditions.

If the stream into which the sewage is to be discharged allows of partial dilution, treatment by some properly designed form of tank may be




FIG. 1.

Type of Septic Tank Suitable for Ordinary Dwelling.

sufficient, but tank treatment alone will not always suffice. Tank treatment is but a step in the purification of sewage and should usually be followed by some form of filtration or after treatment. Various forms of tanks can be used but the type and size that insures the best results can only be determined after proper investigation of all conditions—as the number of people, amount of sewage, the rate and the time of flow, the

location of adjacent property and the size of the water course into which the treated sewage is finally discharged; all are important and enter into the proper solution of this important question. ?

Tank treatment, therefore, is essential as the first step in sewage reduction, and is necessary in order to retain and break down the solids, but it must not be supposed that it purifies the remaining sewage liquor. The tank treatment is necessary in preparing the sewage liquor for further purification. Such tanks can be made in the form of plain settling tanks, a septic tank, or a combination of both.

The public will universally call any tank (even a cesspool) a septic tank, and usually they believe that a septic tank absolutely purifies the sewage. Such is not the case, a reduction from thirty to sixty per cent. of




FIG. 2.

Septic Tank and Natural Sand Filter for Small Installations.

suspended matter and around thirty per cent. in organic matter is usually what takes place. The tank will not take care of very fine particles or colloidal matter. Such matter (colloidal) being in condition just between suspension and solution. The best results are obtained when the solids are taken out or retained as quickly as possible and the subsequent liquor remaining immediately treated. It is important that liquor be not retained too long or it will become in a toxic condition. Time is an important element in the proper design of a tank, also the state of the sewage in reaching the tank.

Septic tanks are usually designed for a rate of flow, of from eight to sixteen hours. The more modern type of tank with two compartments, one for settling and one for sludge digestion, are usually designed with a rate of flow of half (or even less) than the above.

The septic tank is usually a rectangular shaped chamber with several baffle boards, extending across to break the flow of the sewage. Such tanks should be covered as the organisms that break down the solids (known as anaerobic bacteria) thrive best in the absence of air and light. Septic tanks usually take some time to become operative, a scum mat forming at the top and sludge at the bottom. At intervals such tanks must be cleaned of the sludge. It was formerly supposed that just as much solid matter was turned to liquid and gas to offset the amount of solids




FIG. 3.

Imhoff Tank and Sand Filters Installed at Indianapolis Country Club

coming in, however, it has usually been found necessary to clean out the resultant sludge at intervals. (See figure 1.)

The Imhoff type of tank consists of two chambers, one for settling and one for the deposit and digestion of sludge. Such a tank, while somewhat more expensive than the plain tank is smaller and will give more uniform results, besides offering better means for after-treatment of the liquid and assuring a better solution of the sludge problem. (See figure 4.)

This type of tank has an upper settling chamber with a slotted opening at the bottom. The sewage in flowing through the upper chamber deposits

the solids into the sludge chamber below. In the sludge or digestion chamber below, the solids and organic matter is gasified and liquified independent of and without disturbing the settling matter above.

The gases of decomposition and the constant agitation in the lower part does not disturb the sewage in the settling chamber, furthermore albumen from the fresh sewage is not constantly added to the septic sludge; hence there is less odor and the sewage liquor is delivered in a fresher condition for after-treatment. The sludge from this type of tank dries out quicker, is in better condition for disposal, has less water content, and has different characteristics than sludge from a shallow tank which is kept in



FIG. 4.

Imhoff Tank, Julietta, Ind. Note Formation of Sludge on Sides.

constant contact with the sewage. Such double tank sludge soon becomes spadable like garden compost.

The tank treatment should be followed by dilution or some form of filtration. In some cases the sewage liquor from tanks can be discharged into a water-course. Usually it is necessary to use some form of filter or nitrification bed. This can be done in the following manner:

(1) In small plants by discharging the sewage into tile laid near the surface of the ground. Such ground must be suitable for the sewage to percolate through to a subdrainage system below. Such ground should be gravelly or of sand. (See figure 2.)

(2) By discharging the sewage into contact beds, viz., a water-tight bed, filled to a depth of several feet with broken stone or other hard

material, the sewage being automatically discharged on to the bed, retained a fixed period, and then discharged from the bed. In such a bed absorption and oxidation of the organic matter is accomplished by aerobic bacteria, viz., those which thrive in the presence of air. (See figure 5.)

(3) Sand filters: As the sewage from tanks can be discharged on sand filters—automatically dosed as in contact beds. Such sewage covers the surface of the bed and gradually works through to the underdrains below, the action being that of filtration and nitrification. If a very pure effluent is desired the sewage can be discharged from the tank to the con-



FIG. 5.

Contact Beds, Julietta, Ind.

tact bed and then be treated through the sand filters. In a properly designed plant this will give a very clear effluent. (See figure 6.)

Sprinkling Filters: In the larger plants sprinkling filters are largely used. These consist of beds of broken stone, usually of a depth of six feet or more and are arranged for good underdrainage. The sewage is automatically discharged over the top of the bed by sprinkling nozzles; trickles down through the stone and out through the underdrains. Such beds can be worked at a higher rate than any of the preceding methods, hence a smaller area is required, which makes this method more adaptable for large installations.

DISPOSITION OF SLUDGE.

The real problem in sewage disposal plants is the sludge problem. Engineers are just learning how to make sludge but in most cases have not found a satisfactory solution in disposing of it. In larger plants the sludge question is the stumbling block.

Sludge may be roughly divided in two classes, that from shallow tanks and that from deep tanks. Sedimentation tank sludge is a black semi-




FIG. 6.

Sand Filters, Julietta, Ind.

liquid mass which on being exposed to the air becomes offensive, giving off much gas and odor. The water contained from such sludge is usually 90 to 95%.

The sludge from septic tanks ranges all the way from 8 to 45 cubic yards per million gallons of sewage. Septic sludge which has been retained in tanks for a number of months undergoes a great change. The organic matter is attacked and partly gasified and liquified, which reduces the amount of sludge. Such sludge in well operated tanks is a con-

centrated mass containing from 80 to 90% of water; there is not as much odor to septic tank sludge as to the fresh sludge from the plain settling tanks.

The above outlines the principal methods of sewage disposal. It must however be borne in mind that the proper method is wholly decided by local conditions. Care must be exercised in order that the various factors affecting the problem be carefully considered.

A fuller realization of the sewage disposal problem is being evidenced throughout the country as the years go by, both by the state and health officers and city officials. The State has done a great deal of preliminary work in the way of sanitary surveys and this work should be heartily indorsed and commended. However, there is still much to be done, and




FIG. 7.

Water Fowl in Stream, Julietta, Ind., 100 Feet Below Sewage Plant.

Photo taken 2 years after plant was installed. Formerly all waterfowl died from drinking water.

undoubtedly some method of maintaining public control of the streams will have to be devised before the question of pollution can be properly taken care of.

There are so many different factors entering into this question that the best solution can only be worked out with a proper organization which will take into consideration every phase of the question and which can reach every district, affected, whether this territory be in one or more States. The state authorities should be given the power to pass upon every sewage disposal problem and they should have the proper means and support for doing this.

5.

ig.

ives-
and
of a
thor

con-
nor-
the
t be

ola-
aces


(1)
dis-
the
tars
em-
een

first
ally
ure
°C.
t of
nat-
i by

om-
no

6

|



JULIETTA SEWAGE DISPOSAL PLANT

CHAS. BROSSMANN
 INDIANAPOLIS
 CONSULTING ENGINEER
 SCALE 1/8 INCH=FOOT

General Plan. Section of Julietta Sewage Plant Showing Imhoff Tank, Contact Beds and Sand Filters.

Fig. 8.

TAR FORMING TEMPERATURES OF AMERICAN COALS.

OTTO CARTER BERRY, Assistant Professor of Experimental Engineering.

The material presented in this paper is the result of a series of investigations started in the laboratories of the University of Wisconsin and later continued at Purdue University. In form the paper is a brief of a part of a bulletin of the University of Wisconsin published by the author in 1914, with the addition of the results of the subsequent work.

The nature of the volatile matter in bituminous coal is attracting considerable attention at the present time. This is due not only to the enormous amount of coal annually used in the United States, but also to the important part volatile matter plays in determining how the coal must be handled in order to obtain the best results.

One of the most important and troublesome constituents of the volatile matter is tar, especially when the coal must be used in boiler furnaces or in power gas producers.

The investigations to be discussed in this paper had in view: (1) the determination of the temperature limits between which tars are distilled from the various classes of coal; (2) the temperature limits of the maximum rate of evolution of tars; (3) the relative quantities of tars distilled from the various general classes of coal; and (4) the lowest temperature at which one may be certain that the last trace of tar has been driven off from the coal.

Briefly stated, the results show that the temperature at which the first trace of tar appears will range from about 200°C. to about 385°C. usually falling quite near 360°. The maximum deposit will start at a temperature varying between 330° and 450°C. and will end between 430° and 530°C. The last trace of tar will appear between 530° and 680°C. The amount of tar produced seems to vary not so much with the amount of volatile matter in the coal as with the ratio of the carbon to the hydrogen as shown by the ultimate analysis of the coal.

When fresh coal is supplied to a furnace the volatile matter commences to distill off and if properly mixed with air and burned there is no

heat loss. The tarry products do not give trouble in gas producers when the gas is burned hot in ovens or furnaces. If however, the gas is allowed to cool, these products condense and stop up the piping, and unless removed, will clog up the engine valves if used for power purposes. The removal of these tarry products not only involves special and expensive apparatus and the expenditure of power, but also results in a loss of the available heat from the gas.

The problem as here presented is the outgrowth of an attempt to adapt the suction gas producer to the use of bituminous fuel. The type of producer used is what is known as the re-circulating producer such as is represented by the Whitfield and Pintsch patents. In this type of producer an attempt is made to draw off the tarry vapors from the top of the fuel column and introduce them again into the fire at the very bottom of the producer. The finished gas is drawn from the central portion of the fuel column. This location must be chosen with at least three points in mind:

- (1) It must be far enough down in the fuel column to be below the point at which the last trace of tar is driven off from the coal.

- (2) It must not be any farther down than is necessary, or the loss due to the sensible heat in the gas will be excessive. This loss, in percentage of the total heat value of the coal, will equal approximately the number of hundreds of degrees F. at which the gas leaves the producer. Thus, if the gas leaves at 1,200°F., the loss will approximate 12 per cent. of the total heat value of the coal burned.

- (3) The point of exit must be high enough above the bottom to allow ample opportunity for the CO_2 and H_2O resulting from the combustion of the distilled volatile matter to be reduced to free H_2 and CO .

To fulfill these several requirements, it is necessary to know exactly when each factor is operative. The depth of the incandescent zone that is necessary for a producer of a given size and capacity, and the precautions that are necessary to prevent a concentration of draft at any part of the producer, are fairly well known from practice. The most important item that is left for investigation is therefore to ascertain the exact temperature at which the last trace of tar is driven off from the coal.

An attempt was made to follow the temperature conditions met with in the gas producer, in these laboratory tests. This made it necessary to place the following list of requirements on the laboratory apparatus:

- (1) The coal must be heated very slowly and at a uniform rate.
- (2) The heat must be conducted from the outside to the center of

the body of coal by some good conductor, as the coal itself is a very poor conductor of heat and all particles in the body of coal must always be at a uniform temperature during the heating.

(3) The temperature of the coal must be accurately known at all times.

(4) The gases driven off from the coal must be swept out as soon as formed.

(5) The gases must be cooled down and continuously tested for tar.

(6) Any tar deposited in the pipes at a low temperature must not be allowed to re-distill at a higher temperature and then appear in the gas.

1. *The Furnace.* After considering the various possible means of heating the coal it was decided to use an electric resistance furnace. By this means the coal could be heated at any rate desired and the rate of heating could be controlled at all times, or the coal held at any desired temperature for any length of time. The furnace used is shown in cross section in Fig. 1 of the accompanying drawing, and was made as follows: A cylinder, (1), $3\frac{1}{2}$ " inside diameter and 20" long was made out of 16 B. & S. gage sheet iron, riveted at the seam. Around this was wound four thicknesses of wet asbestos paper (2). About 75 feet of 12 B. & S. gage nichrome resistance wire (3), was wound around this, four turns to the inch. The asbestos paper insulated the wire from the cylinder. Around this was wound more wet asbestos paper, to hold the wire in place, and the whole was covered by a layer of asbestos pipe covering (7), about 2" thick. The ends of the resistance wire were fastened to electric terminals on the furnace. The current passed through this furnace was taken from a 110 volt alternating current circuit, and was varied by means of a small water rheostat. The amount of current was measured by an ammeter in the circuit, as is shown in the sketch of the apparatus.

Six or seven amperes were required to bring the temperature up from 20° C. to 600° C. in four or five hours. Direct current would have been somewhat preferable had it been available, but the alternating current used did not jar the coil sufficiently to do any damage.

2. *The Coal Cartridge.* The cartridge in which the coal was placed to be heated was about $2\frac{1}{2}$ " inside diameter by 6" long. It was made up as shown in Fig. 1. (4) is a $2\frac{1}{2}$ " short nipple, having $2\frac{1}{2}$ " couplings (5) screwed on to both ends. Into these couplings pipe plugs (6) were screwed, thus forming a closed cartridge. One plug was drilled and




FIG. 1.

tapped for the $\frac{1}{2}$ " pipe (12), through which the gas from the coal could pass out. The other was drilled and tapped for a thermometer well (8) and $\frac{1}{4}$ " pipe (11). The body of coal heated each time was therefore $2\frac{1}{2}$ " in diameter, and it was necessary to have all the particles of coal in this mass at the same temperature. As coal is a very poor conductor of heat, it was decided to place iron disks (10) $\frac{1}{4}$ " apart throughout the entire length of the cartridge. These disks were large enough to touch the iron cartridge all around, thus taking on its temperature, and were drilled full of small holes to allow the gas to pass through them. As they were $\frac{1}{4}$ " apart, the heat had to be conducted through only $\frac{1}{8}$ " of coal. This fact, together with the very slow rate of heating employed, led us to expect the temperature throughout to be the same, within very close limits. The temperature was read at the very center of the coal body, by means of a thermo couple, the end of which extended down to and touched the end of the thermometer well (8).

3. *Means of measuring the temperature.* The thermo couples used were made of iron and nichrome wires welded together in an electric arc. The couples were used with a Brown millivolt meter, with a resistance of 85 ohms. It was carefully calibrated and checked at the time the experiments were completed. The couples were correct to within 10° C. throughout the range of temperatures here reported. The couples were left in place throughout each entire test, and the temperature readings made whenever desired.

4. *To keep the gases swept out as formed.* To sweep the gases out as they were formed gas free from tar was forced into the cartridge under pressure, through the $\frac{1}{4}$ " pipe (11), and allowed to pass out through (12) in a constant stream. The pressure of this gas inside the cartridge was measured by a mercury manometer, and was kept at about $2\frac{1}{2}$ " of mercury. This gas could not be allowed to contain any O_2 , as it might then burn the coal or tar vapors at the higher temperatures, so air with the O_2 burned out was used. The arrangement of the apparatus as used is shown in Fig. 2. The air was burned in a small furnace made of a piece of 6" pipe about 2' 6" long. This pipe had grates at the bottom and a coupling and plug at the top. The coupling was drilled and tapped $\frac{1}{2}$ " pipe size on one side, and connected to a large coil of $\frac{1}{2}$ " pipe which rested in a tank of cold water. The small furnace thus constructed was filled with an anthracite fire and the plug at the top put in. The air pump then pumped air through the furnace and cooling coil and compressed

it into the large air tank, where it was stored for use. By this means the supply of gas for a complete test could be stored up before the test itself was started. One more precaution had to be taken in using this gas, as it could not be allowed to affect the temperature of the coal as it passed through. To prevent this, a coil of pipe was placed over a gas flame and the gas passed through and heated up to the temperature of the coal, before it was allowed to enter. The temperature of the gas was measured by a thermo couple which extended into it through a tee in the pipe line. To make assurance doubly sure the end of the cartridge itself was filled with steel chips, as is shown at 22 in the sketch in Fig. I. The entering gas was thus forced to pass through a considerable volume of these chips before coming into contact with the coal. It was found difficult to heat the gas up to the highest temperature of the producer. This might tend to affect the seeming temperature at which the last traces of tar appear. The tendency of the gas would always be to be lower than that of the coal. For this reason the end of the thermocouple was placed in the coal at the end where the gas enters it, and therefore at its cooler end, in case there should be any difference at all. Thus the temperature reported as the one at which the last trace of tar appears is as accurate as it is possible to make it.


5. *To test the gases for tar.* The next problem was to find a means of subjecting the gases from the coal to a continuous test for tar. The most searching and satisfactory test known to the author, and the one used by the gas companies over the country, is to allow a small stream of the gas to strike a piece of white paper at a high velocity. If there is any trace of the tar at all in the gas, it soon leaves a spot on the paper. This test was adopted. To use it, the gas must be cooled down before it strikes the paper. This was accomplished by keeping a cloth filled with cold water constantly lying on pipes (12) and (18). It was desired to have the test continuous. The device used to accomplish this is shown in Figs. 1 and 2. The rollers (18) are about 10" in diameter and are supported by steel rods through their centers which turn freely in iron supports at either side. On one end of each of these steel rods was placed a small wooden spool, around which was wound a cord, supporting weights W_1 or W_2 . A long strip of cloth was wound around one of these rollers and its end started around the other. A piece of paper ribbon was wound on with the cloth. W_1 and W_2 tend to turn the rollers in opposite directions, thus keeping the cloth and paper strip tight. W_1 is enough heavier than

W₂ to cause both to turn, unless held back in some way. The key of an alarm clock (21) was fastened to a train of gears (20), and these in turn to the stem of roller (18), so that the rate of motion was held back to a speed governed by the running of the clock. In this case the speed was 30" an hour. The rollers were set in such a way as to cause the paper to pass about $\frac{1}{4}$ " under the end of the orifice (16) in pipe (15). The strip of paper was marked at the beginning and end of the test, so that the exact time when any point on the paper was under the orifice could be determined. Thus a continuous record was kept of the amount of tar in the gas.

It might be well to state here that all of the tar is not taken out of the gas by this means. The tar particles in the cooled gas are very fine and light, and many of them are cushioned off from the paper, and never touch it to stick. Consequently the tar deposited on the paper does not represent all of the tar content of the coal. On the other hand, the slightest trace of tar in the gas will quickly black the paper, and the deposit is probably very nearly proportional to the entire tar content of the gas.

6. *To prevent re-distillation of tar once deposited in exit pipe.* The remaining problem was that of making sure that no tar could be deposited in pipe (15) at a low temperature and be later re-distilled, to show up on the paper at too high a temperature. To accomplish this a large number of duplicate pipes were made and carefully fitted into the threaded bushing (14). By changing these pipes (15) every five or ten minutes, the effect of such a tendency was quite completely eliminated.

The coals tested were chosen to represent the different American grades. They were first ground and screened over a mesh of 20 wires per inch and through one of 10 wires per inch. About 315 grams of coal will fill the cartridge, and it was put in in layers $\frac{1}{4}$ " thick between iron disks. The cartridge was then placed in the middle of the furnace, the thermo couple put in place, and the asbestos packed in against both ends, cutting off the radiation here, and causing the ends and middle, all to keep at the same temperature. A current of 6 amperes was then passed through the resistance wire of the furnace. While the latter was coming up to the temperature where the light oils start to come off, the thermo couples were connected up, the flame placed under the gas pre-heating coil, the gas from the storage tank turned on, the paper rolls connected up to the clock and the clock started. The pipes (15) were cleaned and prepared for immediate use, and the data sheet prepared. As soon as




the thermo couple indicated a temperature close to that at which there was prospect of an oily deposit, the paper rolls were put in place, the paper marked and the test started. When the tar commenced to show up the temperature of the furnace was recorded and the pipe (15) changed every 10 minutes. As the temperature of the furnace rose to 300° C. the current was increased $\frac{1}{4}$ of an ampere, at 400° C $\frac{1}{2}$ of an ampere more, and at 500° C. $\frac{3}{4}$ of an ampere. This was done to take care of the increased radiation, and to keep the rise in temperature at a constant rate. When the paper ceased to show any signs of a tar deposit it was again marked and timed and the current shut off. The strip of paper was then cut up into lengths corresponding to 10 minute periods, and carefully weighed. As the weight of the paper per inch was very constant, the excess in weight over that of clean paper was in each case due to the tar. From this two curves could be drawn, with time plotted horizontally, while one had temperature centigrade and the other grams of tar plotted vertically. These curves when placed one right over the other, as here given in Fig 3, indicate the amount of tar coming off at each temperature. The points where the tar starts and stops can not be indicated by this curve, as the ends of the deposit are too thin to have appreciable weight. They are consequently separately noted elsewhere.

The first condensable gas to be driven off from the coal and appear on the paper record is water vapor. After the last of the water has disappeared there is quite a temperature range through which there is no deposit at all. Then the paper will begin to show a slight trace of oil. This will gradually increase in amount and give the paper the appearance of having been parafined. The deposit will then gradually assume a brownish color, as though engine oil were appearing. Later a temperature will be reached at which the deposit will increase very rapidly in amount, and will assume a distinctly tarlike appearance. The first tar to be deposited is usually very soft and sticky at room temperature. As the temperature rises the tar becomes steadily stiffer, until it is finally hard and brittle when cooled. The temperature range through which the maximum deposit occurs will vary from about 100° C., for some western coals, to 175° for some of the samples from the east. At the higher limit of this range the deposit becomes rapidly smaller in amount until it is too small to weigh, but the paper is still distinctly browned. This discoloration becomes less and less plain, until it finally disappears entirely. There is no definite temperature at which the first and the last

trace of the tar appears, in the same sense that water has a boiling temperature. The first deposit is so indistinct that it is almost impossible to tell whether there is a deposit or not. The increase is also so very gradual that it is difficult to choose the temperature at which to report the first appearance of a deposit. This gradual increase will extend over a temperature range of from 50° to 150° C. before there will be a sufficient deposit of tar to feel sticky to the finger. The determination of the temperatures between which the maximum deposit occurs is likewise an arbitrary matter, and also the temperature at which the last trace of tar appears. Therefore the results as here reported must not be too literally interpreted. However they are a very careful estimate of the facts as they are and the highest temperature reported is one at which one may feel assured that the very last trace of tar has disappeared from the coal.

CARTERVILLE ILLINOIS COAL.



SHAWNEE MOUND, TIPPECANOE COUNTY, AS A GLACIAL ALLUVIAL CONE.

WM. A. MCBETH.

There stands at the northwest corner of section twenty-three (23), town. twenty-one (21) north, range six (6) west, a locally well known hill, quite unusually large and prominent for that part of the country, which generally is a moderately undulating, and, over extensive areas, a quite monotonous plain.

The area of the hill is about thirty-five (35) acres. Its height at the apex is seventy-five (75) feet above the steps of the front door of the residence (facing the road) located near the southwest edge of the hill on the general level of the country. A creek channel at the northwest edge is eighty (80) feet or more below the summit. The long axis of the hill is east west, in which direction it is nearly one hundred (100) rods long, with varying cross-distances averaging a little more than half its length. The high part is near the east end, where the steepest slopes occur. A small basin lies at the foot of this east slope. The distance of the highest point is almost one-third ($\frac{1}{3}$) of the length of the pile from the east end, whence the slopes are much gentler to the western edge. The outline and form of this feature may be properly described as lobate, four leaves, including the west end, showing on the south side, and three on the north side not symmetrical with or opposite those on the south side.

In structure and material the hill is composed of sand, gravel, silt, clay lumps and a very few boulders. The rock fragments are igneous, or crystalline material in great variety, rounded and polished after the manner of stream-worn waste. It is very irregularly bedded with a generally abrupt pitch to the west, or in the direction of the long slope of the hill. The pebbles are not in many cases larger than apples or baseballs, very few exceeding four or five inches in any diameter. Beds of fine sand,



Shawnee Mound. Excavation in North Side showing Material and Arrangement.



Shawnee Mound. (From the South.)


or even silt, are interstratified with layers containing the coarsest materials.

The best, probably the only, interpretation of the observed facts is, that this is a great alluvial cone built in or at the edge of the retreating ice sheet when its edge lay at the prominent Independence-Darlington Moraine, in the range of which this feature stands, although no other of its strong elevations appear in the immediately surrounding landscape, and not within two or three miles.


This great pile, for it is really quite impressive, represents the deposit of a stream of considerable volume flowing off or out of the ice at the time when the height or depth of the ice at this southwestern edge must have been not less than one hundred (100) and possibly several hundred feet. This stream flowed in a channel deep enough to confine it for a long time to this particular place. Possibly the channel was a deep canyon in the ice or a tunnel under it. As it heaped up the material at one point its course was diverted and a new direction of flow and construction was begun in true cone or delta building fashion.

An interesting question arises as to a possible relation between this hill and the gigantic South Raub Esker lying a few miles to the east northeast. The trend of this Esker is directly toward Shawnee Mound and the direction of the esker stream was southwest, but there is a gap of nearly five (5) miles, a distance as great as the length of the esker itself, between the west end of the esker and the Shawnee Mound cone, and no sufficient intermediate correlating features have as yet been found.


This discussion is presented to show the importance of detail work in interpreting the complex and little known phenomena of the great ice sheet.




Shawnee Mound.
Contours Generalized.





Shawnee Mound, S. Raub Esker & Drainage.



Sunman





CORRELATION OF THE OUTCROP AT SPADES, INDIANA.

 H. N. CORYELL.

INTRODUCTION.

The collections forming the basis for the present paper were carefully made by the Rev. T. A. Bendrat, M. S., during the autumn of 1913, from the exposure one mile west of Spades, Indiana. His careful description of the outcrop has been of considerable value in the correlation.

The identifications of the species were made in the laboratories of the Department of Geology of Indiana University under the supervision and direction of Professor E. R. Cumings and Dr. J. J. Galloway. Their assistance and suggestions have been of great value.

THE PRESENT DIVISIONS OF THE RICHMOND.

Richmond—

- Elkhorn (*Platystrophia moritura* zone).
- Whitewater (*Homotrypa wortheni* zone).
- Saluda (*Tetradium minus* zone).
- Liberty (*Strophomena planumbona* zone).
- Waynesville (*Dalmanella meeki* zone).

DESCRIPTION AND STRATIGRAPHY.

The reconstruction of the Cincinnati Division of the C., C., C. & St. L. Railroad through Ripley County, Indiana, in 1902-04 made available for study a section of strata high in the Cincinnati one mile west of Spades. The exposure at present is from five to six feet high and in places partially covered with debris, leaving, however, distinct ledges outcropping. The strata are approximately horizontal. The limestone at the base of the exposure weathers to a bluish-gray; it is very variable in texture, some portions being very finely crystalline, others containing quite large calcite crystals, as well as geode formations and iron concretions. Above this limestone the strata become more argillaceous, thinly bedded and present a distinct shaly structure.

This section possesses a distinct similarity to the upper part of the Whitewater section exposed at Richmond, Indiana, in being a very nodular, shaly limestone.

Though similarity in lithology is not a conclusive proof that two or more separated sections are parts of the same formation, yet the fact that there is a distinct resemblance does tend to lead favor to that decision.

A detailed description of this locality is given in the following section, taken from "The Stratigraphy and Paleontology of the Cincinnati Series of Indiana."*

SECTION ALONG THE WEST FORK OF WHITEWATER RIVER AT
RICHMOND, INDIANA.

(Number 1 is at the top of this section.)

1. Exposures in the bank above Thistlewaite Falls on the west fork of Whitewater River, about one and one-quarter miles north of the National road bridge across Whitewater River. Thin, lumpy limestone. *Rhynchotrema dentata* (aa). Several species of gastropode, including *Salpingostoma richmondense* (c). *Strophomena sulcata* (r)..... 8 feet.
2. Layer in the breast of the falls. Heavier layer at the top. Limestone. *Monticulipora epidermata* (c). *Platystrophia acutilirata senex* (c). *Homotrypa wortheni*..... 5 feet
3. West side of creek just below the falls. Bryozoa (aaa), *Monticulipora epidermata*, etc..... 5 feet.
4. Just north of the C. R. & M. R. R. bridge. Thin, shaly limestone, *Rhynchotrema capax*, the highest specimens. *Plectambonites sericeus* (rr)..... 5 feet.
5. Just south of the C. R. & M. R. R. bridge. *Rhynchotrema capax* (aa) 4 ft. 8 in.
6. About one-eighth of a mile north of the road bridge across the west fork. *Ptylodictya plumaria*, etc..... 5 feet.
7. A short distance north of the junction of the east and west forks of the river. Limestone and intercalated shale. *Hallopora*, very similar to *H. rugosa*. No specimens of *Rhynchotrema dentata* 4 feet.

*Prof. E. R. Cumings. Indiana Dept. Geology and Nat. Resources, 32d Ann. Rept. 1907.

RELATION OF THE SPADES SECTION TO CUT 18 NEAR WEISBURG.

In all parts of the Tanner's Creek section there is a general dip of five feet to the mile, and we may assume that this dip holds over the five miles from Weisburg to Spades. This places the base of the Whitewater, which is exposed in the cut 18 just north of the station at Weisburg, seventy-five feet below the railway grade at Spades. In the Richmond section of the Whitewater, where the whole of the formation is exposed, a similar thickness is found, leading us to conclude that the outcrop one mile west of Spades represents the upper strata of the Whitewater. The position of the latter outcrop in reference to the Whitewater in the Ripley County locality is shown on the accompanying chart. The lines of dip have been extended to Batesville, five and one-half miles northwest of Spades, where there is sixty feet of Whitewater, portions of which are exposed in the stream gullies a few miles south of Batesville.

RANGE OF THE SPECIES FOUND AT SPADES, INDIANA.*

- Byssonychia grandis* Ulrich, Upper Richmond.
Byssonychia obesa Ulrich, Whitewater.
Calymene callicephalo Green. Common throughout the Cincinnati, especially at the top of the Waynesville.
Dalmanella meeki (Miller). Coryville-Arnheim and Waynesville.
Platystrophia laticosta (Meek). McMicken, Maysville and Richmond.
Lophospira trophidophora (Meek). Whitewater.
Platystrophia acutillirata (Conrad), Waynesville, Liberty and Whitewater.
Platystrophia acutillirata senex (Cumings). Upper Whitewater.
Platystrophia laticosta (Meek). McMicken, Maysville and Richmond.
Protarea richmondensis (Foerste). Waynesville, Liberty and Whitewater.
Rafinesquina alternata (Emmons). Throughout the Cincinnati.
Streptelasma rusticum (Billings). Waynesville, Liberty, Saluda and Whitewater.
Arthropora shafferi (Meek). Throughout the Cincinnati.
Bythopora delicatula (Nicholson). Coryville-Arnheim and Richmond.
Dicranopora emacerata (Nicholson). Maysville and Richmond.
Hallopora cf. ramosa. Richmond.
Helopora elegans Ulrich. Liberty and Whitewater.

*See "Stratigraphy and Paleontology of the Tanner's Creek Section of the Cincinnati Series of Indiana."

Heterotrypa sp. undescribed. Upper Whitewater.
 Homotrypa wortheni (James). Richmond and Whitewater.
 Homotrypella hospitalis (Nicholson). Waynesville to Whitewater.
 Nematopora cf. lineata (Billings). Middle and Upper Ordovician.
 Rhinidictya lata (Ulrich). Waynesville.
 Rhombotrypa sp. undescribed. Upper Whitewater.
 Stigmatella sp.

SPECIES FROM NUMBERS ONE AND TWO OF THE WHITEWATER SECTION AT
 RICHMOND, IND. (See section above.)*

Anomalodonta gigantis Miller.
 Byssonychia grandis Ulrich.
 *Byssonychia obesa Ulrich.
 Byssonychia suberecta Ulrich.
 Ischyrodonta modioliformis Ulrich.
 Pterinea demissa (Conrad).
 *Protarea richmondensis Foerste.
 *Streptelasma rusticum (Billings).
 Cyclonema bilix (Conrad).
 *Lophospira tropidophora (Meek).
 Salpingostoma richmondensis Ulrich.
 Cytoceras sp.
 Amplexopora sp.
 *Bythopora delicatula (Nicholson).
 Bythopora meeki (James).
 Ceramoporella ohioensis (Nicholson).
 Corynotrypa inflata (Hall).
 *Dicranopora emacerata (Nicholson).
 Hallopora subnodosa (Ulrich).
 Heterotrypa prolifica Ulrich.
 Heterotrypa subramosa (Ulrich).
 Homotrypa austini Bassler.
 Homotrypa flabellaris Ulrich.
 Homotrypa flabellaris spinifera Bassler.
 *Homotrypa wortheni (James).
 Monticulipora epidermata Ulrich and Bassler.
 Peronopora pavonia (d'Orbigny).

*The species marked with an asterisk are found in the section at Spades, Indiana.

Homotrypella rustica Ulrich.
Rhombotrypa crassimuralis (Ulrich).
Dinorthis subquadrata (Hall).
Hebertella occidentalis (Hall).
 **Platystrophia acutilirata* (Conrad).
 **Platystrophia acutilirata senex* Cumings.
 **Rafinesquina alternata* (Emmons).
 Rhynchotrema denatatum (Hall).
Strophomena vetusta James.
Strophomena planumbona (Hall).
Strophomena sulcata (Verneuil).
Zygospira modesta (Hall).

In comparing the fauna of this cut with the numbers, one and two, of the Richmond type section, we find that many species that appear in the type locality are not found at Spades, but those that do appear at Spades are undoubted Whitewater species. Those that are limited to and characteristic of, the Whitewater, are: *Homotrypa wortheni* (James?) *Platystrophia acutilirata senex*, Cumings. *Byssonychia obesa* Ulrich, and *Lophospira trophidopora* (Meek).

Twenty-four species were identified from Spades, three of which are new and undescribed. The description and discussion of their morphological structure will be given in a subsequent paper.



THE PALEOBOTANY OF THE BLOOMINGTON, INDIANA, QUADRANGLE.

T. F. JACKSON.

The fossil plants herein discussed are, with three exceptions, Pennsylvania forms and were collected principally from two localities in the Bloomington, Indiana, Quadrangle. The greater part of them were obtained from a shale bed about one-fourth mile southeast of the Yoho School. This bed was made up of a succession of thin, bluish-gray clay-shales interstratified with thin sandy layers, with nodules of iron ore irregularly distributed throughout the entire bed. The shale layers were very soft and plastic when wet and both the shale layers and sandy layers were rather hard and very brittle when dry. One of the shale layers was very highly impregnated with iron oxides, and from this layer the best fossils were obtained. The entire bed attains a thickness of eight to nine feet.

The remainder of the Pennsylvanian forms were obtained from a thin, ferruginous sandstone layer and an overlying sandstone layer, about one-fourth mile southeast of Cincinnati. Molds and casts of *Lepidodendron* and *Calamite* forms were collected from the latter. The ferruginous sandstone layer contained a number of *Trigonocarpon* and a few *Carpolithes* forms.

Loose sandstone fragments of fossil plants, apparently of Pennsylvanian age, were noted in a number of places in the southwestern part of the Quadrangle, but, as their exact horizon could not be ascertained, those forms are not included in the following lists of species.

A few fragments of Mississippian forms were noted in the central and northern part of the west half of the Quadrangle. Those plants were very poorly preserved and at but one place were fossils obtained in a state of preservation such that identification was possible. Three species in a fair state of preservation were found in a sandstone layer a few feet above the Mitchell limestone, about one-half mile west of Whitehall.

Although a few of the Pennsylvanian plants examined represent new species and several others differ more or less from previously described forms it is not thought to be advisable to figure and describe those new forms at this time inasmuch as it is planned to include them in a later paper on the flora of the entire Pottsville section as represented in the State.

List of plants from the Yoho School locality :

- Sphenophyllum cuneifolium* (Stb.) Zeill.
- Sphenophyllum bifurcatum* Lx.
- Sphenophyllum tenerrimum*? Ett.
- Asterophyllites erectifolius* And.
- Asterophyllites gracilis* Lx.
- Calamostachys* sp.
- Lepidodendron clypeatum* Lx.
- Lepidodendron* sp.
- Alethopteris Evansii* Lx.
- Alethopteris grandifolia* Newb.
- Alethopteris lonchitica* (Schloth.) Brongn.
- Alethopteris* sp.
- Pecopteris plumosa*? Brongn.
- Pecopteris* sp.
- Neuropteris Elrodi* Lx.
- Neuropteris Jenneyi*? D. W.
- Neuropteris* sp.
- Neuropteris fimbriata* Lx.
- Neuodontopteris*? sp.
- Pseudocopteris decipiens* Lx.
- Sphenopteris* sp.
- Sphenopteris communis* Lx.
- Zeilleria* sp.
- Cardiocarpon annulatum* Newb.
- Cardiocarpon cornutum* Dn.
- Cardiocarpon pachytesta*? Lx.
- Cardiocarpon* sp.
- Rhabdocarpus* sp.
- Carpolithes* sp.

A mere casual inspection of the above list will show the Pottsville aspect of the plants. Of these forms *Sphenophyllum cuneifolium*, *S. bifurcatum*, *S. tenerrimum?*, *Asterophyllites erectifolius*, *A. gracilis*, *Alethopteris Evansi*, *A. grandifolia*, *A. lonchitica*, *Neuropteris Elrodi*, *N. fimbriata*, *N. Jenneyi*, *Pseudocopteris decipiens*, *Sheuopteris communis*, *Cardiocarpon annulatum*, *C. cornutum*, *C. pachytesta?*, have not been reported, I believe, from formations younger than Pottsville age. Of those Pottsville forms *Sphenophyllum bifurcatum*, *Alethopteris Evansi*, *A. lonchitica*, *Neuropteris Elrodi*, *N. fimbriata*, and *Cardiocarpon cornutum* were reported by White* as being confined to the Upper Lykens Coal group of the Pottsville type section. *Sphenophyllum cuneifolium* and *Cardiocarpon annulatum* were reported from both the Upper Lykens Coal group and the Upper Intermediate group of the type section. *Alethopteris grandifolia* was reported from the Lower Intermediate group in the type section. The vertical range of several of the remaining forms is either not definitely known or is too great for correlation purposes; the rest of the forms are too poorly preserved for specific classification. Therefore only those forms reported from the Pottsville type sections are considered of value for correlation. A comparison of the list of Pottsville plants from the Yoho School locality with the list of plants from the Pottsville type section indicates that the Pennsylvanian of the Yoho School area represents an horizon in the upper part of the Middle Pottsville of the type section.

List of plants from the Cincinnati locality:

Calamites approximatus Schloth.

Calamites sp.

Lepidodendron clypeatum Lx.

Lepidodendron sp.

Neuropteris lunata? D. W.


Lepidophloios sp.

Of the above listed forms *Calamites approximatus* and *Neuropteris lunata?* have been reported from the Upper Lykens Coal group of the type section of the Pottsville. *Lepidodendron clypeatum* has too great a vertical range to be of value as a horizon marker. The other forms were too

*White, David. The Stratigraphic Succession of the Fossil Floras of the Pottsville Formation in the Southern Anthracite Coal Field, Pennsylvania. 20th Ann. Rep't. U. S. G. S. Part II, 1900.

poorly preserved for accurate classification. Considering only the forms first mentioned it would appear that the Pennsylvanian of the Cincinnati locality would fall within the Middle Pottsville of the type section.


Three Mississippian fossil plants, *Lepidodendron Volkmannianum* St., *Lepidodendron* sp., and a variety of *Stigmaria ficoides* were obtained from the Chester sandstone a few feet above the Mitchell limestone, about one-half mile west of Whitehall.



VICINITY
 ONTARIO
 MALOTT

GLACIER
 CHANNELS
 WOODS BASIN

DATA IS TAKEN








MAP OF FLATWOODS AND VICINITY

MONROE AND OWEN COUNTIES

DRAWN BY CLYDE A. MALOTT

1918

-  ICE LIMIT OF ILLINOIAN GLACIER
-  PRE-GLACIAL STREAM CHANNELS
-  OUTLINE OF FLATWOODS BASIN
-  PUBLIC ROADS
-  WELLS FROM WHICH DATA IS TAKEN

THE FLATWOODS REGION OF OWEN AND MONROE COUNTIES, INDIANA.

CLYDE A. MALOTT.

EXTENT AND TOPOGRAPHY.

Lying between Ellettsville, Monroe County, and Spencer, Owen County, Ind., is a strip of territory some six miles long and averaging about two miles wide, which has been the object of considerable curiosity and study. It is a low level basin nearly surrounded by higher land, yet having several openings in the surrounding periphery of hills.

The surface of the region is mainly an ash-colored soil of a fine texture, containing very little sand. It is in reality a silt region at the surface, and its outline is clearly discernible at the margin of the basin. This silt region, or its outline, is the principal means of determining the margin of the region, as indicated by the map. It coincides for the most part with the foot of the hills surrounding the basin. Here and there in the basin a hill rises out of the silt region somewhat as an island out of the water, and frequently a hill-like peninsula protrudes out into the region, rising high above the ash-colored silt margin.

The silt margin lying about the foot of the hills is rather uniform in height, averaging close to the 760-foot contour line, excepting at the northeast margin, where it extends much higher. This region of the Flatwoods area is also exceptional in regard to the periphery. Elsewhere, except at the openings, the hills surrounding the area rise rather suddenly above the basin; but here the margin is scarcely discernible, as the slope is very gradual and seems rather to fade out instead of being abrupt. This phenomenon is one of considerable importance and will be discussed later.

Lying in the long axis of the region is McCormicks Creek. This stream drains about nine-tenths of Flatwoods. Its head is about seven hundred fifty feet above sea level, one and one-half miles west of Ellettsville. The first few miles of its course is over the flat plain of the basin, which gives it but little fall. After leaving Section 36, T. 10 N. R. 3 W., the

fall increases slightly, and soon after entering Section 26, T. 10 N., R. 3 W., the stream has cut down to bed rock. The road leading east and west along the north side of Section 26 is practically the margin of Flatwoods in this vicinity. On this road a shallow ford crosses the creek on the 700-foot contour line, over a solid rock floor. From the source of the stream to this ford, a distance of about five miles, the stream has a fall of about fifty feet. It enters White River about two miles below, at an elevation of 540 feet above sea level. Thus the last two miles of the stream have a fall of 160 feet. Practically the entire last two miles of the course is over a solid rock bed. The region presents some of the most rugged and beautiful scenery in the State. The stream courses down a veritable gorge which is but little wider than the stream itself. Many cascades occur, and about a mile below the ford a fall of about 12 feet occurs. In low water the stream cascades over this fall, but when the water is high it rushes over with a roar that can be heard for some distance. Above the falls the floor of the gorge is swept clean of debris, but below, the gorge is wider, and in many places is choked with the rock debris that has been carried from above or has fallen from the almost vertical walls on either side.

Just east of the source of McCormicks Creek is the source of a small branch which leads northeast through an opening in the rim of the basin and empties into Jacks Defeat Creek. This stream drains but little of Flatwoods, as indicated by the map. Its source is about the same height as the McCormicks Creek source, and its mouth, one and one-third miles northeast, comes out at about 670 feet above sea level, thus giving it a fall of eighty feet.

Another break in the rim of the basin occurs in Section 30, T. 10 N., R. 2. W., about two miles southwest of Stinesville. This opening is narrow and its surface is below the 760-foot contour line. To the north of the narrow opening is a wide flat plain similar to the Flatwoods, having a silt surface of the same nature. This flat is drained into Big Creek. The narrow opening itself is practically bed-rock at the surface. Sinks occur in it. A small portion of Flatwoods is drained into a deep sink near the northwest part of this section. The water that goes into this sink undoubtedly passes under the narrow opening and comes out into Big Creek below, as several springs occur in the upper part of this creek. Just to the west of this narrow opening is a high hill capped with sandstone, which is at least sixty feet higher than the opening. The silt line

can be distinctly seen on practically all sides of this hill, coming slightly above the 760-foot contour line.

About one and one-half miles southwest of Ellettsville in the south-east corner of Section 8, T. 9 N., R. 2 W., is a sink which has a small stream entering it, and draining about one-half square mile of Flatwoods. This stream has lowered this corner of Flatwoods considerably below the general level. The water that goes into the sink flows out about a half mile to the southwest from a couple of large springs which drain into Raccoon Creek.

Perhaps the most interesting opening in the periphery of the basin occurs in Section 1, T. 9 N., R. 3 W. This opening leads into a tributary of Raccoon Creek, and is at least a third of a mile wide. To the east of it is a high hill or ridge attaining a maximum height of 910 feet, and on the west another ridge reaches above the 880-contour line. The floor of the opening itself is twenty-five feet or more below the silt-line on the sides of the hills. This opening is really a connection between Flatwoods proper and a continuation of it in the Raccoon Creek Valley. Consideration will be given it later.

There is yet another outlet to the Flatwoods region, which at first was very puzzling to the writer. At the western extremity of the basin Allistons Branch reaches into it by many deep and narrow tributaries. These tributaries are almost invariably headed by seepage springs which come out into the sandy material in which the tributaries are cut. The basin itself is some higher at this western part. The basin cannot be said to have a margin at this western limit; it ends more or less abruptly in the tributaries of Allistons Branch. If it ever had a peripheral margin at this end it has been effaced by the V-shaped valleys leading into Allistons Branch. The writer intends to prove that this western end never had a distinct margin, that is, like the so clearly identified ones on the southern and northern periphery of the region.

From the silt line at the foot of the hills, the slope of the basin is generally inward toward the mathematical center. The lowest part of the basin (not considering the valley and channel of McCormicks Creek) is along the Monroe-Owen County line, between sections 31 and 36, T. 10 N., and branching off from this along the southern part of section 31 and along the northern part of section 36. This region is very fertile, being almost entirely a black loamy soil. The white silt of the bordering

regions passes under this black soil. But farther out into the black soil, the silt underneath almost pinches out. The low region containing the black soil was undoubtedly the centre of the basin in former times, even as it is now. This low-lying, fertile region is very near the 720-foot contour line; thus it is some forty feet below the silt line at the foot of the hills surrounding.

It was said that the slope of the basin is generally toward the mathematical center; this is not true specifically, as there are some exceptions. Several places considerably elevated occur. The large one in Section 31, T. 10 N., R. 2 W. reaches to the height of 795 feet, approximating the peripheral regions. A well shows that bed rock is near the surface of this old monadnock. In section 36, west of the above, a long arm-like island projects out into the basin, and near the south of the middle of the section a notch occurs in the arm, which almost separates the north end, leaving a round-like knob projecting some forty feet above the basin. This elevation also has bed-rock in it near the surface. Section 25, T. 10 N., R. 3 W. has two elevations some twenty feet above the general level of the basin. It was not determined whether these had bed-rock near the surface, but indications are, especially in the western one, that it is there at a shallow depth. The elevation on the section line between sections 26 and 35, T. 10 N., R. 3 W. is a rounded knoll about twenty-five feet above the general level of the basin. Indications are that it contains no bed-rock. The northern part of Section 6, T. 10 N., R. 2 W., contains a slight elevation, perhaps twelve feet above the low-lying area adjacent. A deep well proves that it contains no bed-rock. Southeast in section 5, and entering section 8, is a long elevation parallel to the long axis of the basin and about twenty feet high. A well proves that this one also contains no bed-rock.

While dealing with the irregularities of the surface of the basin, attention must be called to the depression at the southwestern edge of Flatwoods, on the section line between sections 2 and 3, T. 9 N., R. 3 W. This depression, containing about two acres, is the site of a small lake which is being rapidly filled by in-wash and vegetation. The elevation of the surface of this small lake, bearing the name of Stogsdill Pond, is about 770 feet. It is enclosed on three sides by sloping banks which reach thirty feet above the water. It is open on the north.

The south bank of Stogsdill Pond is the lowest opening to a sort of an adjunct to the Flatwoods basin. The surface of this adjunct slopes

gradually from the bank of the pond for about one and one-fourth miles to the south, and in its lowest place, a series of sinks jutting against the bed-rock hills at the south, comes down to the 700-foot contour line. This adjunct basin is about as broad as long. It is drained mainly by two southward extending streams that come to the series of sinks at the south end. A third but much smaller southward flowing stream drains the western side. It also disappears in a sink in the southwest corner of the region. The western edge, near the middle of sections 3 and 10, ends abruptly in the rapidly headward-etching streams of the headwaters of McBrides Creek. The eastern rim is the ridge of highland which has been mentioned as the western rim of the opening extending into the Raccoon Creek Valley. This same ridge turns to the west and forms the southern rim of the adjunct basin also, and beneath which the waters of the region flow in their underground passage. This small adjunct basin undoubtedly once had a smooth and gentle slope from the southern rim of the western part of Flatwoods proper to the high ridge at the south, but subsequent drainage through sinks at the southern end has eroded it into three main grooves with many smaller tributary grooves. The slope began at the north at an elevation of more than 800 feet, and ended at the southeastern corner at 740 feet. The west part of the southern end was somewhat higher, perhaps 760 feet. It was lowest at the southeast corner, because at this place there is an opening in the bed-rock ridge, which leads to Raccoon Creek. This opening will be called into account later.

We are now ready to go back to the broad opening in the middle of the southern periphery of the basin, and see the extension of the Flatwoods basin to the south. As has been said before, the floor of this opening is slightly below 740 feet, and that the silt line extends as high as 760 feet. Soon after leaving the opening, the silt line on the side of the ridges becomes more or less indistinct, since erosion has either erased it or covered it over. Still farther south not only the silt line is removed, but much of the one-time basin-flat itself is removed. The flat, however, can be traced for four and one-half miles south and some west down the valley, or rather above the valley of Raccoon Creek. The creek here turns abruptly and flows to the northwest, at a right angle to the course above. Modified portions of the old flat are distinct for two or more miles northwest of the sharp turn.

It is understood, then, that this Raccoon Creek addition is very much eroded by the present stream and its tributaries. But it is important to

notice the elevation of the flat itself. Where it leaves Flatwoods proper, it is somewhat below the 740-contour. The slope is gradually downward from this place to the south. In the vicinity of Freeman the elevation of the flat is 700 feet. This makes a gentle slope down the valley southwest of a little less than ten feet to the mile. The Raccoon Creek addition extends up the creek almost as far east as the eastern margin of Flatwoods proper. The old flat is recognizable for three and one-half miles up Little Raccoon Creek, which enters Raccoon Creek from the southeast near Freeman. The extent and shape of the addition can be seen by consulting the map. It contains in all about eight square miles. Thus this addition and the adjunct south of Stogsdill pond make an area approximating that of Flatwoods proper.

While dealing with the Raccoon Creek addition of the Flatwoods basin, it must be emphasized that it occurs only in remnants. There are, however, quite large areas, sometimes a quarter-section or more, that have suffered little erosion. In such cases, or in cases where much smaller areas are preserved, there occurs the same flat, ash-colored, crawfish soil flat that is so characteristic of Flatwoods proper. Second to these flats, the most striking physiographic feature is the terraces resulting from the streams cutting down into the flat. The terraces begin almost immediately after entering the gap from Flatwoods proper. Here they begin at zero, but soon become quite a distinct feature. They grow higher very rapidly, so to speak, as the stream cuts down into the flat to the south. At Freeman, four and one-half miles below the gap, the stream has cut down one hundred feet below the old flat, and the terraces are accordingly one hundred feet above the stream. But in this vicinity there are many places where the terraces are indistinct, as they are so eroded that they no longer appear as terraces. This condition occurs in the immediate vicinity of Freeman. Beyond a slight bed-rock hill to the east of Freeman, however, the flat is distinctly discernible, and the terraces show beautifully above the small tributary streams that are etching their way into it.

UNDERGROUND INFERENCES AS REVEALED BY WELLS AND BORDERING REGIONS.

Having dealt somewhat with the extent and topography of the Flatwoods region, we shall now turn to a slightly different phase. Perhaps the most interesting particulars of the region are the underground inferences as they are revealed by the wells of the region and by the places along the western margin, which have suffered erosion by the rapidly

encroaching streams from the west. Data from shallow wells, while revealing interesting sub-surface particulars, are not sufficient to give the shape of the pre-existing basins of the region. The deep wells, the wells which reach bed-rock, are important in this respect. There are only a few such wells; enough, however, were found to reveal an intelligent idea of the shape of the pre-existing basins, and the character of the material filling them. These things are of the utmost importance for working out the history of the region, which was the chief reason of time and attention being given to the area.

The easternmost part of the region is very probably not filled to a great depth. The sinks that abound indicate that bed-rock is near the surface. Sections 4, 9 and 8 contain sinks; these sections are at the eastern margin, and it might be easily deduced that the bed-rock is near the surface.

Well No. 1. Out some distance in the basin, on the southern line of Section 5, T. 8 N., R. 2 W., near the headwaters of McCormick's Creek, is a well at Mr. Fife's which is twenty odd feet deep. This well furnishes a copious supply of water which comes from sand underlying a shallow surface stratum of soil. This well proves that the elongated elevation is a product of the forces which made the topography of the region, and not a remnant or hill in the former basin.

Well No. 2. This well is situated in the middle southern part of Section 6, T. 9 N., R. 2 W., and is on the edge of the silt line at the foot of the hill. It is sixteen feet deep and reaches solid stone. The material through which it passes seems to be entirely the outwash or talus from the hill rising up behind it.

Well No. 3. Middle northern part of Section 6, T. 9 N., R. 2 W., at W. Stone's. Surface elevation 730 feet. The depth of this well was not ascertained but reports indicate that it is of considerable depth to bed-rock.

Well No. 4. Southwestern part of Section 31, T. 10 N., R. 2 W., at H. Heady's. Surface elevation, 723 feet. Depth, 14 feet. Soil, with streaks of yellow and blue containing fine sand, 12.5 feet. Caked sand, of a yellowish sugary appearance when wet, becoming like brittle sandstone upon drying, 1.5 foot.

Well No. 5. This well is in the high hill to the north of well No. 4. It reaches bed-rock at a shallow depth.

Well No. 6. On the county line, middle eastern part of Section 36,

T. 10 N., R. 3 W., at Mr. Whitsell's. Surface elevation, 720 feet. Depth, 12 feet. Under a shallow surface soil occurs a blue clay which contains very fine, angular sand grains. These grains are invisible to the eye, though they may be felt when the dry material is rubbed between the finger and thumb. This very fine sandy clay is very tough, tenacious, and slightly sticky when wet, and of a distinct blue color. On exposure to the air it takes on a brownish-grey hue. When dry it is an ash color. Application of muriatic acid shows that it is highly charged with calcium carbonate. (This detailed description is given because this material is often encountered in the lower parts of wells. Hereafter it will be designated as blue clay. It will be further discussed later.)

Well No. 7. One-fourth mile north of well No. 6, at the residence of B. Smith, is a well which reaches bed-rock at a depth of 12 feet. The surface elevation is 720 feet.

Well No. 8. One-eighth mile northeast of well No. 7, N. W. corner of Section 31, T. 10 N., R. 2 W., at C. Wampler's. Surface elevation, 720 feet.

Soils	17 feet.
Imbedded logs.....	1 foot.
Clay	8 feet.
Gravel and sand.....	1 foot.
Limestone	

Well No. 9. One-eighth mile north of No. 8, southwest corner of Section 30. Surface elevation, 720 feet. Depth to stone, 51 feet.

Soil and clay.....	18 feet.
Sand and gravel.....	4 feet.
Blue clay.....	29 feet.
Limestone	

In the northwest corner of Section 25 and leading far into Section 24, T. 10 N., R. 3 W., is an arm or extension of the Flatwoods basin, which is filled very little. Sinks are very numerous, showing limestone in many places. The general elevation of this extension is 740 feet.

Well No. 10. In the northwest part of Section 26, T. 10 N., R. 3 W., at Frank Marshall's. Surface elevation, 725 feet. Depth to stone, 47 feet.

Sandy clay soil.....	40 feet.
Sand and gravel.....	4 feet.
Soil	3 feet.
Limestone	

Mr. Marshall also has a dug well; it is twenty feet deep. It seems to be in clay soil except near the bottom, where sand occurs. A stick was found in this sand.

Some twenty-five rods east of Mr. Marshall's drilled well is the ford on the road across McCormicks Creek, where rock outcrops at 700 feet. Bed-rock was struck at least twenty-two feet lower than this in the Marshall well. Farther up McCormicks Creek, about three-quarters of a mile, bed-rock is to be seen in the creek bed, and just above, two remarkably large springs pour forth clear, cool limestone water, indicating that bed-rock is near the surface.

Well No. 11. Middle western part of Section 26, one-half mile southwest of No. 10. Surface elevation, 740 feet. Depth to stone, 110 feet.

Soil	20 feet.
Sand	16 feet.
Blue clay.....	74 feet.
Limestone	21 feet

Well No. 12. Northwest corner Section 35, one-half mile south of No. 11. Surface elevation, 740 feet. Depth to stone, 74 feet.

Well No. 13. East of the center of Section 35, T. 10 N., R. 3 W. Surface elevation, 735 feet. Depth, 22 feet. This well seems to be entirely in a reddish sand.

Well No. 14. Center of Section 35, one-eighth mile west of No. 13, at John Leonard's. Surface elevation 735 feet. No stone reached at a depth of 116 feet.

Soil	1 foot.
White sand, with small pebbles infrequently.	.80 feet.
Blue clay	36 feet.

Well No. 15. Southwest corner of Section 35, one-half mile southwest of No. 14, in an open field. Dug well. Surface elevation, 745 feet. Depth, 41 feet.

Well No. 16. One-eighth mile east of Stogsdill Pond, Section 2. Surface elevation, 820 feet. This well is 40 feet deep, and is entirely in a reddish sand containing some water-worn gravels. This well contain no water.

Well No. 17. One-sixteenth mile south of No. 16. Surface elevation, 795 feet. This well was reported by two persons to be 80 feet deep, in a reddish sand its entire depth. The writer is inclined to believe there is some mistake regarding its depth. It is not likely over 50 feet deep.

Well No. 18. One-eighth mile southeast of No. 17, at Geo. Myers'. Surface elevation, 825 feet. Depth to stone, 25 feet, in seemingly the residual limestone soil.

Well No. 19. One-fourth mile southeast of No. 18, on high, rounded hill, at Amos Barker's residence. Surface elevation, 860 feet. This well encountered Chester sandstone at a shallow depth.

Well No. 20. One-fourth mile northeast of No. 19. Surface elevation, 825 feet. This well penetrates almost pure sand to a considerable depth.

Well No. 21. Middle of southeast $\frac{1}{4}$ Section 3, T. 9 N., R. 3 W., at A. Evans'. Surface elevation, 780 feet. Depth, 42 feet.

Soil	18 feet.
Reddish sand (quicksand).....	24 feet.

Well No. 22. One-fourth mile south of No. 21, at C. R. Ellis's. Surface elevation, 775 feet. Depth to stone, 51 feet.

White to yellow soil.....	18 feet.
Water-worn gravel and sand.....	6 feet.
Quicksand	27 feet.
Limestone	92 feet.

Well No. 23. Middle of northern one-half Section 10, one-half mile south of No. 22, at the County Farm. Surface elevation, 765 feet. This well penetrates soil and sand nearly fifty feet.

Well No. 24. Center of Section 23, T. 9 N., R. 2 W., at A. O. Collins'. Surface elevation, 690 feet. Depth to stone, 88 feet.

Red sand and clay.....	35 feet.
Blue clay.....	53 feet.
Limestone	1 foot.

This completes the number of wells from which data was secured. The reader can see at once that the greater number of them reveal the fact that Flatwoods was in former times a basin much deeper than it is now. It seems that the basin was rather deep at the northern part of Section 6, as indicated by well No. 3. This deep portion extended northwest, entering Section 36, and thence northward, but went northward for only a short distance, along the present channel of McCormicks Creek, until it turned westward as indicated by the shallow bed-rock in well No. 7. Wells No. 8 and No. 9 indicate that a tributary channel passed near the southwest corner of Section 30. This channel probably entered the

main channel near the northern middle of Section 36. The bed-rock outcrop along McCormicks Creek in the southeast part of Section 26 indicates that the region to the northeast was high, very probably a divide between the tributary just mentioned and the one that undoubtedly came from the northeast of section 24. Wells Nos. 10, 11 and 12, by their depth to bed-rock, reveal a channel region running south and southwest from section 23, from near where McCormicks Creek leaves the Flatwoods region. A small tributary in bed-rock flowing its water in the up-stream direction, just west of where McCormicks Creek turns west in section 23, also indicates that a channel once went southwest from this region. There is no doubt in the mind of the writer that this region and also the long extension north into section 24, was drained to the west and north through a well developed underground system. Well No. 14 shows that the ancient surface was more than 116 feet below the present surface, and more than eighty-five feet below the bed of McCormicks Creek a mile to the north. At this place the old stream channel must not have been far from the 600 foot contour line.

The western border of Flatwoods in the region of the headwaters of Allistons Branch reveals facts in harmony with those shown by the wells. The tributaries of this creek are etching their way slowly into the Flatwoods region. The etching is slow because the slope is away from them, thereby causing practically all the water, except that which falls immediately into them, to flow in the opposite direction. These tributaries are deep, V-shaped valleys or ravines in the sandy material at this margin of the region. A typical ravine may be found just south of the middle of section 27. It begins at an elevation of 760 feet, and descends rapidly to a depth of sixty feet or more. The sandy banks on either side are very steep. Small erratic boulders may be found in the narrow bottom, amid which trickles water seeping from the sandy banks near the bottom. The descent continues rapidly until near the 620 foot contour. A small valley flat then begins to appear, and soon the main stream is reached. The entire length of this ravine is less than one-fourth mile, and a descent of at least 150 feet has been made. The slope away from the Flatwoods region towards White River, much cut up by etching ravines, is very rapid; this makes the banks or sides of the etching ravines higher near their heads than farther down.

The great amount of sandy material at the head waters of Allistons Branch indicates that the old channel found and traced westward by the

wells had its course entering Allistons Branch near its head-waters. The present Allistons Branch, then, is the lower part of the main stream that in former times principally drained the area now occupied by Flatwoods. The map showing the main drainage, the main channel and its tributaries, are inferences that can scarcely be avoided. There are indications, however, that portions were drained by underground channels to other streams; as, for instance, the extreme southeast corner, the eastern part of section 30 and the portions already mentioned near the northwest corner of the region. The sinks of these portions are evidences that their drainage was as it is now, and was only temporarily interfered with by the forces that destroyed the old drainage. Very probably other portions of considerable area were drained into the main channel from underground passages, the water coming to the surface farther down in the form of springs.

Next, the relation of Flatwoods proper with the adjunct south of Stogsdill Pond will be considered. The lack of any deep well immediately west of Stogsdill Pond leaves the data somewhat incomplete, but this portion has an elevation of 820 feet, and it seems likely that bed-rock is much higher here than either to the north or to the south. Bed-rock out-crops at 760 feet one-half mile west of this portion, and it is found at 800 feet in Mr. Myer's well one-half mile southeast. Thus it is very probable that there was a divide between Flatwoods proper and the small adjunct to the south. It must have been very low near Stogsdill Pond, for wells Nos. 16 and 17 penetrate sand their entire depth, and No. 17 was said to be eighty feet deep, but, as said before, the writer doubts the reputed depth of this well.

The region east of the Myer's well seems to have been considerably filled, as indicated by well No. 20, and the sharp ravines which are etching their way in the flat area near by. Examinations of these ravines show that they contain no bed-rock, but were rather grooved into the stratified sand and fine gravel of which the flat area is composed.

In the extreme southwest corner of section 1 is also a filled area, but farther east bed-rock is found, and on top of the ridge there is no sign of silt or sandy material. Undoubtedly the head of a stream reached into this corner, near where the present stream is endeavoring to clean out the filled-in material. Thus the evidence shows that a divide with a very irregular summit once separated the adjunct basin from Flatwoods proper.

Turning to the western edge of the adjunct basin, we find sharp V-shaped valleys or ravines of McBrides Creek etching their way into the

body of the flat in the same manner that the tributaries of Allistons Branch are etching their way into the western edge of Flatwoods proper. These ravines descend almost suddenly a hundred feet below the level of the flat. In several places the structure of the material can be seen. The upper fifteen to twenty feet is a fine, white soil, characteristic of the surface of the Flatwoods region. Underneath this, is reddish sand with layers of fine gravel alternating with the much thicker layers of sand. Water comes from the sand and gravel into the ravine, making them miry in the bottom. These ravines, in conjunction with the wells (Nos. 21, 22, and 23) clearly reveal that the region has been filled, and that the adjunct basin had a broad outlet, or opening, to the west. McBrides Creek must have extended much farther east, draining in all probability the greater part of the adjunct, and having its tributaries reaching to the divide between the adjunct and Flatwoods proper.

It might be mentioned that the streams in section 11 have cut themselves down into the filled material at least fifty feet and leave the old flat above as a beautiful terrace. The material of this terrace is shown in an excellent manner along the steep western side of the middle stream. It is as follows:

Soil	12 feet
Red sand.....	25 feet
Blue clay.....	5 feet

McBrides Creek undoubtedly had its upper portion and upper tributaries taken away from it by the forces that remade the topography of the region. But it is rapidly working its way back into its old domain in the same manner that Allistons Branch is trying to recapture its old basin. The rapidity with which these tributaries etch back into the filled material can be seen in a single ravine just north of C. R. Ellis' house in the middle of the southeast quarter section 3. The main stream flows parallel with the road, and the short tributaries come into it at right angles. These tributaries are, in fact, just immense gullies only a few rods long, but with the depth easily forty feet. The water during showers gushes into these ravines and carries away the easily transported sand at the bottom, leaving the soil above to slump into the ravine, which is then in turn rapidly carried away. These ravines grow directly in proportion to the amount of water entering them at their head. Mr. Ellis stated that eleven years ago, the head of the particular ravine mentioned above was at least sixty feet

west of the road. Now, it is at the very road ditch, and immediately descends twenty-five feet. If nothing should interfere with it, it would in a short time destroy the road here by etching into it.

There is no doubt in the mind of the writer that McBrides Creek at one time extended eastward to the very divide between the adjunct basin and Raccoon Creek, but evidence shows that probably most of the water that fell into its upper portion was carried by underground drainage either to the lower part of the stream or to the east and to the south to Raccoon Creek. At present practically all the water falling into the adjunct basin goes into Raccoon Creek, mainly through the sinks at the southeast corner of the adjunct region. Two very large springs come out about a mile south of where the water enters these sinks, which in all probability are outlets of underground channels beginning at the sinks in section 11.

The Raccoon Creek addition of Flatwoods may be treated in a few words. This region was the site of an old stream which followed the same valley that the present stream does. The present stream, as stated before, has cut itself down into the old flat and in some places has removed much of the filled-in material. Raccoon Creek, however, does not reach bed-rock until it is at least 100 feet below the old flat, as found in the southern part of section 26. Yet there is a short distance that the stream passes over rock in a rather constricted place in the middle northeast part of the same section. This portion of the stream has evidently been cut since the time that the old valley was filled. The old stream evidently passed to the east of this place, as indicated on the map.

The faces of the terraces have but few portions that show the material of the terraces, but the few that are shown, and the rock structure of the higher areas, along with the sinks in the terrace flat, reveal the ancient drainage lines in a very able manner. The small tributary entering Raccoon Creek just north of the mouth of Little Raccoon Creek once extended nearly three-fourths of a mile farther east than it does now. It is making heroic efforts to recapture its old drainage basin; it is being aided by underground drainage, much material having been carried away leaving great sinks in the one-time flat. A small tributary no more than one-fourth mile long enters Raccoon Creek from the east in the middle of section 23. This tributary is a very small one in comparison to its predecessor. The old tributary extended nearly two miles eastward. Practically all of this region is now drained by sinks, which have caused the old flat to be considerably depressed locally. Well No. 24 is near the site of this old

tributary. It shows the filled-in material to be eighty-eight feet deep; that makes the old tributary somewhat below the present level of Raccoon Creek, where the present tributary enters. But not more than thirty rods to the northeast of the well, limestone outcrops. This indicates that the long hill protruding westward from section 24 was continued as the northern divide of the old tributary.

THE PRE-GLACIAL GEOLOGY OF FLATWOODS.

In considering the geology of the Flatwoods region, it is thought best to divide it into two main divisions. The first division includes the rock structure of the region and the subsequent history up to the Pleistocene. The second division begins with the Pleistocene and takes in all up to the present. The rock structure of the region and immediate vicinity is of Middle Mississippian age with some Pennsylvanian bordering closely on the west. The stratigraphy of the region will be given a brief treatment.

Knobstone Group. Just to the east of the region in the valley of Jacks Defeat Creek, are the upper portions of the thick Knobstone Group of shales and sandstones. The area of its outcrop in Indiana is a strip of territory some twenty-five miles wide, extending north, northwest from Floyd County to Benton County. This formation consists of compact, insoluble, impervious sandstones and shales, aggregating a thickness from 400 to 600 feet. The topography of the outcrop, resulting from the peculiar weathering of the rocks, is of a distinct type. The rocks absorb water readily, but transmit it poorly, so that they are easily shattered by freezing and thawing. The region is weathered and eroded into deep, steep-sided valleys of very pronounced relief. Brown County affords a typical example of Knobstone topography. Another characteristic of this group of rocks is the general absence of fossils, such being present only very locally. It seems that the shales and sandstones were laid down in impure, muddy waters, which were not on the whole very favorable for the life of water-breathing animals.

Harrodsburg Limestones. Overlying the Knobstone is the Harrodsburg limestone, with a thickness varying from sixty to 100 feet. This limestone consists of several heavy bedded layers of hard, gray to blue, often highly crystalline stone. There are occasional intercalated thin beds of shale. In some sections a very cherty layer occurs. The top member of this stone often is very massive, and its texture is very similar to the oölitic bed overlying. Geodes are characteristic. The limestone as a whole is

very fossiliferous, showing that life was abundant when it was laid down. The area of outcrop is narrow and intermingles with the western edge of the Knobstone, often capping outliers of the latter. It has no distinct topography outside of the fact that sinks occur in it, which never occur in the Knobstone, and rarely in the formation above it.

Salem Limestone. Superior to the Harrodsburg formation is the famous Salem limestone known as the Oölitic, or Bedford limestone. It is an excellent building stone, and is known as such all over the United States. Typical outcrops of it occur along the valley of Jacks Defeat. The stone is usually massive, with few indications of bedding, varying from a few feet up to eighty and ninety feet in extreme cases. The stone typically is a porous stone composed of nearly pure calcium carbonate. It is made up principally of broken animal remains and several species of Protozoa, among which the main one is *Endothyra baileyi*. These have all been cemented together in a loose manner. The area of its outcrop is characterized by long, gentle slopes, rounded hills and general undulating topography.

Mitchell Limestone. The Mitchell limestone is the one that we are mostly concerned with, since practically the entire basin of the Flatwoods region is in this stone. It ranges in thickness from a few feet in its northern outcrop to 250 feet in the southern part of the State. The stratigraphy of this formation is rather varied, as there are rarely two successive layers alike. In general it consists of impure limestones and calcareous shales, usually thin. Many layers are very hard, and weather white, sometimes small slabs having the appearance of bleached bone, and on being struck have a metallic ring. Such layers usually have numerous right-angled joint cracks, and are semi-lithographic, breaking with a sub-concoidal fracture. The upper members of this formation are usually a beautiful oölitic structure. As a rule the limestone is fossiliferous. The area of the outcrop of the Mitchell limestone extends over a broad plain which narrows to the north and pinches out in Montgomery County. It is essentially a cave-bearing formation, containing some of the most famous caves in the world. Wyandotte and Marengo caves of Crawford County and Mammoth cave of Kentucky are in this stone. The region of its outcrop is pitted with sinks, and underground drainage is a distinct and noticeable feature. Lost River of Orange County is a typical example of an underground stream in the Mitchell area. Hundreds of sinks occur on the borders of the Flatwoods region, where the filled material is relatively

thin. These sinks connecting with under-channels are the sources of many springs that abound in and at the borders of the region.

Chester Group. On the high hills at the borders of the Flatwoods region shales and shaley sandstones overlie the Mitchell. These are usually about twenty-five feet thick, and upon them is a limestone usually about three feet thick in the Flatwoods exposures. This group is known as the First, or Lower Chester. On top of the First Chester limestone occur a series of shales and sandstones, consisting of a portion of the Second Chester group. This group is capped by a limestone, its last member. At no place in the immediate vicinity of Flatwoods was the top of the Second Chester found.

The above general outlined stratigraphy is found in the Flatwoods region and portions directly connected with it. Some attention now will be given to local details.

The general dip of the rock of the Bloomington Quadrangle, in which a portion of Flatwoods occurs, is on the average twenty feet to the mile to the west, southwest. In regard to the dip of the rock structure in the Flatwoods region, the following data reveal an interesting feature:

Contract of Mitchell and First Chester, south side Flatwoods:

In 920-foot hill, N. W. $\frac{1}{4}$, sec. 16, T. 9 N., R. 2 W.....	870
In 900-foot hills, S. $\frac{1}{2}$, sec. 7, T. 9 N., R. 2 W.....	845
In W. part of hill N. W. $\frac{1}{4}$, sec. 7, T. 9 N., R. 2 W.....	820
In centre of S. $\frac{1}{2}$, sec. 1, T. 9 N., R. 3 W.....	800
One-half mile N. of above.....	805
S. W. side of hill, S. E. $\frac{1}{4}$ sec. 2, T. 9 N., R. 3 W.....	800
Middle N. of N. W. $\frac{1}{4}$ sec. 3, T. 9 N., R. 3 W.....	720

North side of Flatwoods:

In 810-foot hill, Middle sec. 30, T. 10, N., R. 2 W.....	790
Chambers Hill, middle of line, sec. 24-25, T. 10 N., R. 3 W.....	810

The data along the southern side of Flatwoods reveals an average dip of thirty-feet to the mile along a line which is as much in the direction of the strike as in the direction of the dip. Again, going from the eastern contacts toward the western, the data indicates that the first two-thirds of the distance has a dip of twenty feet to the mile, and the remaining distance a dip of fifty feet to the mile. The data on the north side is very meagre, as there are only a few hills high enough to reach the Mitchell and Chester

contact. Only the eastern half of the north side is represented by contacts. Here, two contacts were found, but these reveal a rather striking feature. The two contacts are one and one-fourth miles apart, the west one being a mile west and one-half mile north of the eastern contact. Judging from the general dip of the rock structure, one would expect the western contact to be several feet lower than the eastern one. But the reverse is true, the eastern contact is actually twenty feet lower.

Having absent contacts for the western part of the north side, the data does not give absolute information, but the indications point decidedly to the fact that a monocline or an anticline, with its long axis extending parallel with McCormicks Creek, exists in the Flatwoods region. This means that Flatwoods proper is an eroded anticline. This interesting structure indicates that the possibilities for oil are much stronger than a guess. However, it is possible and even probable that the anticline is superficial, and does not extend to strata of oil-bearing properties. The only sure method for determining the presence of oil is the drilling of a hole the required depth. The writer, however, is of the opinion that the Flatwoods region has sufficient superficial indications of oil to warrant an experimental hole being made.

An interesting chapter in the pre-glacial geology of the Flatwoods region is found in the physiographic development. Several million years of time lapsed between the time when the later Mississippian deposits of the region were first lifted above sea level and the advent of the glacier in quaternary times. During this time, no deposits were made, indicating that the region was never again subject to such a depression that would reduce it below sea level. There is no rock structure present representing the late Paleozoic, the entire Mesozoic and the early Cenozoic systems. During the lapse of all this time, the area was subject to all the forces of weathering and erosion. Undoubtedly much must have been accomplished in that long interval of time.

It is very likely that the new land surface of the upper Mississippian strata was not raised to any great height above sea level for a long time. The old sea in which the Pennsylvanian rocks west of the area were deposited came and went many times before it left the region forever. In the ages that followed the withdrawal of the old sea, perhaps several peneplains were formed, and each, in turn, destroyed by the subsequent erosion, following the successive uplifts of the area. Only the later of these would be preserved in any recognizable degree.

A study of the topography of the Bloomington Quadrangle shows that there are several rather flat-topped, isolated hills and several long, irregular, flat-topped ridges, all of practically the same height. The hills reaching the 900-foot contour and forming the southern rim of Flatwoods proper, are examples of the isolated hills. A typical example of the flat-topped ridges is found in the region of Kirksville, where the irregular ridge extends for several miles in a north and south direction. These high, flat-topped hills and ridges are undoubtedly the remnants of a former peneplain, which may be correlated; but not absolutely beyond doubt, with that at the base of the Cumberland Plateau, and with its continuation southward and westward into Tennessee, thence northward into Kentucky, where it becomes known as the Lexington Plain. This correlation makes it of early Tertiary age. (J. W. Beede, *Features of Subterranean Drainage in the Bloomington Quadrangle*. Proceedings of the Indiana Academy of Science 1910. Ditney Folio, Indiana, U. S. Geol. Sur.) Beede has named the peneplain, which these isolated heights represent, the Kirksville peneplain, after the typical development of it at the little village of Kirksville in southwestern Monroe County. Representatives of it occur along the south side of Flatwoods and between the two main branches of Raccoon Creek. Chambers Hill on the north side is also a representative of it.

Succeeding this peneplanation there was an uplift of the region of about 175 feet. The streams went to work again, cutting deep valleys in the Kirksville peneplain. In time the stream reached base-level, and by lateral erosion and beveling by the minor tributaries, local peneplains were developed. The wide expanse west of Bloomington, which continues southward through Lawrence, Washington and Crawford Counties, is the best and most strikingly preserved area representing this peneplanation. This plain is in the Mitchell limestone, and is well represented at Mitchell, Lawrence County. It had its maximum development in late Tertiary times, and is the Mitchell plain of Beede, being so designated by him. (*Features of Subterranean Drainage in the Bloomington Quadrangle*. The Proceedings of the Indiana Academy of Science, 1910.) Flatwoods at this time was peneplaned, the rim of the higher land and the monadnocks being the remnants of the old Kirksville peneplain. Flatwoods is really a portion of the Mitchell plain. The drainage in late Tertiary times was into White River. The main stream was probably in the long axis of the

region with its outlet through Allistons Branch. This main stream and its principal tributaries were in their old age, and were wandering about the plain, being separated from each other and adjoining stream basins by low divides, except locally where remnants of the preceding peneplain persisted.

Near the end of Tertiary time there occurred another upheaval; this time of about 300 feet. The streams immediately began to corrade their channels, and in the course of time the main streams cut their channels to base level in their upper and middle courses. In early Pleistocene times there occurred a depression of about 150 feet, which caused the base level portions of the streams to become filled. Wabash and the White rivers show this in an ideal manner. Bean Blossom and Jacks Defeat creeks are excellent examples of smaller streams which have their lower and middle courses filled as a result of the depression of the land. But even these streams are still corradng their channels in their upper courses. Examples of streams which are still cutting their channels, down in the late Tertiary or the Mitchell plain are found in Stouts Branch and Rocky Branch, north of Bloomington, Clear Creek south of Bloomington, and many other small streams reaching into the Mitchell plain. However, such streams are confined to the margins of this plain, because of the peculiarity of the Mitchell limestone in its tendency for the formation of sinks and subterranean drainage. Only the major drainage lines cross this formation with an open channel. Beede has treated this subject thoroughly in the paper referred to above.

In the Flatwoods region the main stream and main tributaries were about as indicated on the map showing the pre-glacial drainage. Considerable portions of the region, however, were drained by sinks and underchannels, as is characteristic of the Mitchell plain west and southwest of Bloomington. But despite underground drainage, the lower part of the main streams and principal tributaries were cut down to base-level. No doubt many springs came into the streams, being the outlets of the underground channels. At the margins underground drainage undoubtedly carried water to other streams. Instances of this kind have already been given.

At this point it is deemed advisable to give some attention to White River, near the lower end of Flatwoods. Collet in his report on the geology of Owen County (Seventh Annual Report Indiana Geol. Sur., 1875), makes note of the extreme narrowness of the White River valley between Romona

and Spencer. He accounts for "The Narrows", as this very constricted portion of the river valley is called, by asserting that this portion of the valley is new, having been formed since the Illinois glaciation. The old channel, he says, was up McCormicks Creek, through the Flatwoods basin, and back to the present channel by way of Raccoon Creek. Leverett says: "The stream (White River) is now occupying a pre-glacial valley for a few miles in southwest Morgan County, and is also in a pre-glacial valley throughout much of its course below Owen County. But in its passage across Owen County it is opening a new valley. It has been suggested that this stream had a subterranean passage across the sink-hole region of Owen County, in which case no well-defined surface channel may have been opened prior to the glacial invasion." (The Illinois Glacial Lobe pp. 104, Monograph XXXVIII, U. S. Geol. Surv.)

Both Collet and Leverett have expressed their belief that White River in its present passage across the Mitchell limestone region is in a new valley. Collet says that the old channel was through McCormicks Creek, Flatwoods basin and Raccoon Creek. Siebenthal, in regard to Collet's idea, says: "The Pleistocene terraces of Bean Blossom Creek clearly prove the pre-glacial valley of that creek to have been practically as it is at present. It is impossible to imagine how it could be cut down to its present depth, while White River, into which it emptied, was running at a level 150 feet higher than now, as it is alleged to have done. Moreover the gorge of McCormicks Creek is clearly post-glacial. And further, it empties into White River at least a mile below the upper end of the "Narrows," whose existence it was brought forward to explain." (Twenty-first Annual Report, Ind. Geol. Surv., 1896, pp. 302). Thus Seibenthal makes it clear that Collet was in error in regard to the ancient channel of White River in the Mitchell limestone region.

Leverett asserts that White River in Owen County is post-glacial, and suggests that the pre-glacial drainage was a series of channels through the limestone region. The writer in his examination of the area found no evidence of an ancient channel on either side of the present river in the limestone region of Owen County; and there is little, if any, evidence of its passage through the region ever having been subterranean.

The constriction of the river just above Spencer is undoubtedly remarkable, and not geologists alone have asked the why of it. Puzzling as the "Narrows" are, they have a rather simple explanation. The valley here is very narrow in comparison to the extremely wide portions above

and below. The valley is wide in Morgan County because of the easily eroded Knobstone sandstones and shales, through which the valley has been cut. It is wide in Greene County again on account of the same fundamental reason. Here the valley is in the soft sandstones and shales of the Chester Group and the Coal Measures. In this county, however, the Illinois glacier undoubtedly was an important factor in widening the valley. In Owen County the strata in which the "Narrows" occur, are hard, resisting limestones, which are little disintegrated by weathering and suffer even less by abrasions. The widening of the valley in this region must be carried on mainly by solution, which is a much slower process than those involved in the region above and below. This narrowing of the valley is identical with the appearance of the limestone bluffs in the vicinity of Gosport. The same condition is to be seen on the East Fork of the White River, where the valley is exceedingly wide in the Knobstone region, and becomes almost gorge-like in the limestone region.

Furthermore, if the valley in the limestone region were post-glacial there would be very little alluvium below the present channel. This is not the case. Wells at Spencer prove that the alluvium is at least 100 feet deep, just as it is in the wide regions of the valley.

White River Valley, then, in its passage across the limestone region of Owen County is not a new opening. It is the same valley that is seen in the wide portions of both Morgan and Greene counties. It is the same valley that has carried the waters of the basin above since the time that the present fundamental topographic features were initiated. In fact this part of the valley and channel is more nearly where it has always been than any part either above or below, for the simple reason that at this point the Illinois glacier but little more than crossed the valley, while both above and below, it crossed for many miles farther, and deranged the drainage accordingly.

We are now ready for the final chapter of the history of the Flatwoods region, the chapter which really gives the explanation of the Flatwoods phenomena.

THE GLACIAL HISTORY OF FLATWOODS.

It is not the writer's purpose to give here a treatise on glaciers and glaciation, nor to give an intricate and detailed history of the period of glaciation known to have been present in the Flatwoods region. The purpose here rather is to show the relations of the edge of the Illinois ice-

sheet to the Flatwoods region and immediate vicinity, and show how it was responsible for the peculiar phenomena of the area. For an intimate knowledge of glaciology and its broad relationships, the reader's attention must be given to the many text-books and matter dealing specifically with such phases. That glaciation has taken place over very large areas of the world is no longer a theory. The most obtuse have long been convinced of that fact. The most important phase concerning glaciation before the scientific world today, is the manner in which it took place in specific areas. The most interesting features of this phase of glaciation occur in the phenomena existing along the border, or near the border, of the one-time ice-sheet edge. The Flatwoods region belongs to this phase.

Leverett has given a detailed account of the drift border in southwestern Indiana. The drift border through Greene, Owen and Monroe counties, he credits to C. E. Siebenthal. "From near Scotland (southern Greene County) it has a course slightly east of north to the valley of Plummers Creek, in section 9, T. N., R. 4 W. North of this creek it makes an eastward protrusion of about two miles into a lowland tract known as the American Bottom, reaching section 36, T. 7 N., R. 4 W. North of this lowland the course of the boundary is west of north to the valley of Richland Creek, in section 9, T. 7 N., R. 4 W. It follows the east bluff for about three miles and crosses to the west side of the creek in section 35, T. 8 N., R. 4 W. It follows nearly the west bluff to section 17, T. 8 N., R. 3 W., passing about a mile southeast of the village of Newark. The boundary makes an eastward protrusion of about a mile into Richland Creek valley in section 16, from which the course is northward into Owen County. Entering Owen County in section 33, T. 9 N., R. 3 W., the boundary leads north-eastward past Freeman post office and crosses into Monroe County in section 6, T. 9 N., R. 2 W. The course continues northeastward through northern Monroe County, the boundary being about two miles north of Ellettsville and one mile north of Modesto, and coinciding nearly with Indian Creek valley from mouth to source." (Monogr. XXXVIII, U. S. Geol. Surv., pp. 34-38.)

The above detailed line of the glacial limit places the whole of the Flatwoods region well within the limits of the drift line. While it is not the intention of the writer to disprove the general limit of the drift as interpreted by Leverett, there must be some variation made in the Flatwoods region. Perhaps it should be made clear that in the detailed work done by the writer, searching attention was given to the probable advance

of the ice-sheet itself, rather than to the exact drift line. In interpreting the Flatwoods phenomena the position of the advance of the ice-sheet itself is of fundamental importance. This line of advancement, as given below, was determined not only by the presence of erratic boulders and rocks, but by stratified outwash material as well. Of the two phenomena, perhaps the latter is the more important.

A close examination of the hills or rather ridge extending eastward along the northern part of section 15, T. 9 N., R. 3 W., thence northeast through section 11 and into section 1, shows that the ice-sheet never crossed beyond. Evidence is plentiful in showing that the ice rested against this ridge and remained close to it for some time. The west side of the small adjunct is an outwash plain, which in the headwaters of McBrides Creek shows the coarse layers of stratified gravel alternating with both coarse and fine sand. About a mile west of the Stogsdill Pond is the remnants of an old moraine, showing the last stand made by the ice front. The coarse sand found in the ridge, dividing Flatwoods proper from the adjunct, is partly outwash material and partly the result of the ice-front itself in pushing material against the ridges, which the water later worked over. Undoubtedly a tongue of the ice-front pushed up to the very upper tributaries of the old stream, the lower part of which is now represented by McBrides Creek, but that it never crossed the ridge between the old stream and Raccoon Creek to the east is certain.

To the north the ice came up the White River slope and pushed up the old stream, draining the region now occupied by Flatwoods proper. It may have come up this old stream as far east as the Owen-Monroe County line, and even some distance farther, but for the most time it must have remained near the western border of the present Flatwoods. Alliston's Branch has been eating its way into an old outwash plain since the withdrawal of the ice from the region. It has erased the moraines, if any were formed, and has taken considerable of the head of the outwash plain. It is this outwash material, covered with later silt, that makes the slope of the western part of Flatwoods toward the centre of the region. It is evident from this that the ice-front for the most part did not extend beyond the headwaters of Alliston's Branch, in sections 26 and 35, T. 10 N., R. 3 W.

No outwash material or erratic boulders were found in the extension of the Flatwoods basin into section 24, or on Chambers Hill, but erratic boulders were found in the northern part of section 23 on the White River

slope, away from the high periphery at this part of Flatwoods. It is improbable that this immediate portion was ever covered with ice, or if it was, only for a short time.

The headwaters of Big Creek in sections 24, T. 10 N., R. 3 W., and 19, T. 10 N., are in outwash material, and erratic boulders are common. But these conditions are not found on the Flatwoods side of the divide. It is very probable that the ice-front never got over this divide, although the waters came freely into the Flatwoods region over this divide in the southwest corner of section 19.

But little attention was given to the probable front of the ice east of Big Creek, but indications are that it pushed up Jacks Defeat valley to near the middle of section 28, T. 10 N., R. 2 W. Its maximum advance east of Jacks Defeat was probably where indicated by Leverett. Yet it seems to the writer, for reasons that will appear later, that for the greater time it must not have been farther south than the middle of sections 21 and 22, T. 10 N., R. 2 W.

The line of the advance of the ice-sheet south of Freeman was probably as indicated by Leverett. The advance was somewhat east of Freeman and continued northeast to near the southwest corner of section 23, T. 9 N., R. 3 W., where it turned northwest, jutting against the high ridge extending northward and northwest to the southeast corner of section 9. Here the ice crossed the ridge and protruded eastward, as already described. But it is not likely the ice-front was east of Freeman during the time of the formation of the basin flat. The material of the flat west of Freeman is principally sand, and quite likely is the lower end of an outwash plain.

Although the above details show the tracing of a line for the advance of the ice-front, it must be continually borne in mind that the ice-front was not irregular in outline, but that it was constantly changing in a sort of backward and forward movement, due to the seasonal changes in temperature. This line, as described above, has practically all of the Flatwoods region outside of the ice limit, and in some places several miles within the limit placed by Leverett. Some glacial material was found well outside the limit as traced above, but it was material carried there by currents of water, and is not to be interpreted as being direct evidence of the presence of ice. Again, it is possible that the extreme limit of the ice-sheet came as far as indicated by Leverett. But it is quite evident that it

was not advanced that far when the flat topography of Flatwoods was formed.

Having determined the advance of the ice-front, it is easy to picture the fluvial conditions that existed during the presence of this great ice barrier. Streams of water, sometimes constricted and sometimes in broad sheets, poured out from the ice sheets in the summer seasons, and worked over the sandy debris, which the ice continually brought forward from the nearby sandstone hills. This debris was made into the outwash plains already mentioned as being along the headwaters of Alliston's Branch and McBride's Creek. In the adjunct basin the water from the ice carried material south and southeast from the region of accumulation, filling this basin to the level of the narrow outlet south of section 11. The water in being concentrated through this opening undoubtedly cut it down considerably. The material in the wide terrace southeast of this opening contains but little material foreign to the immediate vicinity, but there is an abundance of limestone, chert and sandstone material in the debris. Where the stream has cut a fresh place in the terrace these materials are shown in abundance.

Turning to the eastern side of the region, we find conditions which aided greatly in the making of the Flatwoods region as it appears today. White River was within the ice limit of the glacier and was at this time frozen with all of its upper tributaries, and incorporated within the glacier itself. Practically all of the tributaries beyond Bean Blossom were within this great ice clasp. The greater part of the Bean Blossom area, however, was free to gather its waters before the ice-front in the summer seasons at least. But the outlet and a few miles of the lower portion of this stream were within the ice limit. Consequently a lake gathered before the ice-front, both from the melting ice and the drainage of Bean Blossom basin. The water soon reached the height of the lowest point in the divide between Bean Blossom and Jacks Defeat Creek. This seems to have been about one and a half miles southeast of Stinesville, near the middle of sections 21 and 22, T. 10 N., R. 2 W., on the farm of Jack Litten. Here occurs a col nearly one-fourth mile wide, which comes down at least 100 feet below the height of the divide south of this place. The elevation of the col is 715 feet. Locally this col is known as the "Valley." Since it is on the farm of Mr. Litten, the writer proposes to call it the Litten Col.

The water on passing through the Litten Col came into Jacks Defeat valley, but again it had to lodge against the ice wall until it had found an

outlet. This was found just northwest of Ellettsville through the deep col, which has been described as the southeastern opening of the Flatwoods region. This opening shall be henceforth called the Ellettsville Col. The water however was not confined to this col alone; it came over the divide to the north, especially through the region where a small stream leads from the extreme southwest corner of section 33, T. 10 N., R. 2 W. This brought the water coming from the Bean Blossom region and also that coming from the melting ice into the Flatwoods region. In the first part of this paper it was mentioned that the slope of the eastern Flatwoods region was very gentle from the periphery, and that the deposits extended much higher at this side of the basin than anywhere else. These features, along with the presence of geode fragments in Raccoon Creek valley, were instrumental in the investigation of the possibility of the entrance of the Bean Blossom waters.

After entering the Flatwoods region an outlet for the inflowing waters, mingling with that which was coming from the western ice-front, was found through the opening near the Owen-Mouroe County line in section 1. This outlet, which might be designated as the Raccoon Creek Col, has already been described. But again the waters were to be checked by the ice-front in the Raccoon Creek Valley near Freeman. There must have been an outlet, either under the ice or around it, in the vicinity of Freeman near the 700-foot contour line; but such an outlet was not discovered. The terraces come to that height here, and slope from the above regions, indicating that the outlet at the time that the terrace-flat was formed could not have been either higher or lower.

Since the outlet in the Freeman vicinity was at the 700-foot contour line, it is easy to see the terraces representing the old lake bottom would not be any higher at this point, but that they should be higher above this point. This is true, as has already been described. The terraces up Little Raccoon Creek also get twenty-five or thirty feet higher in the upper region, where they fade out into the recent alluvium.

It is quite probable that for some time the waters entered Raccoon Creek just south of the Reeves School through a col on the section line, between sections 8 and 17, T. 9 N., R. 2 W. This col is about 770 feet in elevation, and is barely above the level of the silt line on the periphery of the region. During the time the water went through this col, the ice was advanced far enough east to obstruct the passage-way through the Raccoon Creek col.

In section 30, T. 10 N., R. 2 W., there is a connection through a narrow opening in the periphery of the Flatwoods basin to the outwash plain at the headquarters of Big Creek. No doubt waters came through this opening into the Flatwoods region. The silt flat on the Big Creek side of the opening comes up to the level of the opening, while on the Flatwoods side it is much lower, due to the sinks at this place. No doubt that at one time the slope was gradually away from this opening on the south side.

It is evident that the fluvial conditions existed for a long time, long enough to fill practically the entire stretch from the place of the entrance of the waters into the region to where they left in the vicinity of Freeman, with the material which has been described. These waters, being so intimately connected with the glacier, carried much of its material, and it is possible that at times large pieces of floating ice carried glacial material; thus one may expect to find glacial pebbles and even small boulders in any part of the region covered by these waters.

Indications show that for a long time there were no rapidly flowing waters in the Flatwoods region. In nearly all of the deeper wells and in many of the shallow ones the very fine sandy blue clay, which has been described in well No. 6 is characteristic of the lower material. This blue clay is thicker, perhaps, than any other material laid down. It can be seen in the bottom of the middle branch in the small adjunct of the Flatwoods region. Here it is as described, but reveals another feature quite important. Where it has weathered, or has been eroded, it shows laminae, indicating that it is a water-laid material. This delicate bedding of lamination cannot be detected in the clay when it is cut into. The elevation of this material as found in the wells is as follows (See map for location of wells):

Well No. 5.....	715
Well No. 7.....	690
Well No. 10.....	680
Well No. 11.....	700
Well No. 14.....	660
Well No. 24.....	655
In Flatwoods Adjunct.....	705

In other wells it never occurred, or was not identified as such. These figures show that the surface of this material was not actually flat, but that it was even flatter than the present surface of Flatwoods. It is very probable that before later material was deposited upon it, it was eroded in

places and in other places was entirely washed away. Generally, either sand or gravel overlies it, indicating that rapid flowing waters followed the long period of quiet waters in which the clay was deposited.

The next step in the history of the Flatwoods region comes with the deposition of the loess material. This fine, close-textured, ash-gray-to-white material composes the present surface of the region, except where the long duration of swamps has built up a black soil, which in some places is several feet deep over the top of the loess. The loess is somewhat uncertain as to its exact time and manner of deposition, and also as to the origin of the material. It was in all probability deposited not so very long after the disappearance of the Illinois glacier. Its thickness varies greatly in the Flatwoods region. Usually it is very thick at the margin of the old lake, being as much as twenty or even thirty feet; but in the centre of the region it is much thinner, at times being scarcely discernible. It is thicker at the margin because it has been washed from the hills adjacent. In the interior, perhaps much of it has been incorporated with organic material in the making of the black soils.

In connection with the history of Flatwoods, McCormicks Creek adds a very fascinating chapter. After the withdrawal of the ice, the lowest outlet was still the Raccoon Creek col. The region in the vicinity of the headwaters of Alliston's Branch was thirty feet or more higher than the Raccoon Creek col, having been built up by a possible moraine and the outwash plain already described. Therefore, the waters lowered to the level of the Raccoon Creek col. The water within the region was the site of a shallow lake, not being any deeper than the lowest place below the Raccoon Creek col. But the waters soon fell below the level of this opening, because of the opening of the old sinks near the margin of the region. These sinks were not covered deeply by the silt and sand, and consequently soon opened their old channels which had been filled or partly filled. It is very likely that the sinks formed in the northwestern margin or portion of the region in section 23, T. 10 N., R. 3 W., were the lowest and were, perhaps, the first to be opened on account of the static head at this place. These sinks had been draining this immediate portion of the Flatwoods basin long before the advent of the Illinois glacier, and it is quite likely that a thorough drainage, though underground, was already established along the line of the present McCormick's Creek Gorge. It can be easily seen that below the lower part of the gorge the valley becomes wider, and on approaching the river it is a filled valley. This clearly shows

that a pre-glacial tributary of White River occupied the region below the gorge of McCormick's Creek. The underground drainage merely had to develop through the narrow divide, composed entirely of Mitchell limestone. The great static head given the waters which reached into the southern part of section 23, in connection with a favorable dip of the rock, insured underground drainage. Moreover, the nature of the rock as exposed in the gorge would make this drainage not only apt, but very rapid. The hard Mitchell limestone in the upper part of the gorge is a continual occurrence of small irregular folds and dips, and even small faults, all being more or less disrupted and broken. Then below comes in a structure of loose, brecciated, highly argillaceous limestone, which is very easily eroded and carried away. This material is so loose that it can actually be torn from place by the bare hands. It is exposed in the falls and in the sides of the perpendicular cliffs below.

Thus, the present drainage of Flatwoods was initiated through the present McCormick's Creek by underground drainage. On account of the peculiar structure of the rock and the great fall, the drainage soon became open, somewhat as it is at present.

While the above outlined drainage through McCormick's Creek was in progress the middle part of Flatwoods remained a shallow lake. But finally the stream cut down enough to drain the region, with the exception of the middle portion and several small isolated depressions. Stogsdill Pond is one of the remaining representatives of these isolated depressions. The middle portion remained a great swampy morass for ages, and was even such at the time of the coming of the white man. It has since had the timber removed and better drainage instituted. The fine muck soil resulting from the long continued swampy conditions is now the most fertile part of the Flatwoods region.

(NOTE 1. The writer wishes to acknowledge his indebtedness to Dr. J. W. Beede, and Dr. E. R. Cumings, of the Department of Geology of Indiana University, under whose general charge the work of securing the data for the above report was carried on, as part of the work of the Department of Geology, Indiana University. The writer was also aided by his brother, Burton J. Malott, who proved himself valuable in the gathering of certain data.)

(NOTE 2. It was the writer's intention that a contour map accompany this paper, and data was gathered with that end in view, but press of time has precluded its preparation.)

MECHANICAL DEVICE FOR TESTING MERSENNE NUMBERS FOR PRIMES.

THOS. E. MASON.

Lucas,* in a note in "Récréations Mathématiques," gives a method of testing numbers of the form $2^{4q+3}-1$ for primes. The purpose of this note is to show how the labor of that method can be shortened, and how a machine could be constructed which would do most of the labor. If such a machine were constructed the labor of verifying the Mersenne numbers would be reduced to hours where it now requires weeks and months, for example, for numbers like $2^{127}-1$.

Lucas makes use of the following theorem: In order that the number $p=2^{4q+3}-1$ shall be prime, it is necessary and sufficient that the congruence

$$\sqrt{-1} \equiv 2 \cos \frac{\pi}{2^{4q+2}}, \pmod{p},$$

shall be satisfied, that is, that

$$\sqrt{-1} \equiv \sqrt{2 - \sqrt{2 + \sqrt{2 + \sqrt{2 + \dots}}}}, \pmod{p},$$

shall be verified after the successive removal of the radical. In other words, if we form the set of numbers V_n ,

$$V_0=1, V_1=3, V_2=7, V_3=47, V_4=2207, \dots,$$

such that each after the second is the square of the preceding diminished by 2 units, and then consider only the residues, modulo p , if the residue of the number V_n , where $n=4q+2$, is zero the number p is prime.

The process of Lucas makes use of the binary system of numeration. In this system multiplication consists simply in the longitudinal displacement of the multiplicand. It is evident also that the residue of the division of 2^m by 2^n-1 is equal to 2^r , r designating the residue of the division of m by n ; consequently, in trying 2^7-1 , it is sufficient to operate upon numbers having at most 7 of the figures 0 or 1. Figure I gives the calculation of V_4 deduced from the calculation of V_3 by the formula

$$V_4 = V_3^2 - 2 \pmod{2^7 - 1};$$

the dark squares represent the units of different orders of the binary system

*Lucas, Récréations Mathématiques, Vol. 2, pp. 230-235.

and the white squares the zeros. The first line is the residue of V_3 ; the 7 lines numbered 0 to 6 represent the residues (mod 2^7-1) of the partial products in




FIG. 1.

squaring V_3 ; the lines below, numbered 0 and 1 represent the addition of the partial products above and reduction modulo 2^7-1 , and line 2 represents the addition of 0 and 1; the single line below gives the residue of the square of V_3 with 2 subtracted, which is the residue of V_4 . In order to complete the test of 2^7-1 it is necessary to find V_6 . If the residue of V_6 is zero, then 2^7-1 is prime. This briefly is the plan as given by Lucas except that he used a different illustrative example.

In order to test a large number, say $2^{127}-1$, it would be necessary to make 126 of these square tables such as Figure I, each having 127^2 small squares. This would entail considerable labor and require a great deal of time. A simple device will reduce the labor of writing each line in the large square (A) to counting. Let us set down the work of squaring a number written in the binary system, for example, 13, which in the binary system is 1101.

$$\begin{array}{r}
 \hline
 1011 \qquad (S) \\
 \hline
 1101 \\
 1101 \\
 \hline
 1101 \\
 1101 \\
 1101 \\
 \hline
 10101001
 \end{array}$$

Now, if we write the number with the digits in reverse order on a slip of paper (S) and place it above the number itself as shown above, we see that the digits which occur in the third column of the partial products are the digits which come together, or correspond, when the slip of paper (S) is placed so that the first digit of the number, when digits are reversed, is placed over the third column. This is possible in no other scale, because the product of two digits in the binary scale does not give a number of more than one place. We can give the following rule for squaring a number written in the binary scale:

Write the number with the digits equally spaced and write the same number with the digits reversed on a slip of paper, using the same spacing. Place the slip of paper above the number so that the first digit in the reversed order comes above the last digit of the number. Move the slip of paper a single space to the left each time. Count the correspondences at each step. The number of correspondences at each step is the number which belongs in that place in the result which is immediately beneath the first digit on the slip. Continue this until there are no more correspondences.

It is easily seen that by means of the above rule the process described by Lucas can be followed out by counting the correspondences and will lead to the result in the lines marked (B) in Figure I, without having to write the part (A).

It would be possible to construct a machine which would have two parallel bars in which could be set pins for the places where 1 occurs in the number. The pins on one bar would be in reverse order. The bars could be turned over and the number of pins striking could be recorded automatically. At the same time one bar could be moved along one place and be in readiness for the next turn. From the machine then would come the data for compiling the part (B) of Figure I. Or, a more complicated machine could be constructed which would give the part (C) at once. This would so shorten the work of testing the Mersenne numbers that it would be possible to check the results on all of them again with a reasonable expenditure of time.

*Purdue University,
LaFayette, Indiana.*

SOME PROPERTIES OF BINOMIAL COEFFICIENTS.

A. M. KENYON.

§1.

The binominal coefficients of the expansion

$$(x + y)^k = \binom{k}{0}x^k + \binom{k}{1}x^{k-1}y + \binom{k}{2}x^{k-2}y^2 + \dots + \binom{k}{k}y^k$$

were known to possess a simple recursion formula

$$(1) \quad \binom{k}{n} + \binom{k}{n+1} = \binom{k+1}{n+1} \quad k, n = 0, 1, 2, 3, \dots$$

by means of which Pascal's Triangle*

	$n = 0$	$n = 1$	$n = 2$	$n = 3$	$n = 4$	<i>etc.</i>
$k = 0$	1					
$k = 1$	1	1				
$k = 2$	1	2	1			
$k = 3$	1	3	3	1		
$k = 4$	1	4	6	4	1	
<i>etc.</i>	—	—	—	—	—	—

could be built up, before Newton showed that they are functions of k and n :

$$(2) \quad \left. \begin{aligned} \binom{k}{n} &= \frac{k(k-1) \dots (k-n+1)}{n!}, \quad n=1, 2, 3, \dots \\ \binom{k}{n} &= 1 \end{aligned} \right\} \quad k=0, 1, 2, \dots \quad n=0$$

A great number of relations involving binomial coefficients have been discovered**, some of the most useful of these are

$$(3) \quad n \binom{k}{n} = k \binom{k-1}{n-1}; \quad \binom{k}{n} = \frac{k}{k-n} \binom{k-1}{n}; \quad \binom{k}{n} = 0 \text{ if } n > k.$$

*See Chrystal: Algebra I, p. 81.

**See Chrystal: Algebra II, Chaps. XXIII, XXVII. Hagen: Synopsis der hoeheren Mathematik, p. 64; Pascal: Repertorium der hoeheren Mathematik I, Kap. II, Sec. 1.

From (1) and (3) it follows that $\binom{k}{n}$ satisfies the linear difference equation

$$(n + 1)f(n + 1) + (n - k)f(n) = 0$$

It is well known that the sum of the coefficients $(x + y)^k$ is 2^k and that the sum of the odd numbered coefficients is equal to the sum of the even numbered ones; the following are perhaps not so well known:

(4) If, beginning with the second, the coefficients of $(x - y)^k$ be multiplied by $c^n, (2c)^n, (3c)^n, \dots, (kc)^n$ respectively; c being arbitrarily chosen different from zero, the sum of the products will vanish for $n = 1, 2, 3, \dots, k - 1$ but not for $n \geq k$, e. g.

$$\begin{array}{cccccccc} k = 8 & -8, & 28, & -56, & 70, & -56, & 28, & -8, & 1 \\ c = 2 & 2^n, & 4^n, & 6^n, & 8^n, & 10^n, & 12^n, & 14^n, & 16^n \end{array}$$

The sum of the products vanishes for $n = 1, 2, \dots, 7$; but not for $n > 7$; for $n = 8$ it is 10,321,920.

(5) If the first k coefficients of $(x - y)^{k+1}$ be multiplied term by term, with $k^n, (k - 1)^n, (k - 2)^n, \dots, 1^n, (n, k = 1, 2, 3, \dots)$ the sum of the products will be

$$(-1)^{k+n} \text{ if } n < k \quad \text{and } (k + 1)! - 1 \quad \text{if } n = k + 1;$$

in particular

$$k^k \binom{k+1}{0} - (k-1)^k \binom{k+1}{1} + (k-2)^k \binom{k+1}{2} - \dots + (-1)^{k-1} \binom{k+1}{k-1} = 1$$

e. g. take $k = 5$.

$$\begin{array}{cccccc} 1, & -6, & 15, & -20, & 15, & \\ 5^n, & 4^n, & 3^n, & 2^n, & 1^n, & \end{array}$$

The sum of the products is +1, -1, +1, -1, +1, 719, for $n = 1, 2, 3, 4, 5, 6$, respectively.

Both (4) and (5) are special cases of

(6) If the coefficients of $(x - y)^k, (k = 1, 2, 3, \dots)$ be multiplied term by term by the n th powers $(n = 0, 1, 2, \dots)$ of the terms of any arithmetic progression with common difference $d \neq 0$, the sum of the products will vanish if $n < k$; will be $(-d)^k (k!)$ if $n = k$; and if $n = k + 1$ will be the product of this last result and the sum of the terms of the arithmetic progression.

e. g. take $k = 6, d = -1$, a.p., 4, 3, 2, etc.

$$\begin{array}{ccccccc} 1, & -6, & 15, & -20, & 15, & -6, & 1 \\ 4^n, & 3^n, & 2^n, & 1^n, & 0^n, & (-1)^n & (-2)^n \end{array}$$

The sum of the products vanishes for $n = 1, 2, 3, 4, 5$, but not for $n > 5$; for $n = 6$, it is $(-1)^6 (6!) = 720$; and for $n = 7$, it is $720(4 + 3 + 2 + 1 + 0 - 1 - 2) = 5040$.

The third conclusion of (6) shows that if

$$(I) \quad a + (a + d) + (a + 2d) + \dots + (a + kd)$$

and

$$(II) \quad \binom{k}{0} a^k - \binom{k}{1} (a + d)^k + \binom{k}{2} (a + 2d)^k - \dots + (-1)^k \binom{k}{k} (a + kd)^k$$

be multiplied term by term and the $(k + 1)$ products be added, the result will be the same as though (II) be multiplied through by the terms of (I) in succession and the $(k + 1)^2$ products be added; e.g. take $k = 4, a = 1, d = 2$

$$(I) \quad 1 \quad , \quad 3 \quad , \quad 5 \quad , \quad 7 \quad , \quad 9$$

$$(II) \quad 1 \cdot 1^4 \quad , \quad -4 \cdot 3^4 \quad , \quad 6 \cdot 5^4 \quad , \quad -4 \cdot 7^4 \quad , \quad 1 \cdot 9^4$$

$$1 \quad - \quad 972 \quad + \quad 18750 \quad - \quad 67228 \quad + \quad 59049 = 9600$$

	$1 \cdot 1^4$	$-4 \cdot 3^4$	$6 \cdot 5^4$	$-4 \cdot 7^4$	$1 \cdot 9^4$	
1	1	-324	3750	-9604	6561	384
3	3	-972	11250	-28812	19683	1152
5	5	-1620	18750	-48020	32805	1920
7	7	-2268	26250	-67228	45927	2688
9	9	-2916	33750	-86436	59049	3456
	25	-8100	+93750	-240100	+164025	9600

§2.

The properties noted above, and many others, can be made to depend upon those of the sum

$$(1) \quad S(k, n) = \sum_{i=0}^k (-1)^i \binom{k}{i} i^n \quad k, n = 0, 1, 2, 3, \dots$$

It is readily shown that

$$(2) \quad S(k, n) \text{ vanishes for } k > n$$

$$(3) \quad \left. \begin{aligned} S(k, n) &= -k \sum_{i=k}^n \binom{n-1}{i-1} S(k-1, i-1) \\ &= - \sum_{i=k}^n \binom{n}{i-1} S(k-1, i-1) \end{aligned} \right\} k, n = 0, 1, 2, 3, \dots$$

whence $S(k, n)$ is divisible by $k!$ and in fact $S(n, n) = (-1)^n n!$ Also, since $S(1, n) < 0$, it follows that for fixed k , $S(k, n)$ preserves a constant sign (or vanishes) for all values of n ; and this sign is the same as that of $(-1)^k$.

These numbers possess a recursion formula

$$(4) \quad S(k, n) = k[S(k, n-1) - S(k-1, n-1)] \quad n, k = 0, 1, 2, \dots$$

by means of which may be constructed,

A TABLE OF VALUES OF $S(k, n)$

	$k=0$	$k=1$	$k=2$	$k=3$	$k=4$	$k=5$	$k=6$	$k=7$	$k=8$
$n=0$	1								
$n=1$	0	-1							
$n=2$	0	-1	2						
$n=3$	0	-1	6	-6					
$n=4$	0	-1	14	-36	24				
$n=5$	0	-1	30	-150	240	-120			
$n=6$	0	-1	62	-540	1560	-1800	720		
$n=7$	0	-1	126	-1806	8400	-16800	15120	-5040	
$n=8$	0	-1	254	-5796	40824	-126000	191520	-141120	40320

Subtract any entry from the one on its right, multiply by the value of k above the latter.

- (5) $\sum_{k=0}^n S(k, n) = (-1)^n$ $\sum_{k=2}^n S(k, n) = 1 + \cos n\pi$
- (6) $\sum_{k=1}^n \frac{S(k, n)}{k} = 0$ $n = 2, 3, 4, \dots$
- (7) $\sum_{i=k}^n \binom{n+1}{i} S(k, i) = (k+1) \sum_{i=k}^n \binom{n}{i} S(k, i)$
- (8) $\sum_{i=k}^n \binom{n}{i} S(k, i) = S(k, n) - S(k+1, n)$

Setting $n = k + 1$ in (7) we obtain

$$(9) \quad S(k, k+1) = \binom{k+1}{2} S(k, k)$$

and similarly we can express $S(k, k+2), S(k, k+3)$, etc. in terms of $S(k, k)$.

From (4)

$$S(k, n) = S(k+1, n) - \frac{1}{k+1} S(k+1, n+1) \quad k, n = 0, 1, 2, 3, \dots$$

By applying this m times, we obtain

$$(10) \quad S(k, n) = \sum_{i=0}^m (-1)^i H_i S(k + m, n + i)$$

$k, n = 0, 1, 2, \dots; m = 1, 2, 3, \dots$

where H_i is the sum of the products of the fractions $1/(k + 1), 1/(k + 2), 1/(k + 3), \dots, 1/(k + m)$, taken i at a time; $H_0 = 1$.

The proof of (6) §1 is as follows. If the first term of the arithmetic progression is zero,

$$\sum_{i=0}^k (-1)^i \binom{k}{i} (di)^n = d^n S(k, n)$$

and this vanishes if $n < k$; is $(-d)^k (k!)$ if $n = k$; and is

$$(-d)^k (k!) [d + 2d + 3d + \dots + kd] \quad \text{if } n = k + 1.$$

If the first term of the arithmetic progression is $a \neq 0$,

$$\sum_{i=0}^k (-1)^i \binom{k}{i} (a + di)^n = d^n \sum_{i=0}^k (-1)^i \binom{k}{i} (x + i)^n$$

where $x = a/d \neq 0$.

If we use the notation

$$f(n, x, k) \equiv \sum_{i=0}^k (-1)^i \binom{k}{i} (x + i)^n$$

expand $(x + i)^n$ by the binomial formula and reverse the order of summation, we obtain

$$(11) \quad f(n, x, k) = \sum_{i=0}^n \binom{n}{i} x^{n-i} S(k, i)$$

Therefore

$$\begin{aligned} f(n, x, k) &= 0 \quad \text{when } n < k, \text{ since all the summands vanish} \\ &= S(k, k) \quad \text{when } n = k \\ &= \sum_{i=k}^n \binom{n}{i} x^{n-i} S(k, i) \quad \text{when } n > k \end{aligned}$$

In particular, when $n = k + 1$

$$f(k + 1, x, k) = (x + \frac{k}{2}) (k + 1) S(k, k) \quad \text{and on putting } a/d \text{ for } x,$$

$d^{k+1} f(k + 1, x, k) = d^k S(k, k) [a + (a + d) + (a + 2d) + \dots + (a + kd)]$
 and from these follow the three conclusions* of (6) §1.

*Chrystal: Algebra II, Sec. 9, p. 183, gives the proof of a slightly less general theorem.

Cauchy: Exercices de mathematiques, 1826, I, p. 49 (23), obtains as a by-product the second conclusion of the theorem for the case $d = -1$, and remarks that it is well known.

§3.

In finding the sum of certain series by the method of differences** it is convenient to express positive integral powers of x in terms of the polynomials

$$(1) \quad \begin{aligned} x^{(n)} &= x(x-1)(x-2)\dots(x-n+1) & n &= 1, 2, 3, \dots \\ x^{(0)} &= 1 \end{aligned}$$

If we set

$$(2) \quad x^n = A(0, n)x^{(0)} + A(1, n)x^{(1)} + \dots + A(k, n)x^{(k)} + \dots + A(n, n)x^{(n)}$$

it is easily shown that

$$(3) \quad A(k, n) = S(k, n)/S(k, k)$$

whence

(4) $A(k, n)$, $k, n = 0, 1, 2, 3, \dots$, vanishes if $n < k$; is always positive if $n \geq k > 0$; in particular $A(n, n) = 1$; and the following relations come from those given in §2 for $S(k, n)$:

$$(5) \quad A(k, n) = \sum_{i=k}^n \binom{n-1}{i-1} A(k-1, i-1) = \frac{1}{k} \sum_{i=k}^n \binom{n}{i-1} A(k-1, i-1)$$

The recursion formula

$$(6) \quad A(k, n) = k A(k, n-1) + A(k-1, n-1)$$

by which may be constructed

A TABLE OF VALUES OF $A(k, n)$

	$k=0$	$k=1$	$k=2$	$k=3$	$k=4$	$k=5$	$k=6$	$k=7$	$k=8$
$n = 0$	1								
$n = 1$	0	1							
$n = 2$	0	1	1						
$n = 3$	0	1	3	1					
$n = 4$	0	1	7	6	1				
$n = 5$	0	1	15	25	10	1			
$n = 6$	0	1	31	90	65	15	1		
$n = 7$	0	1	63	301	350	140	21	1	
$n = 8$	0	1	127	966	1701	1050	266	28	1

To any entry add the product of the one on its right and the value of k above the latter.

**See for example Boole's Finite Differences, Chap. IV.

$$(7) \quad \sum_{i=k}^n \binom{n}{i} A(k, i) = A(k+1, n+1) \quad n > k = 0, 1, 2, \dots$$

$$(8) \quad \sum_{k=1}^n A(k, n) S(k-1, k-1) = 0 \quad n = 2, 3, 4, \dots$$

Inversely, since

$$x^{(n+1)} = x(x-1)(x-2)\dots(x-n) \quad n = 0, 1, 2, \dots$$

if we set

$$(9) \quad x^{(n+1)} = x[B(0, n)x^n - B(1, n)x^{n-1} + \dots + (-1)^k B(k, n)x^{n-k} + \dots + (-1)^n B(n, n)]$$

it is evident that $B(0, n) = 1, n = 0, 1, 2, \dots, B(k, n) =$ the sum of the products of the numbers $1, 2, 3, \dots, n$, taken k at a time; in particular $B(k, k) = k! = (-1)^k S(k, k)$ and $B(k, n) = 0$ if $k > n$. For convenience define $B(p, n) = 0$, if p is a negative integer.

If we multiply both sides of

$$x^{(n)} = x[B(0, n-1)x^{n-1} - B(1, n-1)x^{n-2} + \dots + (-1)^{n-1}B(n-1, n-1)]$$

by $x-n$, and equate the coefficients of x^{n-k} , we obtain the recursion formula

$$(10) \quad B(k, n) = B(k, n-1) + n B(k-1, n-1)$$

by means of which may be constructed

A TABLE OF VALUES OF $B(k, n)$

	$k=0$	$k=1$	$k=2$	$k=3$	$k=4$	$k=5$	$k=6$	$k=7$	$k=8$
$n = 0$	1								
$n = 1$	1	1							
$n = 2$	1	3	2						
$n = 3$	1	6	11	6					
$n = 4$	1	10	35	50	24				
$n = 5$	1	15	85	225	274	120			
$n = 6$	1	21	175	735	1624	1764	720		
$n = 7$	1	28	322	1960	6769	13132	13068	5040	
$n = 8$	1	36	546	4536	22449	67284	118124	109584	40320

Multiply any entry by the number $(n+1)$ of the next row, and add to the entry on its right.

$$(11) \quad B(k, k+n) = \sum_{i=k}^{n+k} \binom{i}{k} B(k+n-i, k+n-1) \quad k, n = 0, 1, 2, 3, \dots$$

The equation

$$B(0, n) x^n - B(1, n) x^{n-1} + \dots + (-1)^n B(n, n) = 0$$

has 1, 2, 3, n, for roots. If we set

$$S_k = 1^k + 2^k + 3^k + \dots + n^k \quad k = 1, 2, 3, \dots$$

and solve Newton's formulae* we obtain

$$B(k, k) B(k, n) = \begin{vmatrix} S_1 & 1 & 0 & 0 & \dots & 0 \\ S_2 & S_1 & 2 & 0 & \dots & 0 \\ S_3 & S_2 & S_1 & 3 & \dots & 0 \\ S_4 & S_3 & S_2 & S_1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ S_k & S_{k-1} & S_{k-2} & S_{k-3} & \dots & S_1 \end{vmatrix} \quad k, n = 1, 2, 3, \dots$$

This determinant vanishes when $k > n$.

Inversely,

$$S_k = \begin{vmatrix} B(1, n) & B(0, n) & 0 & \dots & 0 \\ 2B(2, n) & B(1, n) & B(0, n) & \dots & 0 \\ 3B(3, n) & B(2, n) & B(1, n) & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ kB(k, n) & B(k-1, n) & B(k-2, n) & \dots & B(1, n) \end{vmatrix}$$

$k, n = 1, 2, 3, \dots$ (even if $k > n$)

These sums of the powers of the first n natural numbers are connected by the following relations, in which $I(k/2)$ signifies the integral part of $k/2$:

$$\sum_{i=0}^{I(k/2)} \binom{k}{2i+1} S_{2k-1-2i} = 2^{k-1} S_1^k$$

$$\sum_{i=0}^{I(k/2)} \frac{2k+1-2i}{1+2i} \binom{k}{2i} S_{2k-2i} = (2n+1) 2^{k-1} S_1^k$$

whence

$$\sum_{i=0}^k c_i \binom{k}{i} S_{2k-i} = 0 \quad \text{where } c_i = \frac{2k+1-i}{1+i} \text{ when } i \text{ is even}$$

$$= -(2n+1) \text{ when } i \text{ is odd}$$

*See, for example, Ca'ori's Theory of Equations, pp. 85-86.

†Stern, Crelle's Journal, Vol. 84, pp. 216-218.

Also

$$\sum_{i=0}^k \binom{k+1}{i} S_i = ** (n+1)^{k+1} - 1$$

Relations between the A's and the B's:

$$x^m = \sum_{i=1}^m A(i, m) x^{(i)} \quad m = 1, 2, 3, \dots$$

$$x^{(i)} = \sum_{j=0}^{i-1} (-1)^j B(j, i-1) x^{i-j} \quad i = 1, 2, 3, \dots$$

Therefore

$$x^m = \sum_{i=1}^m A(i, m) \sum_{j=0}^{i-1} (-1)^j B(j, i-1) x^{i-j}$$

the coefficient of x^k on the right is

$$\sum_{i=0}^{m-k} (-1)^i A(k+i, m) B(i, k+i-1)$$

and this must vanish $k = 1, 2, 3, \dots, m-1$, and be equal to 1, for $k = m$.

Whence, setting n for $m - k$,

$$\sum_{i=0}^n (-1)^i A(k+i, k+n) B(i, k+i-1) = 0, \quad \begin{cases} k = 0, 1, 2, \dots \\ n = 1, 2, 3, \dots \end{cases}$$

or, setting i for $k + i$, and n for m ,

$$(12) \quad \sum_{i=k}^n (-1)^i A(i, n) B(i-k, i-1) = 0. \quad \begin{cases} k = 0, 1, 2, \dots, n-1 \\ n = 1, 2, 3, \dots \end{cases}$$

Similarly, starting from

$$x^{(m)} = \sum_{i=0}^{m-1} (-1)^i B(i, m-1) x^{m-i}$$

we obtain

$$(13) \quad \sum_{i=0}^n (-1)^i A(k, k+n-i) B(i, k+n-1) = 0, \quad \begin{cases} k = 0, 1, 2, \dots \\ n = 1, 2, 3, \dots \end{cases}$$

This relation may be generalized as follows:

Set

$$C(k, n, p) = \sum_{i=0}^n (-1)^i A(k, k+n-i) B(i, k+n-p)$$

**Prestet, Elements de Mathematique, p. 178.

then directly and by (13)

$$(a) \quad \left. \begin{aligned} C(k,0,p) &= 1 & p &= 0, 1, 2, \dots \\ C(k,n,1) &= 0 & n &= 1, 2, 3, \dots \end{aligned} \right\} k = 0, 1, 2, \dots$$

making use of (10) we obtain

$$(b) \quad C(k,n,p) = C(k,n,p-1) + (k+n-p-1) C(k,n-1,p-1)$$

The left side vanishes when $p = 1$; therefore

$$C(k,n,0) = -(k+n) C(k,n-1,0)$$

By repeating this $(n-1)$ times and noting that $C(k,0,0) = 1$, we obtain

$$(c) \quad C(k,n,0) = (-1)^n (k+1) (k+2) \dots (k+n) \quad \begin{cases} k = 0, 1, 2, \dots \\ n = 1, 2, 3, \dots \end{cases}$$

Setting $p = 2, 3, 4, \dots, n$, in (b), we find

$$(d) \quad \begin{aligned} C(k,n,p) &= 0 && \text{for } p = 1, 2, 3, \dots, n \\ &= k^n && \text{when } p = n + 1 \end{aligned}$$

Therefore for all values of $k = 0, 1, 2, \dots$; and $n = 1, 2, 3, \dots$

$$(14) \quad \sum_{i=0}^n (-1)^i A(k, k+n-i) B(i, k+n-p) = \begin{aligned} &(-1)^n (k+1) (k+2) \dots (k+n) && \text{when } p = 0 \\ &= 0 && \text{when } p = 1, 2, 3, \dots, n \\ &= k^n && \text{when } p = n+1 \end{aligned}$$

Example illustrating (14) for $k = 2, n = 3$.

		$i=0$	$i=1$	$i=2$	$i=3$	
$p = 0$	$A(2,5-i)$	15	-7	3	-1	sums of products
	$B(i,5)$	1	15	85	225	$(-1)^3 3^3 4^3 5$
$p = 1$	$B(i,4)$	1	10	35	50	0
$p = 2$	$B(i,3)$	1	6	11	6	0
$p = 3$	$B(i,2)$	1	3	2	0	0
$p = 4$	$B(i,1)$	1	1	0	0	2^3

In particular, when $p = n$,

$$\sum_{i=0}^n (-1)^i A(k, k+n-i) B(i, k) = 0$$

or, setting $n \rightarrow k$ for n

$$(15) \quad \left. \begin{aligned} \sum_{i=0}^{n-k} (-1)^i A(k, n-i) B(i, k) &= 0 \\ \sum_{i=0}^k (-1)^i A(k, n-i) B(i, k) &= 0 \end{aligned} \right\} \text{provided } n > k = 0, 1, 2, 3, \dots$$

The two sums are equivalent since for $i > k$, $B(i, k)$ vanishes and for $i > n-k$, $A(k, n-i)$ vanishes.

From (15)

$$A(k, n) = \sum_{i=1}^k (-1)^{1+i} A(k, n-i) B(i, k), \quad n > k = 0, 1, 2, \dots$$

whence

$$B(k, n) = \sum_{i=1}^k (-1)^{1+i} B(k-i, n) A(n, n+i), \quad n > k = 0, 1, 2, \dots$$

Solving for the successive A 's and B 's, and for brevity writing A_1, A_2 for $A(n, n+1), A(n, n+2)$ etc., and B_1, B_2 , for $B(1, k), B(2, k)$ etc.,

$$\begin{aligned} A(k, k) &= 1 \\ A(k, k+1) &= B_1 \\ A(k, k+2) &= B_1^2 - B_2 \\ A(k, k+3) &= B_1^3 - 2B_1 B_2 + B_3 \\ A(k, k+4) &= B_1^4 - 3B_1^2 B_2 + 2B_1 B_3 - B_4 + B_2^2 \\ A(k, k+5) &= B_1^5 - 4B_1^3 B_2 + 3B_1^2 B_3 - 2B_1 B_4 + B_5 + 3B_1 B_2^2 - 2B_2 B_3 \\ &\text{etc., etc.} \end{aligned}$$

$$\begin{aligned} B(0, n) &= 1 \\ B(1, n) &= A_1 \\ B(2, n) &= A_1^2 - A_2 \\ B(3, n) &= A_1^3 - 2A_1 A_2 + A_3 \end{aligned}$$

etc., etc., in exactly the same form as the B 's.

$S(k, n)$ satisfies the linear difference equation of order k ,

$$(16) \quad S(k, n+k) - B(1, k) S(k, n+k-1) + \dots + (-1)^i B(i, k) S(k, n+k-i) + \dots + (-1)^k B(k, k) S(k, n) = 0$$

of which the characteristic equation has for roots $1, 2, 3, \dots, k$; and the conditions

$$S(k, n) = 0; \quad n = 1, 2, 3, \dots, k-1; \quad S(k, k) = (-1)^k k!$$

are exactly sufficient to determine the constants. These two equations, therefore, completely characterize

$$S(k, n) = \sum_{i=0}^k (-1)^i \binom{k}{i} i^n$$

In like manner, the difference equation

$$(17) \quad A(k, n+k) - B(1, k) A(k, n+k-1) + \dots + (-1)^i B(i, k) A(k, n+k-i) + \dots + (-1)^k B(k, k) A(k, n) = 0$$

and the conditions

$$A(k, n) = 0, \quad n = 1, 2, 3, \dots, k-1; \quad A(k, k) = 1$$

completely characterize $A(k, n) = \frac{1}{S(k, k)} \sum_{i=0}^k (-1)^i \binom{k}{i} i^n$

$B(k, n)$ satisfies the difference equation of order $2k + 1$,

$$(18) \quad B(k, n + 2k + 1) - \binom{2k+1}{1} B(k, n + 2k) + \dots + (-1)^i \binom{2k+1}{i} B(k, n + 2k + 1 - i) + \dots - B(k, n) = 0$$

of which the characteristic equation is

$$(x - 1)^{2k+1} = 0$$

Whence $B(k, n)$ is a polynomial of degree $2k$ in n , but the $k + 1$ obvious conditions

$$B(k, n) = 0, \quad n = 0, 1, 2, 3, \dots, k - 1, \quad B(k, k) = k^k$$

are not sufficient to determine the constants. It is possible, however, by the successive application of the method of differences, since

$$\Delta B(k, n) = (n + 1) B(k - 1, n)$$

to determine these constants for any particular value of k .

Thus:

$$B(1, n) = \frac{1}{2} (n+1)n$$

$$B(2, n) = \frac{1}{24} (n+1)n(n-1) (3n+2)$$

$$B(3, n) = \frac{1}{48} (n+1)^2 n^2 (n-1) (n-2)$$

etc., etc.

§4.

The properties of

$$f(n, x, k) = \sum_{i=0}^k (-1)^i \binom{k}{i} (x+i)^n \tag{§2}$$

and an application of $\sum_{i=0}^k (-1)^i \binom{k}{i} \frac{1}{x+i}$ in the theory of gamma functions suggests the generalization:

$$(1) \quad f(t, x, k, n) = \sum_{i=0}^k (-1)^i \binom{k}{i} i^n (x+i)^t$$

$k, n = 0, 1, 2, 3 \dots \dots ; t = 0, \pm 1, \pm 2, \dots \dots$

Whence

- (2) $f(0, x, k, n) = S(k, n) \quad k, n = 0, 1, 2, \dots \dots \dots$
- (3) $f(t, x, 0, n) = x^t \quad \text{when } n = 0$
 $\quad \quad \quad = 0 \quad \quad \quad \text{when } n > 0$
- (4) $f(t, x, 1, n) = x^t - (x+1)^t \quad \text{when } n = 0$
 $\quad \quad \quad = -(x+1)^t \quad \quad \quad \text{when } n > 0$

When $t < 0$, this function has poles at $x = -1, -2, \dots \dots \dots -k$, and also when $n + t < 0$, at $x = 0$.

Since $f(t, x, k, n) = \sum_{i=0}^k (-1)^i \binom{k}{i} i^n (x+i)^{t-m} (x+i)^m$

we have the recursion formula

$$(5) \quad f(t, x, k, n) = \sum_{i=0}^m \binom{m}{i} x^i f(t-m, x, k, m+n-i)$$

$t = 0, \pm 1, \pm 2, \dots \dots ; k, n = 0, 1, 2, 3, \dots \dots ; m = 1, 2, 3, \dots \dots$

If t is not negative, we have on setting t for m in (5)

$$(6) \quad f(t, x, k, n) = \sum_{i=0}^t \binom{t}{i} x^i S(k, t+n-i) \quad k, n, t = 0, 1, 2, 3 \dots$$

If $0 < n < k$

$$\sum_{i=0}^k (-1)^i \binom{k}{i} i^{(n)} (x+i)^t = (-1)^n k^{(n)} f(t, x+n, k-n, 0) \quad n = 1, 2, 3 \dots \dots k$$

Whence, making use of (2) §3,

$$(7) \quad f(t, x, k, n) = \sum_{i=0}^n (-1)^i A(i, n) k^{(i)} f(t, x+i, k-i, 0) \quad n = 1, 2, 3 \dots \dots k.$$

In (5), setting $n = 0, m = 1$, and $t+1$ for t :

$$f(t+1, x, k, 0) = f(t, x, k, 1) + x f(t, x, k, 0)$$

but by (7)

$$f(t, x, k, 1) = -k f(t, x+1, k-1, 0)$$

Therefore,

$$(8) \quad x f(t, x, k, 0) = f(t+1, x, k, 0) + k f(t, x+1, k-1, 0), \quad k = 1, 2, 3 \dots$$

In (5) setting $t = 0$:

$$(9) \quad \sum_{i=0}^m \binom{m}{i} x^i f(-m, x, k, n+m-i) = S(k, n)$$

$$k, n = 0, 1, 2, \dots; \quad m = 1, 2, 3, \dots$$

Now $S(k, n)$ vanishes if $k > n$; therefore $f(-m, x, k, n)$ satisfies the linear homogeneous difference equation of order m :

$$(10) \quad \sum_{i=0}^m \binom{m}{i} x^i f(-m, x, k, n+m-i) = 0.$$

$$k > n = 0, 1, 2 \dots \quad m = 1, 2, 3, \dots$$

of which the characteristic equation is

$$(r + x)^m = 0$$

whence the complete solution is

$$(11) \quad f(-m, x, k, n) = (c_0 + c_1 n + \dots + c_{m-1} n^{m-1}) (-x)^n$$

$$m = 1, 2, 3 \dots; \quad n = 0, 1, 2, \dots, k-1; \text{ not for } n \geq k;$$

however, the equation (10) itself will give $f(-m, x, k, n)$ for

$$n = k, k+1, \dots, k+m-1.$$

For $m = 1$, we have

$$f(-1, x, k, n) = c_0 (-x)^n \quad n = 0, 1, 2, 3, \dots, k.$$

and setting $n = 0$, we determine

$$c_0 = f(-1, x, k, 0).$$

setting $t = -1$ in (8)

$$f(-1, x, k, 0) = \frac{1}{x} [S(k, 0) + k f(-1, x+1, k-1, 0)]$$

$$= \frac{1}{x} \text{ when } k = 0$$

$$= \frac{k}{x} f(-1, x+1, k-1, 0) \quad k = 1, 2, 3 \dots$$

whence by repetition, and noting (3)

$$f(-1, x, k, 0) = \frac{k!}{x(x+1)(x+2)\dots(x+k)}^*$$

and

$$f(-1, x, k, n) = \frac{(-x)^n k!}{x(x+1)\dots(x+k)} \quad n = 0, 1, 2, 3 \dots k-1$$

therefore, since by (10), $f(-1, x, k, k) = -x f(-1, x, k, k-1)$,

$$(12) \quad f(-1, x, k, n) = \frac{(-x)^n k!}{x(x+1)\dots(x+k)} \quad n = 0, 1, 2, 3 \dots k, \text{ but not } n > k.$$

Example:

$$\begin{aligned} x(x+1)(x+2)(x+3)(x+4) \sum_{i=0}^4 (-1)^i \binom{4}{i} \frac{i^n}{x+i} &= 24 && \text{when } n = 0 \\ &= -24x && n = 1 \\ &= 24x^2 && n = 2 \\ &= -24x^3 && n = 3 \\ &= 24x^4 && n = 4 \\ \text{but} &&& = 240x^4 + 840x^3 + 1200x^2 + 576x, \quad n = 5 \end{aligned}$$

To find the value of $f(-1, x, k, n)$ for $n > k$, set $m = 1$ in (9) and multiply through by

$$(x+1)(x+2) \dots (x+k) / S(k, k) = \sum_{i=0}^k B(i, k) x^{k-i} / S(k, k)$$

and set

$$\begin{aligned} g(-1, x, k, n) &\text{ for } f(-1, x, k, n) \sum_{i=0}^k B(i, k) x^{k-i} / S(k, k): \\ g(-1, x, k, n+1) &= A(k, n) \sum_{i=0}^k B(i, k) x^{k-i} - xg(-1, x, k, n) \\ & \quad k, n = 0, 1, 2, \dots \end{aligned}$$

Setting $n = k, k+1$, we verify that

$$(13) \quad g(-1, x, k, k+n) = \sum_{j=1}^n (-1)^{j-1} A(k, k+n-j) \sum_{i=j}^k B(i, k) x^{k+j-i-1}$$

holds for $n = 1, n = 2$; and a complete induction shows, on taking account of (14) §3, ($p = n$), that it holds for all positive integral values of n . On

*See Chrystal: Algebra II, Ex. 26, p. 20.

changing the order of summation and replacing $g(-1, x, k, n)$ by its value, we have

$$(14) \quad f(-1, x, k, n) = \frac{\sum_{j=1}^k x^j \sum_{i=1}^j (-1)^{i-1} B(k-j+i, k) S(k, n-i)}{x(x+1)(x+2)\dots(x+k)}$$

$n > k = 0, 1, 2, \dots$

the numerator being a polynomial arranged according to ascending powers of x ; on arranging this in descending powers of x , taking account of (14) §3.

$$(15) \quad f(-1, x, k, n) = \frac{\sum_{j=0}^{k-1} x^{k-j} \sum_{i=0}^j (-1)^i B(j-i, k) S(k, n+i)}{x(x+1)(x+2)\dots(x+k)}$$

$n > k = 0, 1, 2, 3, \dots$

It is obvious that (14) does not hold for $n \leq k$, since in that case $S(k, n-i)$ vanishes, $i = 1, 2, \dots, n$; on the other hand, noting that $B(k, n)$ and $S(k, n)$ both vanish if $k > n$ and taking account of (15), §3, it results that in the numerator on the right side of (15), when $n \leq k$, the coefficient of every power of x vanishes except that of x^n and this turns out to be

$$(-1)^{k-n} B(0, k) S(k, k) = (-1)^n k! \text{ which agrees with (12).}$$

Therefore,

$$(16) \quad \sum_{i=0}^k (-1)^i \binom{k}{i} \frac{i^n}{x+i} = \frac{\sum_{j=0}^{k-1} x^{k-j} \sum_{i=0}^j (-1)^i B(j-i, k) S(k, n+i)}{x(x+1)(x+2)\dots(x+k)}$$

$k, n = 1, 2, 3, \dots$

but for the case where $n \leq k$, (12) is simpler.

Setting $m = 2$ in (11)

$$(17) \quad f(-2, x, k, n) = (c_0 + c_1 n) (-x)^n \quad n = 0, 1, 2, \dots, k-1.$$

Put $n = 0, n = 1$, and determine

$$\begin{aligned} c_0 &= f(-2, x, k, 0) \\ c_1 &= -\frac{1}{x} f(-2, x, k, 1) - f(-2, x, k, 0), \quad \text{which by (7)} \\ &= \frac{k}{x} f(-2, x+1, k-1, 0) - f(-2, x, k, 0) \end{aligned}$$

In (8) set $t = -2, k = 1$

$$f(-2, x, 1, 0) = f(-1, x, 1, 0) + f(-2, x+1, 0, 0)$$

whence by (12) and (3)

$$\begin{aligned} f(-2, x, 1, 0) &= \frac{1}{x^2(x+1)} + \frac{1}{x(x+1)^2} \\ &= \frac{1!}{x^2(x+1)^2} \sum_{i=0}^1 (1+i) B(1-i, 1) x^i \end{aligned}$$

Again, setting $k = 2$ in (8)

$$\begin{aligned} f(-2, x, 2, 0) &= \frac{1}{x} f(-2, x, 1, 0) + \frac{2}{x} f(-2, x+1, 1, 0) \\ &= \frac{2!}{x^2(x+1)^2(x+2)^2} \sum_{i=0}^2 (1+i) B(2-i, 2) x^i \end{aligned}$$

Assume

$$(18) \quad f(-2, x, k, 0) = \frac{k!}{x^2(x+1)^2 \dots (x+k)^2} \sum_{i=0}^k (1+i) B(k-i, k) x^i$$

and a complete induction, on taking account of (11) §3, shows that this holds for all positive integral values of k .

Therefore:

$$\begin{aligned} c_0 &= \frac{k!}{x^2(x+1)^2 \dots (x+k)^2} \sum_{i=0}^k (1+i) B(k-i, k) x^i \\ -c_1 &= \frac{k!}{x^2(x+1)^2 \dots (x+k)^2} \sum_{i=0}^k B(k-i, k) x^i \end{aligned}$$

and

$$(19) \quad f(-2, x, k, n) = \frac{(-x)^n k!}{x^2(x+1)^2 \dots (x+k)^2} \sum_{i=0}^k (1+i-n) B(k-i, k) x^i$$

$k = 0, 1, 2, \dots; n = 1, 2, 3, \dots, k-1$

On computing, by means of (10), the values of $f(-2, x, k, k)$ and $f(-2, x, k, k+1)$, we verify that (19) holds for $n = 1, 2, 3, \dots, k+1$ but not for $n > k+1$,

Therefore,

$$(20) \quad \sum_{i=0}^k (-1)^i \binom{k}{i} \frac{i^n}{(x+i)^2} = \frac{(-x)^n k!}{x^2(x+1)^2 \dots (x+k)^2} \sum_{i=0}^k (1+i-n) B(k-i, k) x^i$$

$k = 0, 1, 2, \dots; n = 0, 1, 2, \dots, k+1; \text{ not } n > k+1$

The corresponding results for $n = k + 2, k + 3$, etc., may be found by putting these values successively for n in

$$(21) \quad f(-2, x, k, n+2) = S(k, n) - 2x f(-2, x, k, n+1) - x^2 f(-2, x, k, n)$$

which results from setting $m = 2$ in (9). The general result may be put into the form

$$(22) \quad f(-2, x, k, n) = \frac{\sum_{j=0}^{2k-2} x^{2k-j} \sum_{i=0}^{k-1} D(i, j, k) S(k-i, n)}{x^2(x+1)^2 \dots (x+k)^2}; \quad k, n = 1, 2, 3 \dots$$

in which the coefficients D , are independent of n :

$$\begin{aligned} D(i, 0, k) &= 1 \quad \text{when } i = 0 \\ &= 0 \quad \quad \quad i = 1, 2, 3 \dots \dots \dots \\ D(0, j, k) &= \sum_{t=0}^j B(t, k-1) B(j-t, k-1) \quad \quad \quad j = 1, 2, 3 \dots \end{aligned}$$

but I have not been able to determine a general formula for $D(i, j, k)$ by means of which to calculate the coefficients of $f(-2, x, k, p)$, $p > k+1$, without first calculating successively those for $n = k+2, k+3, \dots \dots p-1$.

By making use of (10) § 2, (21) may be reduced to

$$(23) \quad f(-2, x, k, n) = \frac{\sum_{j=0}^{2k-2} x^{2k-j} \sum_{i=0}^{k-1} E(i, j, k) S(k, n+i)}{x^2(x+1)^2 \dots (x+k)^2}; \quad k, n = 1, 2, 3 \dots$$

with which compare (16)

Example:

$$\begin{aligned} x^2(x+1)^2(x+2)^2(x+3)^2(x+4)^2 \sum_{i=0}^4 (-1)^i \binom{4}{i} \frac{i^n}{(x+i)^2} &= S(4, n) x^8 + \\ [12 S(4, n) + 8 S(3, n)] x^7 + \\ [58 S(4, n) + 76 S(3, n) + 36 S(2, n)] x^5 + \\ [144 S(4, n) + 272 S(3, n) + 288 S(2, n) + 96 S(1, n)] x^5 + \\ [193 S(4, n) + 460 S(3, n) + 780 S(2, n) + 720 S(1, n)] x^4 + \\ [132 S(4, n) + 368 S(3, n) + 840 S(2, n) + 1680 S(1, n)] x^3 + \\ [36 S(4, n) + 112 S(3, n) + 312 S(2, n) + 1200 S(1, n)] x^2 \\ n = 1, 2, 3 \dots \dots \dots \end{aligned}$$

also:

$$\begin{aligned}
 &= S(4,n) x^8 + [20 S(4,n) - 2 S(4,n+1)] x^7 + \\
 &[170 S(4,n) - 40 S(4,n+1) + 35 S(4,n+2)] x^6 + \\
 &[800 S(4,n) - 340 S(4,n+1) + 60 S(4,n+2) - 4 S(4,n+3)] x^5 + \\
 &[2153 S(4,n) - 1350 S(4,n+1) + 335 S(4,n+2) - 30 S(4,n+3)] x^4 + \\
 &[3020 S(4,n) - 2402 S(4,n+1) + 700 S(4,n+2) - 70 S(4,n+3)] x^3 + \\
 &[1660 S(4,n) - 1510 S(4,n+1) + 476 S(4,n+2) - 50 S(4,n+3)] x^2 \\
 &\quad n = 1, 2, 3 \dots \dots
 \end{aligned}$$

These results are consistent with (20) for $n = 1, 2, 3, 4, 5$ and for $n = 6$ give

$$\begin{aligned}
 &1560 x^8 + 14400 x^7 + 51672 x^6 + 59520 x^5 + 100320 x^4 + 57600 x^3 \\
 &\quad + 13824 x^2.
 \end{aligned}$$

Purdue University.

RADIOACTIVITY OF SPRING WATER.

R. R. RAMSEY.

Since the discovery of the Becquerel rays in 1896 by Henri Becquerel a great amount of work has been done on radioactive bodies; i. e., bodies which give out a radiation which, among other things, renders the air conducting. Madam Curie discovered polonium and radium in 1898. After the discovery of radium a great many workers contributed to our knowledge of radioactive bodies. Radium and polonium are now known to be transformation products in the radioactive series headed by uranium. Besides the uranium-radium series we have the thorium series, the actinium series, and the potassium series in the radioactive list.

Very early in the history of radioactivity, tests were made on ordinary matter to see if all matter is radioactive. Although there is some evidence to show that all matter is radioactive, i. e., is disintegrating, it has been found that a great part of the effect is due to slight traces of radium and other radioactive substances which are mixed with matter. Thus the surface of the earth is covered with slight traces of radium. The exact distribution of radium on the surface of the earth is not known, determinations having been made in a relatively few localities. Besides the scientific interest in the distribution of radium there is another. It has been found that a great many of the celebrated European springs and baths show an unusual amount of radioactivity. The theory has been advanced that the curative properties of these springs are due to the radioactivity of the water.

Table I gives a partial list of the measurements made on noted springs also a short list of ordinary springs, etc.

TABLE I.
RADIOACTIVITY OF NOTED SPRINGS, ETC.


Kings well, Bath, Eng.		173. X10 ⁻¹² Gm. Ra. per liter.
Brembach (Saxe.).....	36000. to 720000.	X10 ⁻¹² Curies per liter.
Schweizergang, Joachimsthal.		98000.
Lake Balaton, Hungary.....	10300. to 36000.	
Potable waters of Mulhause (Alsace).....		2800.
Evaux-les-Bains.	1060. to 2340.	
Evaux-les-Bains gas.....	3440. to 80090.	
Japanese hot springs.....	237. to 13800.	
Colorado Springs, Manitou.....	120. to 4730.	
Colorado Springs, Manitou, gas.	470. to 20500.	
West Canada, Fairmount, Sinclair.....	3500. to 4000.	
Yellowstone Park.	2.26 to	10.4 Mache units.
Yellowstone Park gas.....	6.25 to	118.3 Mache units
Sixty Springs, Tyrol.....	.06 to 89.	Mache units.
Saratoga, N. Y., springs.....	39. to 880.	X10 ⁻¹² Curies per liter.
Saratoga, N. Y., springs gas, Max.....		847.
Williamstown, Mass.	13. to 216.	
Williamstown, Mass., gas.....	759. to 7290.	
Caledonian Springs, near Ottawa, Can.....		15.
St. Lawrence River.....	.25 to	1.1
Sea water.....		.9
Air, Montreal, Cambridge, etc.....		.1

One Mache unit equals 364.X10⁻¹² (Curies per liter).

The radioactivity of water may be due to traces of radium salts dissolved in the water. It may be due to some other product of the uranium-radium series, to radium emanation, usually, or to some product of the thorium or actinium series. The greater amount is usually due to radium or radium emanation dissolved in the water.

In the uranium-radium series (Table No. 2), it will be noted that when one substance changes into another a radiation of α , β , or γ rays, in some cases all three, are given off. This radiation ionizes the air and renders it conducting. The conductivity of the air becomes a measure of the radioactivity of the substance. This is proportional to the rate which a charged body loses its charge.

The ionization produced by the three sets of rays is about in the following proportions: $\alpha = 100\%$, $\beta = 1\%$, $\gamma = .01\%$. The penetrating powers are in the inverse proportion. Electroscopes for radioactive measurements are known as α ray electroscopes, β ray electroscopes, γ ray electroscopes according to the amount of material that must be penetrated by the radiation in order to get into the electroscope. Thus in an α ray electroscope the substance tested is placed in the electroscope or very near to a window covered with a very thin sheet of aluminum or paper. The rays pass in



36. 4

5-h

Figure I.

without absorption and practically all, at least 99%, of the ionization is produced by the α rays. In the β ray electroscopes the radiation must pass through .05 mm. aluminum, which absorbs all the alpha rays and the ionization is produced by the β rays. In the γ ray electroscopes the radiation must pass through 2 mm. of lead, which completely absorbs the α and β radiation leaving the γ rays to produce the ionization. Thus for very weak radioactive bodies the α rays are used, to produce the ionization.

TABLE 2.
URANIUM RADIUM SERIES.

SUBSTANCE.	Radiation.	Half-value Period.	Transformation Constant sec ⁻¹ .	Range of α rays in cms. (15° C.).	Absorption Coefficients.	
					β rays Alum. cm ⁻¹ .	γ rays Lead cm ⁻¹ .
Uranium 1.....	α	5×10^9 years.	4.6×10^{-18}	2.50		
Uranium 2.....	α	2×10^6 years.	1.1×10^{-14}			
Uranium X.....	β, γ	24.6 days.	3.26×10^{-7}		14.4 and 510	72
(Uranium Y).....	β	1.5 days.	5.34×10^{-6}		360	
Ionium.....	α	2×10^5 years.	1.1×10^{-13}	3.00		
Radium.....	α, β	2000 years.	1.1×10^{-11}	3.30	200	
Radium Emanation.....	α	3.85 days.	2.085×10^{-6}	4.16		
Radium A.....	α	3.0 min.	3.85×10^{-3}	4.75		
Radium B.....	β, γ	26.7 min.	4.33×10^{-4}		13 and 91	4-6
Radium C ₁	α, β, γ	19.5 min.	5.93×10^{-4}	6.94	13 and 53	50
(Radium C ₂).....	β	1.4 min.	8.25×10^{-3}		13	
Radium D.....	β	16.5 years.	1.33×10^{-9}		very soft.	
Radium E.....	β, γ	5 days.	1.60×10^{-6}		43	very soft.
Radium F (Polonium).....	α	136 days.	5.90×10^{-8}	3.77		

The substances in parentheses are products not in the direct line of transformation.

Makower & Geiger's Practical Measurements in Radioactivity.

One notes in the radioactive series (Table 2), that the disintegration product of radium is emanation, a gas, which gives off an α particle and changes into Rad. A. This emanation is a gas and obeys all the gas laws. Rad. A has a half value period of three minutes and gives off an α particle and changes into Rad. B. Rad. B has a half value period of 26.8 minutes and gives off β and γ radiation and changes into Rad. C. Rad. C has a half value period of 19.5 minutes, gives off α, β and γ particles and changes into Rad. C₂ and Rad. D. Rad. D has a slow half value period of 16.5 years. This is so slow that the ionization produced by this change can be neglected in com-

parison with the others. Thus some time, about three hours, after the emanation has been placed in a vessel we have Rad. Em. changing through the intermediate products into Rad. D. giving off three α particles, one from Rad. Em; one from Rad. A; and one from Rad. C. This complex radiation has after the first few hours the half value period of the longest of the series, which is that of Rad. Em., 3.85 days. Thus if a quantity of radium emanation gas is placed in an electroscopes the rate of "leak" of the electroscopes in-




FIG. 2.

creases for three hours and then slowly decreases, dropping to one-half value of the maximum in 3.85 days from the time it reached the maximum.

The rise of activity during the first few hours is shown by the curves in Fig. I. These curves are plotted from data given in Rutherford's *Radioactive Substances and Their Radiations*, and in Makower & Geiger's *Practical Measurements in Radioactivity*. The final values (4 hours) are based on the number of ions produced by α particles from the various products. Thus

Rad. Em., 29.1%, Rad. A, 31.4%, Rad. (B) and C, 39.5%. Total, 100%. The line marked Rad. Em. starts initially at 30% and in four hours has diminished to 29.1% according to the half period of 3.85 days. Rad. A initially is zero, because initially emanation alone is placed in the chamber. Curve A rises to half value in three minutes and in 20 or 30 minutes becomes in equilibrium, that is, it disintegrates into Rad. B as fast as it is formed from the emanation. The latter part of the curve is practically a straight line parallel to the curve for emanation.

Rad. B does not give off α particles. The ionization due to the β radiation can be neglected. Rad. B changes into Rad. C whose half period is 19.5 minutes. Thus the curve (B) and C depends upon the amount of Rad. C present. This initially depends upon the formation of Rad. A and B. The




FIG. 3.

curve starts from zero and reaches its equilibrium in about four hours. The total ionization depends upon all three, so the current in the chamber, assuming that all ions capable of being produced by the α particles are used, increases according to the curve $Em. + A + B + C$, which is formed by summing the ordinates of the three curves. This reaches 100% in about three hours. In a chamber of smaller dimensions the effect of the slower electrons will be greater than the above, since a greater number of the high velocity ones will be absorbed by the walls of the chamber before they have produced their maximum number of ions.

The quantity of emanation gas associated with or occluded in, or in equilibrium with, a quantity of radium has been found to be directly proportional to the mass of radium. This is so true that the amount of emanation in equilibrium with one gram of radium has been measured very exactly and is called the curie. Thus one gram of old radium contains or is in equilibrium with one curie of radium emanation gas. The volume of this gas under standard conditions is .62 cu. mm.

To collect this gas the radium is put into solution, boiled and the gas diluted with air is collected over mercury and then introduced into the electroscope. The radium solution after standing one month is again in equi-

librium with the emanation and can be used again. By noting the ionization current or the "leak" of the electroscope other samples of radium can be compared with the first by putting sample No. 2 through the same process. The Bureau of Standards at Washington is prepared to standardize radium solutions by comparing them with a standard in its possession.

If no standard is at hand the electroscope can be standardized by using Duane's empirical formula. (Le Radium Vol. XI, P. 5, 1914; Ann. der Phys. Vol. 38, P. 959, 1912; Compt. Rendus Vol. 150, P. 1421, 1910; Jour. de Phys. Vol. 4, P. 605, 1905), which is,

$$e = \frac{i_0}{2.49 \times 10^6 (1 - 0.517 S/V)} \text{ curies.}$$

or,

$$e = \frac{i_{\max.}}{6.31 \times 10^6 (1 - 0.572 S/V)} \text{ curies.}$$

Where, e = amount of emanation in the electroscope.

i_0 = initial current, expressed in E. S. units.

i_{\max} = maximum current (current at end of three hours) expressed in E. S. units.

S = inside surface of ionization chamber of electroscope.

V = volume of ionization chamber.

This equation applies to a cylindrical ionization chamber with a central rod. The volume of the chamber must be about one liter and the height is from one to three times the diameter.

The ionization chamber can be a cylindrical metallic chamber with an insulated rod extending through the center. This rod can be connected to an electrometer or to an electroscope in order to determine the potential of the rod. For very delicate measurements of small amounts of emanation a sensitive electroscope is better than an electrometer. In an electroscope the ionization current, i , is measured by knowing the capacity, C , of the electroscope; the change of potential, dV , of the insulated rod, in the time, t ; according to the equation,

$$i = \frac{C dV}{t}.$$

In order to measure a small current in a short time, C , the capacity of the electroscope must be small and dV , the change of potential, must be

small. Therefore we wish a sensitive electroscope of small capacity. The cylindrical chamber is in reality a cylindrical condenser. Therefore we want the diameter of the rod small compared to the diameter of the cylinder, also the collar holding the insulating material should be short and have a large diameter compared to the rod. In short the dimensions of the insulated system should be as small as rigidity and other considerations will permit.

To make the electroscope sensitive the "gold leaf" should be very light and narrow. With Dutch leaf a strip between one and two millimeters wide will give a change of one mm. for a change of potential of five volts. By




FIG. 4.

using a reading microscope with graduated eye piece a fraction of a volt can be detected. The electroscope should be made of brass tubing the thickness of whose walls is one mm. or more. Where such material is not to be had sheet tin can be used and will give satisfactory results, especially if used in a laboratory free from penetrating radiation. I will describe one which I made at a cost of a few cents and have found to give consistent results when compared with one made in Germany which cost \$50.

The cylinder A, Fig. 2, is a 1 lb. coffee can. B, is a tin can made in a local tin shop, 3x3x5 inches with a lid at the top. The lid of the coffee can and the lid of the rectangular can are soldered together and a short cylinder

of brass, S, is soldered in a hole in the center. In this short cylinder the central rod system, which can be made of two or more sections, is insulated by pouring melted sulphur into the cylinder while the rod is supported in proper position by a cork, Fig. 3, which extends a short distance into the cylinder. After a few hours the cork can be removed, leaving the sulphur plug. In melting the sulphur care must be used not to get the sulphur too




FIG. 5.

hot or to burn it. The melted sulphur should be a clear amber liquid. If the sulphur takes the waxy condition it should be discarded. The leaf is attached to a narrow plate which is attached to the lower end of the rod. The leaf which is a narrow strip of Dutch foil should be attached to the plate so as to be straight and to swing freely bending at a point near the plate.

The deflection of the leaf is observed through a small window at one

side. A similar window should be placed on the opposite side to admit light. To read the amount of deflection a scale *S*, as shown in Fig. 4, is mounted on one side on or near the back window. A strip of cross-section paper stuck to the glass with paraffin while hot will answer. The paraffin serves two purposes: it sticks the paper and renders the scale translucent. Half way between the scale and the plane of the leaf system a mirror, *M*, is mounted. Through the front window one sees in the mirror images of the plate, *P*, and the foil, *L*, at P^1 and L^1 . These images are in the same plane as the scale, *S*, which can be viewed by looking over the mirror, *M*. The position of L^1 can be read on the scale, *S*, and at the end of a convenient interval of time its position can be noted again. By comparing the two positions with the calibration curve of the electroscope the change of potential, dV , can be obtained.

The system is charged by means of a rubber rod through the charging system, *C*, Fig. 2. This consists of an insulated rod with a bent wire connected so as to be in contact with the central rod while charging. While not in use this rod is rotated so as to break connection with the rod and then to come into metallic contact with the case of the electroscope.

Two $\frac{1}{4}$ -inch brass drain cocks are soldered to the emanation chamber to admit the emanation.

The data of the following experiment, Table 3, carried out by two students using the "tin electroscope" and a Schmidt electroscope made by Spindler & Hoyer, Göttingen, will give an idea of the accuracy of this electroscope and also the accuracy of Duane's formula.

TABLE 3.

Electroscope.....	"Tin".....	Schmidt.
Observer.....	F. G. T.....	W. D. S.
Diameter of chamber.....	10.8 cm.....	7.8 cm.
Height of chamber.....	12.1.....	20.3
Volume of chamber.....	1102 cc.....	968 cc.
Surface of chamber.....	594 sq. cm.....	586.6 sq. cm.
Capacity of electroscope.....	17.1 cm.....	6.3 cm.
Observed emanation, Curies per liter.....	206000. $\times 10^{-12}$	200000. $\times 10^{-12}$

The two electroscopes were connected together and connected to a vessel containing emanation and pumped causing the air and emanation to pass in a circle through the three chambers until the three contained emanation of the same density.

Calibration of Leaf.—The instrument can be calibrated by connecting to known potentials and noting the deflections of the leaf. A storage battery of three or four hundred volts is convenient. Readings should be taken for every few volts from 0 to the maximum and a curve plotted. X = deflection, Y = volts. If a large voltage battery is not at hand a 110 volt D. C. circuit can be used making connection to a resistance as in Fig. 5. The voltmeter, V , should be read at the same time that the deflection of the leaf is read. A calibration curve from 0 to 110 volts can then be obtained. For the higher points proceed as follows: Charge the leaf to maximum voltage by means of a rubber rod. A body of small capacity, small compared to the capacity of the electroscope, 1 or 2 cm., say, is mounted on an insulated handle. A coin on a small rubber rod will answer. This is first grounded and then touched to the charged system. The gold leaf falls. The capacity is removed, grounded, and the position of the leaf noted. The operation is repeated until the leaf falls to 0 on the scale.

If C is the capacity of the electroscope, and
 c is the capacity of the coin,
 Q , the quantity of electricity,
 $V_1 V_2$ is 1st, 2nd, potential of the leaf,
 $d_1 d_2$ is 1st, 2nd, deflections of leaf,

$$\begin{aligned} \text{then, } Q_1 &= CV_1 = (C + c)V_2 \\ Q_2 &= CV_2 = (C + c)V_3 \\ Q_n &= CV_n = (C + c)V_{(n+1)} \\ \frac{C + c}{C} &= \frac{V_1}{V_2} = \frac{V_2}{V_3} = \frac{V_n}{V_{(n+1)}}. \end{aligned}$$

The last three or four deflections should be on the part of the scale already calibrated. That is, the potentials should be less than 110 volts. If V_n and V_{n+1} are known by comparing with d_n and d_{n+1} on the calibration curve. Since,

$$\frac{V_n}{V_{n+1}} = \frac{V_{n-1}}{V_n}$$

then,

V_{n-1} can be calculated. V_{n-1} being known, then V_{n-2} can be determined. In like manner all V s can be determined up to V_1 . Knowing V and the corresponding deflection, d , the curve can be extended up to the maximum deflection.

Determination of Capacity.—In the same equations if c is known, that is, if c is a cylindrical condenser, then C can be obtained. Note that C is the capacity of the "leaf" system plus the charging system. Knowing the sum, the capacity of the "leaf" can be had by getting the ratio of the two by an operation similar to the above.

Removing the Emanation Gas from the Solution.—The emanation can be removed from a solution by the boiling method. The solution is boiled, driving off the dissolved gases with the steam. The steam is condensed and the gases are trapped in suitable glass tubes over mercury. The ionization chamber is then evacuated and the emanation is sucked into the electroscope. The entire amount of emanation is placed in the chamber by washing the




FIG. 6.

glass tube with air until the pressure of the ionization chamber of the electroscope is at normal pressure. This method is accurate but requires elaborate apparatus which can be used only in the laboratory.

Where the greatest accuracy is not wished Schmidt's shaking method can be used. (*Phys. Zeit.*, Vol. 6, p. 561, 1905.) This method admits of determinations being made at the spring with apparatus which easily can be carried by the observer. The shaking method consists of taking a known volume of water and shaking it vigorously for two minutes in a closed vessel with a known volume of air. Then the emanation which was originally dissolved in the water is mixed in the air and water in a known proportion, depending upon the temperature of the water. Then this air is pumped through

rubber tubing from the shaking vessel into the ionization chamber and back again to the shaking vessel until the emanation is mixed through the air of both chambers in the same proportion. Knowing the constants of the electroscope and the observed change of deflection of the leaf, the amount of emanation in the ionization chamber is known. Knowing this and the various volumes of air and water the amount of emanation per liter of water can be calculated. The shaking vessel is made of a can with two brass stop cocks soldered into it. One cock is placed near the top the other is placed on the side about half way up. For convenience the position of the lower stop cock can be calculated so that the vessel will hold a certain quantity of water when the vessel is filled full and then placed on a level stand with both stop cocks open. In this manner the volume of the water is determined easily and can be made the same in each experiment. The volume of the air above the water can be had by determining the total volume of the can. To pump the air around a rubber bulb pump such as is used in pyrography outfits answers well. The volume of the air in the tubes and pump must be estimated and used in the calculations.

The formula for calculating the amount of emanation per liter, which can be derived easily in connection with Fig. 6, is as follows:

$$E = \frac{I}{V_1} \left(\frac{V_2 + \alpha V_1}{V_1} \right) \left(\frac{V_2 + V_3 + V_4}{V_4} \right) e,$$

Where V_1 = Volume of water in shaking can, expressed in liters.

V_2 = Volume of air in shaking can, expressed in liters.

V_3 = Volume of bulb, pump, and connection tubes.

V_4 = Volume of ionization chamber.

α = Absorption coefficient of water for radium emanation.

e = Amount of emanation in chamber, V_4 .

E = Amount of emanation per liter of water.

The quantity alpha, α , has been determined experimentally and has been found to depend upon the temperature. The value at any temperature can be had by referring to the curve (Fig. 7). The data for this curve is taken from M. Kofler (Akad. Wiss. Wien, Ber. 121, 2a pp. 2193; Sci. Abs. Vol. 16, 1742, 1913), and Boyle (Phil. Mag., 22, p. 840, 1911.)

As a test of the above equation the following will serve (Table 4). Three tests were made at the spring under the ordinary conditions. The

same electroscope was used, but two shaking cans were used, the larger of which had three stop cocks so that two volumes of water could be had.

TABLE 4.

C. J. S. Spring. August 5, 1914. Temperature of water 12.5° C. Temperature of air 30° C.

	I.	II.	III.
Time of beginning.....	11.03 a. m.....	11.46 a. m.....	2.00 p. m.
Volume of water.....	.707 liters.....	3.00 liter.....	5.00 liter.
Volume of air.....	.943.....	4.10.....	2.10.
Curies $\times 10^{-12}$ per liter.....	425. $\times 10^{-12}$	435. $\times 10^{-12}$	443. $\times 10^{-12}$.

These observations were taken every minute and the mean deflection from 15 minutes to 20 minutes from the time of putting the emanation in the ionization chamber, was used. By referring to an experimental curve (Fig. 8), the maximum deflection per minute or the deflection at the end of three hours was calculated. A better agreement could have been obtained if the interval from the end of one experiment to the beginning of the next had been three hours or more.

Fig. 8 is a curve showing actual observations during a period of three hours taken with a sample of water from Hottle Spring. The observations have been reduced to a scale of 100% for the maximum. The curve marked "Decay A and C" is made by observing the deflections after the emanation has been pumped out. By means of this experimental curve observations at any time can be reduced to the maximum or three-hour values. For exact work the emanation should be placed in the electroscope and allowed to stand for three hours and several observations made and the mean used. The curve is for all practical purposes horizontal from three to four hours.

In all these observations the deflections have been corrected for the natural leak of the electroscope due to the natural ionization of the air.

Before giving results, I shall speak of some factors which may influence the results. Since the emanation gas is dissolved in the water and is removed by boiling or by shaking, care should be used in filling the shaking can. Immerse the can in a pool as close as possible to the source and allow the water to flow in gently. Filling by dipping and pouring with a smaller vessel removes some of the gas. If before the water issues from the ground it trickles over rocks in the presence of air which is not charged with emana-




Fig. 7.

Absorption of Radium Emanation by Water at Various Temperatures.

⊙, Kofler, Akad. Wiss. Wien, Ber. 121, 2a F. 2193.

X, Boyle, Phil. Mag. 22, P. 840, 1911

tion it must lose some of the emanation. This may explain the variation of springs in the same locality.

Observations made at the spring simply show the emanation content of the water. This may be due to three things. The emanation which is continually forming from traces of radium in the soil and rocks through which and over which the water passes is dissolved in the water and passes out with the water. It may be due to radium salts dissolved in the water. Or it may be due to some product of the thorium or actinium series. In the first case the water will show radioactivity by the emanation method and after standing in a closed vessel for a month will not show any emanation. In the second case it may show radioactivity the same day as taken from the spring and after standing a month in a closed vessel it will show more or less emanation than at first. In any case the emanation content after standing one month is equal to the amount of radium dissolved in the water, since one curie is the amount of emanation which is in equilibrium with one gram of radium.

All the observations given below are for the emanation content of the water as taken from the spring or well. These observations were taken from time to time on various springs and wells in Indiana and Ohio. The date of observation, approximate location of the spring, and temperature of the water at the spring is given.

TABLE 5.
SPRINGS.

NAME.	Location.	Date.	Temp. C.	Curies, Per Liter.
Ill. Cent.....	Bloomington.....	Mar. 4, 1914.	12.5°	600.X10 ⁻¹²
Youno.....	Brown County, Indiana.....	Mar. 6, 1914.	16.°	355
J. C. S.....	Two miles southeast of Bloomington.....	Mar. 13, 1914.	10.3°	430
J. C. S. Old.....	Two miles southeast of Bloomington.....	Mar. 14, 1914.	11.5°	660
J. C. S.....	Two miles southeast of Bloomington.....	May 16, 1914.	11.5°	170
Ill. Cent.....	Bloomington.....	May 23, 1914.	12.2°	265
Stone.....	Two miles southwest of Bloomington.....	May 23, 1914.	11.°	77
Weimer.....	Three miles southwest of Bloomington.....	May 23, 1914.	12.3°	175
Hottle.....	Bloomington.....	Sept. 24, 1914.	13.°	650
South.....	Morning Sun, Ohio.....	Aug. 24, 1914.	13.°	420
C. McQ.....	One mile southeast of Morning Sun.....	Sept. 2, 1914.	16.°	560
I. B.....	One-half mile west of Morning Sun.....	Sept. 7, 1914.		100
C. D. McQ.....	One mile west of Morning Sun.....	Sept. 7, 1914.	15.8°	250
C. D. McQ.....	(Wood) one mile west of Morning Sun.....	Sept. 7, 1914.	19.5°	300
W. P. McQ.....	One mile west of Morning Sun.....	Sept. 7, 1914.	17.°	610
C. W.....	Two miles west of Morning Sun.....	Sept. 7, 1914.	19.5°	140
Fal. No. 1.....	One mile northeast Col. C. O.....	Sept. 7, 1914.	17.°	350
Tal. Upper.....	One mile northeast Col. C. O.....	Sept. 7, 1914.	17.°	350

CITY WATER.

Location.	Date.	Temp. C.	Curies Per Liter.
Bloomington, Ind.....	Feb. 24, 1914.	5°	27. X 10 ⁻¹²
Bloomington, Ind.....	Mar. 2, 1914.	Hot.	41.
Indiana University.....	Mar. 2, 1914.	5°	45.
Oxford, Ohio.....	Aug. 12, 1914.	19°	70.
Union City, Ind.....	Aug. 18, 1914.	19°	45.
Celina, Ohio.....	Aug. 20, 1914.	26°	7.

WELLS.


S. R. R., Morning Sun, Ohio.....	Aug. 27, 1914.	15°	95.
J. S. R., Farm. One mile north Morning Sun, Ohio.....	Aug. 27, 1914.		70.
C. McQ. One mile south Morning Sun, Ohio.....	Sept. 2, 1914.	13°	200.
Forest Park. Six miles east Union City, Ind.....	Aug. 18, 1914.	12°	185.
Melted snow water.....			6.
Boiled water.....			00.

In some papers the radioactivity of springs is given in Mache units. One mache unit = 364. X 10⁻¹² curies per liter. (Le Radium, Vol. XI, p. 5, 1914.) In several papers higher ratios have been used, 500. X 10⁻¹² in some cases.

The radioactivity of these springs is not great, but it is high compared with ordinary springs of some localities. There is a great deal of variation from spring to spring, but this may be explained by assuming that some emanation has been given up before issuing from the ground. The variation of the same spring from time to time is dealt with in another paper.

Department of Physics,

Indiana University, December 21, 1914.



A TORNADO AT WATERTOWN, SOUTH DAKOTA, JUNE 23, 1914.

J. GLADDEN HUTTON.

A tornado occurred at Watertown, South Dakota, late in the afternoon of June 23, 1914. A large number of dwelling houses and barns were destroyed, telephone and telegraph poles were razed and many gardens ruined. More than a score of people were more or less seriously injured and a number of others were slightly hurt by flying debris. No one was killed outright, though one child was reported to have died of its injuries.

The writer was passing through the city on June 25th and spent the day collecting data relative to the storm. Had more time been available, further information could have been secured. However, it seems worth while to give a brief report of the tornado, notwithstanding the fact that the data are incomplete.

The Watertown Daily Public Opinion issued June 24th said: "People watched the approach of what looked like an ordinary thunder storm following a hot day* yesterday afternoon. Wind clouds formed about 6:30 o'clock and gradually developed into a heavy line to the north. The first indication of the formation of a cyclone was noticed in the continuous change of the light wind. Those watching next turned their attention to clouds forming fast in the northwest, and as a twister was developing the approach of the cyclone which went through the city was noticed.

"The path of the storm embraced an area about three blocks wide the entire length of the city east and west. The worst section in the south part of town was in the three blocks north and east of the corner of Seventh avenue and Maple street S. From there the cyclone took a course east and a little northerly sweeping everything in its path and wrecking homes and barns between Third and Fourth avenues and Fifth and Sixth streets almost entirely. It continued across Seventh, Eighth, Ninth and Tenth streets E. and between Ninth and Tenth streets reached as far

*Mr. R. Q. Wood cooperative observer at Watertown reported the maximum temperature for June 23, 1914, as 83° and the minimum temperature 54°.




FIG. 1. Map of Watertown, S. D., showing path of tornado of June 23, 1914. See text for explanation.

north as First avenue N., badly wrecking homes on First avenue and Kemp avenue E."

In the issue of the same paper for June 25th there is a brief account of the storm at Goodwin, fourteen or fifteen miles a little south of east of Watertown where houses and barns were damaged. Some damage was also reported at Altamont, about ten miles south of east of Goodwin.

Mr. Ray stated that a thunder storm was approaching against a light east wind. At about 6:30 p. m. some hail fell, after which the temperature rose and a light east wind was blowing. About thirty minutes after the hail ceased falling, he noticed a great turmoil in the clouds and a funnel formed which struck the earth near the South Dakota Central roundhouse. (From this point the course of the tornado is indicated on the map shown in Fig. 1.)

When the tornado passed through the city Mr. Ray was at the Elks' Hall, four blocks north of the path of the storm. He stated that there was no wind where he was standing. After the tornado passed the wind changed to the northwest and blew hard. Fifteen minutes later a heavy shower occurred. The tornado passed through the city in ten or fifteen minutes. Mr. Ray had previously witnessed storms of this kind in Iowa.

Mr. Mitchell, agent for the Rock Island Railroad, stated that he first observed the storm over Pelican Lake, about one mile southwest of Watertown. It was traveling in a northeasterly direction and was drawing up water from the lake. Rock Island train No. 417 was pulling into town from the east at 7:05 p. m. The engineer saw the funnel and backed his train hoping to miss it. The train, however, was caught in the storm and had twenty-five panes of glass broken and the coaches were unroofed. One passenger who jumped from the train was injured by flying debris. The storm struck the city at 6:50 p. m. and was twenty minutes in passing through the city, a distance of one and one-fourth miles.

A number of persons corroborated these statements as to the length of time required for the tornado to pass through the city. Mr. H. Dietz stated that the hail came while a gentle southeast breeze was blowing and that there was little or no wind just before the tornado appeared. He saw the twister coming like a black smoke and it appeared to be about ten feet in diameter at the bottom. There was no rain or thunder or lightning accompanying the storm according to his testimony and this statement was verified by other persons questioned concerning it.

There were varying statements as to the presence of more than one



FIG. 2. Map of eastern Codington County, S. D., showing path of tornado. The heavy dark line beginning at the northeast end of Lake Pelican indicates the path of the principal tornado. A lighter line crossing the heavy line at the eastern boundary of the city represents the path of the second funnel referred to in the text. Mapped for the writer by Mr. H. Mathiesen.

funnel, though a number of people said they observed one or more funnels which did not reach the earth. Mr. Dietz said that after the funnel left the city there was another one southeast of it which was white instead of black and that it dipped up and down but did not reach the earth. Mr. Mathiesen, who prepared the map of the storm's path through Codington County (Fig. 2), said that there were two funnels and his map shows the path of the second one, which, peculiarly, seems to have crossed the path of the main storm. He was at his farm about three miles east of Watertown and witnessed the storm as it passed by.

Mr. J. B. Kintsley saw the cloud just before it reached the city and he said that it seemed to be about the size of a box car and looked like a



FIG. 3. View showing wreckage of house in southwest part of Watertown, S. D. General character of houses indicated.

whirling column of mud. After the tornado passed by all of the clouds in the sky seemed to be rushing after it.

The writer carefully examined the path of the storm from the point where it entered the city to the point where it left the city and passed out into the open prairie. The two outside lines shown in Figure 1 indicate the boundaries of the zone of damage, while the middle line is the locus of points where the greatest destruction occurred. The small arrows indicate the direction in which objects moved as assumed from their position before and after the storm. The writer is aware of the fact that in cases where objects were moved for some distance the arrows may not indicate the direction of movement, but where houses were only moved slightly

from their foundations, and in similar cases, the arrows indicate the actual direction of movement. On the right of the axis of the path objects seem to have moved forward and to the left, while on the left side of the path they moved generally backward and to the right with reference to the advance of the tornado, although there are exceptions to this general statement.

Houses on the edges of the path had their chimneys damaged. In fact the outside lines might be designated the chimney lines. Inside the chimney lines, shingles were removed in patches and usually on the side of the house nearest the axis of the storm. Farther in more shingles were removed, porches were blown away, roofs entirely removed and in



FIG. 4. House with end blown outward.

the middle of the path total destruction occurred, though not at all points. Greater destruction seems to have occurred on the right side of the storm path than on the left, and at some points the axis lies to the left of the middle of the path. The destruction seems to have been greatest where the storm entered the city and where it left it. The light construction of many of the houses in the part of the city traversed seems partly responsible for the damage. (See Fig. 3.)

The following incidents are of interest and may be briefly noted: The violent expansion of air in closed buildings was observed everywhere. Shingles were blown from roofs by the sudden expansion of air in the garrets. Windows were blown outward. Mr. Kintsley, who was in a cellar, said that the southwest window was the first one to blow outward.

Hollow cylindrical porch posts were split in at least one instance. Walls or foundations made of hollow cement blocks or hollow tile failed in many instances. (See Figures 9 and 10.)

The entire end of the house shown in Figure 4 was blown outward. The end of the house may be seen lying in the foreground.

Figure 5 shows a similar condition, though the house was greatly damaged otherwise.

Two boys who were in Oak Park in the southwest part of the city just outside the path of the storm said that when the funnel passed by it




FIG. 5. House in east Watertown, S. D., showing explosive effect of air during passage of tornado.

looked like an elephant's trunk and that hot and cold blasts of air passed over them "sometimes hot enough to roast them and sometimes cold enough to freeze them."

At Goodwin, east of Watertown, clouds of soot rushed from the chimneys "as if everyone had a roaring fire." Here "the storm appeared to stay higher up in the air, though chimneys toppled and smaller buildings were overturned."

Figure 6 is from a photograph taken by Mr. Ward Carr who was at a farm house three miles west of Watertown. The tornado is moving toward the left and seems to be at the forward point of a crescent-shaped cloud. The writer does not know whether this is the squall cloud of the thunder storm or not. The hour-glass shape of the tornado is notable.

Figure 7 is from a photograph said to have been taken by a traveling salesman at the corner of Maple street and Second avenue south. This point is about four blocks north of the axis of the storm path. The other photographs were made by the writer.

The weather map of June 23, 1914, reproduced herewith (Fig. 8), shows the weather conditions prevailing on the morning preceding the storm. A trough of low pressure extends toward the southwest from a low central in Canada between an area of high pressure central in southern Alabama and an area of high pressure on the South Pacific States.



FIG. 6. View of tornado Watertown, S. D., June 23, 1914. Photo by Ward Carr.

All students of meteorology are familiar with the atmospheric conditions which prevail when tornadoes occur as well as with the usual freakish behavior of storms of this type. The writer has not discussed these points for this reason, nor does he wish to make a comparative study of this storm in this report. He has aimed only to state as many of the facts concerning this one meteorological event as he was able to ascertain in the brief time at his disposal, trusting that they may add a small part to the great fund of information already recorded concerning these small but violent atmospheric disturbances.

State College,

Brookings, South Dakota.




FIG. 7. View of tornado cloud, Watertown, S. D., June 23, 1914. Name of photographer unknown




FIG. 8. Weather map of June 23, 1914. Conditions preceding Wauertown, S. D., tornado



FIG. 9. Collapse of hollow tile structure. Watertown, S. D., June 23, 1914.



Fig. 10. The house from the foundation in the foreground was carried bodily across the street and then dropped to the earth and broken to bits. Wreckage may be seen in front of cottage on the right.



FIG. 11. General view of wreckage where destruction was greatest.
Watertown, S. D., June 23, 1914.



FIG. 12. House moved from foundation. Watertown, S. D., June 23, 1914.



FIG. 13. House moved from foundation. Watertown, S. D., June 23, 1914.



FIG. 14. Effects of tornado, Watertown, S. D., June 23, 1914.



FIG. 15. The tornado passed to the left of this house moving toward the reader. The porch was torn away and deposited in the rear of the house. Watertown, S. D., June 23, 1914.



FIG. 16. House moved from its foundation. Shingles stripped from roof. Tornado passed to the right of it. Watertown, S. D., June 23, 1914.

A SIMPLE FORM OF THE CAREY FOSTER BRIDGE.

J. P. NAYLOR.

Probably there is no better method than the Carey Foster method for comparing nearly equal resistances or for measuring small changes in resistances such, for instance, as those due to change of temperature. While many arrangements have been devised for interchanging the resistances, as is necessary in this method, the following arrangement is believed to be sufficiently novel to make it worth while to present it to the Academy and it is also thought to possess some advantages over other forms of the apparatus.

Its advantages are that it is readily adapted to any width of resistance terminals; the arrangement of connections and relative position of the resistances is such that they can be seen at a glance; and the apparatus is so simple in construction that it can easily be made by anyone who can solder a little and can drill a few screw-holes in the straight copper strips of which it is made up. The whole apparatus can be built up on well varnished hardwood as a base, the copper strips being fastened to the wood base with wood screws. To help the insulation, it has been found to be a good plan, after the screws have been screwed home, to remove them, one at a time, put a drop or two of thick shellac in the hole, and then put the screws back again.


No. 10 or 12 gauge copper strip, two centimeters wide, can be used for the conductors. In the figure, the double, concentric circles represent binding posts and the single circles mercury cups. The latter are made from .41 copper rim-fire cartridge shells.

It may not be generally known that these shells, although quite thin, make very satisfactory mercury cups if soldered directly on top of such copper strips as here mentioned. If blank shells cannot be obtained, the charges may be drawn from the loaded shells. The charges should not be shot out, for the firing pin will nick the shells and cause them to leak. In order to prevent any mercury from coming in contact with the solder and loosening the shells, they and the strips, after soldering is done,


should be thickly coated with shellac varnish which, after drying, should be baked on by heating until it begins to smoke. Mercury cups made in this way have been in use in our laboratory for twenty years and are still doing good service.

The figure practically explains itself. By means of the links, L and L prime, the apparatus can be connected with any calibrated bridge wire, thus adapting it to the Carey Foster method. By changing the thick connectors, C and C prime, to the dotted positions it will be seen, by tracing the connections, that the resistance in the gaps, R and R prime, are interchanged to opposite ends of the bridge wire. These gaps can be adjusted to any width by means of the slotted support shown in the figure. A piece of mica, M, is used to separate the strips at the point of crossing.

As the conductors are so arranged that the arms of the bridge are at all times symmetrical, the dimensions of the apparatus are immaterial. However, a scale of centimeters is added as a suggestion of size for those who may care to construct apparatus of the size figured. It might be added, that apparatus similar to the above has been used at the DePauw physics laboratory for a number of years and has proved very satisfactory for student work.



A Simple Form of the Carey Foster Bridge.



VARIATION OF THE EMANATION CONTENT OF CERTAIN SPRINGS.

R. R. RAMSEY.

While measuring the radioactivity of springs in the neighborhood of Bloomington, Indiana, I found that the measurements made at certain times did not agree with those made at other times. The discrepancy was so great I set out to see if the difference was due to errors or was a real difference. I selected two springs and have tested them once a week for about three months. One, the Hottle Spring, is due north of the square near the corporation line. The other is due west of the square a little beyond the corporation line, across the Illinois Central Railroad from the railroad pumping station. These springs are 1.3 miles apart. Each has a flow of about 15,000 gallons per day. The flow of the Illinois Central Spring has perceptibly lowered during October and November. The Illinois Central Spring issues from the ground at the root of a beech tree through coarse gravel or stones. The Hottle Spring issues from a crack in solid rock.

The results are shown in the following table:

VARIATION OF EMANATION CONTENT OF CERTAIN SPRINGS NEAR BLOOMINGTON, IND. EXPRESSED IN CURIES PER LITER.

Date.	Temp. C.	J. C. S.	Hottle.	Ill. Cent.
Mar. 4	12.0°	X 10 ⁻¹²	X 10 ⁻¹²	600. X 10 ⁻¹²
Mar. 13	10.3°	430.
May 16	11.5°	170.
May 23	12.2°	265.
July 24	330.
Aug. 5	12.5°	425.
Sept. 24	13 and 13°	650.	445.
Oct. 9	15°	530.
Oct. 16	13 and 12.8°	695.	166.
Oct. 23	13.3 and 13°	700.	120.
Oct. 30	13 and 12.7°	665.	20.
Nov. 6	13 and 12.6°	650.	40.
Nov. 13	13 and 12.6°	705.	20.
Nov. 20	13 and 13°	520.	20.
Nov. 26	13 and 13°	550.	33.
Dec. 3	13 and 13°	535.	60.
Dec. 11	13 and 13°	510.	20.
Dec. 18	13 and 13°	450.	00.

The Hottle Spring has remained almost constant, while the Illinois Central has fallen from $600. \times 10^{12}$ curies to almost zero. Four readings taken on another (J. C. S.) spring are added. Taken all together it is noted that in May there was a low value and an increase during September and another decrease October and December. In a rough way an increase coincides with a season of rain and a decrease with dry weather. It may be that due to a small flow there is more splashing and trickling over rocks near the mouth of the spring and the emanation is lost.

Indiana University,
December 21, 1914.

THE CONSTRUCTION OF A RUTHERFORD'S ELECTROSCOPE.

EDWIN MORRISON.

INTRODUCTION.

At the suggestion of Dr. R. A. Millikan the author recently undertook the problem of constructing an electroscope for general laboratory work in radio-active measurements. The general outlines and plans suggested by Dr. Rutherford in his original papers published in the *Philosophical Magazine* and in his work entitled *Radio-active Transformations* have been followed. Suggestions have also been taken from the following works: *Studies in Radio-activity*, by Bragg; *Conduction of Electricity Through Gases and Radio-activity*, by McClung; and *Practical Measurements in Radio-activity*, by Makower and Geiger.

The purposes have been, first, to show, in greater details than the original papers give, the methods of constructing a successful electroscope; and, second, to embody in one instrument as wide a range of experimental work as possible.

CONSTRUCTION.

A diagram of the electroscope is shown in Fig. 1. The dimensions of the gold leaf chamber (E) are 10 x 10 x 10 cm. This chamber is constructed from sheet brass 1.7 mm. in thickness. The four plates for the sides, top and bottom are first carefully jointed by means of a file and then soldered together as shown in Fig. 2 (A). To facilitate the process of soldering two right angle pieces of metal are joined together forming a right angle frame as shown in Fig. 2 (B). When two pieces of the box are to be joined together they are carefully adjusted upon the frame, a few small pieces of solder and soldering fluid are placed along the joint and a pointed flame is directed along the joint in the inside angle until the solder is thoroughly fused. In this way the parts of the electroscope box can be joined together square and straight.

The front side of the electroscope box is a hinged door. This door has a window in it 6.5 cm. square covered with mica. Through this window the gold leaf may be observed by means of a reading microscope. A dia-

gram on the back side of the door is shown in Fig. 3 (A). The mica is held in place by means of four pieces of brass, 1.7 mm. thick and 1.5 cm. wide. These pieces are screwed onto the door in such a way that they not




FIG. 1.

only hold the mica in place over the window but they also form a close fitting rabbeted joint of the door to the box.

The back side of the electrostatic experiment is constructed like the front side, with a mica window of the same dimensions for illumination. This side may be either hinged to the box or it may be held in place by means of two little hooks.

Using the same methods as described above, the ionisation chamber (I) is constructed. No mica windows are needed in the front door or the back side of (this ionisation) chamber. The dimensions of (I) are $12 \times 12 \times 12$ cm., and the thickness of the walls is 1.7 mm. With the exception of the





FIG. 2.

mica window the door to the ionisation chamber is constructed like that of the door to the gold leaf box. The front door should be hinged to the box and the back side can either be permanently soldered to the box or it can be fastened by means of hinges.

The two boxes (E) and (I) are fastened together by screws, the right hand sides being flush with each other as shown in Fig. 1. This arrange-

ment places the condenser in the center of the ionisation box and the gold leaf support is on one side of the box thus giving more free space to the gold leaf.

Through openings in the top and bottom of the electroscop box (E), and the top of (I) a brass rod (D), about 4 mm. in diameter is adjusted being insulated from the boxes by means of two amber plugs (R) and




FIG. 3.

(S). The upper end of the rod (D) is covered by a metal cap (C). The upper opening of the cap is closed by means of an ebonite plug (G). The cap can be removed for charging purposes. The rod (D) may be extended and enlarged by fitting to the upper end a brass cylinder (H). This cylinder acting with the metal cap (C) forms a condenser which increases the capacity of the electroscop. Upon the side of (D) a brass

strip (S), which is about 6 mm. in width, is fastened by means of a small screw at (F). To this strip the gold leaf is fastened by means of wax or shellac. The rod (D) is terminated at its lower end by means of a




FIG. 4.

disk (A), which is 8 cm. in diameter. This disk is screwed onto the end of (D), and larger or smaller disks may be used, thereby varying the capacity of the electroscope. Supported upon a screw (T) which passes through the bottom of the ionisation chamber is a disk (B) the same

size as that of (A). By turning the screw (T) the position of (B) can be made to change through a range of about 8.5 cm. Its distance in cm. or mm. from (A) can be determined by reading the position of the disk (Q) with reference to the scale (Z).

To the bottom of the ionisation chamber four brass rods 1.2 cm. in diameter and 15 cm. in length are securely fastened by means of screws for legs. Leveling screws are adjusted to the lower ends of these legs.

The electroscopes can be converted into an emanation electroscopes by placing stopcocks (M) and (N) in the sides of the ionisation chamber. The active gas can be admitted to the electroscopes by exhaustion through one stopcock and attaching the source of gas to the other stopcock. Or the active gas can be forced into the ionisation chamber by means of a pressure bulb. In testing active gases it is usually necessary to reduce the capacity to the lowest amount possible. This can be done by removing the cylinder (H) and the condenser plate (A). A small rod should be screwed onto the end of (D) in place of the disk (A). Figure four shows the instrument mounted ready for use.

The instrument can be changed into one for measuring the "Variations of the ionisation produced by an alpha particle along its path," by enclosing the condenser plate (A) in a metal box which has a number of short, small, brass tubes passing through the bottom. (See Makower and Geiger's Practical Measurements in Radio-activity, article 32, page 46.) A drawing of the box is shown in Fig. 3 (B) and a cross section in Fig. 3 (C).

After a thorough test it has been shown that this one instrument with its attachments can be used for a wide range of radio-active measurements, and that it is well adapted to general laboratory work.

*Physical Laboratory,
Earlham College.*

I N D E X

A.

	PAGE
Active Members	22
Alcohol Problem, The, In the Light of Coniosis. Robert Hessler.....	85
Alundum, The, Crucible as a Substitute for the Gooch Crucible. George L. Clark	351
Apparatus, An, for Aerating Culture Solutions. Paul Weatherwax...	157
Appropriation for 1913-1914.....	9

B.

B. Fluorescens and B. Typhosus, Antagonism on, in Culture. P. A. Tetrault	161
Binomial Coefficients, Some Properties of. A. M. Kenyon.....	433
Birds' Eggs, The Alba B. Ghere Collection of, Presented to the Museum of Purdue University. Howard E. Enders.....	273
Birds, Public Offenses—Hunting Wild,—Penalty.....	10
Birds, Their Nests and Eggs, An Act for the Protection of.....	9
Birds, Why Do Our, Migrate. D. W. Dennis.....	145
Black Locust, A New Enemy of the. Glenn Culbertson.....	185
Botanical Problems, Some Large. J. C. Arthur.....	267
Business Session, May 29, 1914.....	40
By-laws	7

C.

Carey Foster Bridge, A Simple Form of the. J. P. Naylor.....	485
Chemistry, The Correlation of High School and College. James Brown	355
Chimney Swift, A Note on a Peculiar Nesting Site of the. Glenn Culbertson	279
Civilization, Conservation and. Arthur L. Foley.....	133
Coals, Tar Forming Temperatures of American. Otto Carter Berry...	373
Cold Storage is Practical Conservation. H. E. Barnard.....	101
Committees, Academy of Science. 1915.....	12
Conservation and Civilization. Arthur L. Foley.....	133
Conservation of Human Life, Science in its Relation to the. Severance Burrage	55
Constitution	5
Contents, Table of	3
Corn Pollination, Report on, IV. (Final). M. L. Fisher.....	207
Curators	11

	PAGE
E.	
Earthworms, Notes on Indiana. H. V. Heimburger.....	281
Electroscope, The Construction of a Rutherford's. Edwin Morrison..	491
Emanation Content, Variation of the, of Certain Springs. R. R. Ramsey	489
Executive Committee	11
F.	
Feeble-minded and Delinquent Girl, The. E. E. Jones.....	71
Feeble-minded and Delinquent Boy, The. Franklin C. Paschal.....	63
Feeble-mindedness in the Public Schools. Katrina Myers.....	79
Feeble-mindedness, The Problem of. G. S. Bliss.....	61
Fellows	14
Flatwoods Region, The, of Owen and Monroe Counties, Indiana. Clyde A. Malott	399
Flood Protection In Indiana. W. K. Hatt.....	149
Forest Trees, Notes Upon the Destruction of, in Indiana. Stanley Coulter	167
G.	
Genus Rosellinia, The, in Indiana. Glenn D. Ramsey.....	251
Gravimetric Analysis, Stirring as a Time Saver in. W. M. Blanchard.	349
I.	
Insects, Some, of the Between Tides Zone. Chas. H. Arndt.....	323
K.	
Kentucky Mountains, Changing Conditions in the. B. H. Schockel....	109
M.	
Members	14
Members, Active	22
Mersenne Numbers, Mechanical Device for Testing, for Primes. Thos. E. Mason	429
Minutes of the Spring Meeting.....	39
Minutes of the Thirtieth Annual Meeting.....	45
Mosses, Corrections of the Lists of, of Monroe County, Indiana, I and II. Mildred Nothnagel and F. L. Pickett.....	179
Mosses, The, of Monroe County, Indiana, III. F. L. Pickett and Mildred Nothnagel	181
N.	
Nummularia, Some Species of, Common in Indiana. Claude E. O'Neal.	235

O.		PAGE
Oat Smut in Indiana. F. J. Pipal.....		191
Officers 1914-1915		11
Officers of the Indiana Academy of Science, 1885-1915.....		13
Orthoptera and Orthopteran Habitats, Notes on, in the Vicinity of Lafayette, Indiana. Henry Fox.....		287
Outcrop, Correlation of the, at Spades, Indiana. H. N. Coryell.....		389

P.

Paleobotany, The, of the Bloomington, Indiana, Quadrangle. T. F. Jackson		395
Plants New or Rare to Indiana. No. V. Chas. C. Deam.....		197
Program of the Thirtieth Annual Meeting.....		50
Public Offenses—Hunting Wild Birds—Penalty.....		10

R.

Radioactivity of Spring Water. R. R. Ramsey.....		453
Rust Propagation, Continuous, Without Sexual Reproduction. C. A. Ludwig		219
Rusts, Correlation of Certain Long-cycled and Short-cycled. H. C. Travelbee		231

S.

Science in its Relation to the Conservation of Human Life.....		55
Sewage Disposal. Chas. Brossmann.....		365
Shawnee Mound, Tippecanoe County, as a Glacial Alluvial Cone. Wil- liam A. McBeth.....		385
Snakes, The, of the Lake Maxinkuckee Region. Barton Warren Ever- mann and Howard Walton Clark.....		337
Soils, The Chemical Composition of Virgin and Cropped Indiana. S. D. Conner		359
Spirogyra Dubia, Some Peculiarities in. Paul Weatherwax.....		203
Springs, Variation of the Emanation Content of Certain. R. R. Ram- sey		489
Spring Water, Radioactivity of. R. R. Ramsey.....		453

T.

Table of Contents.....		3
Tornado, A, at Watertown, South Dakota, June 23, 1914. J. Gladden Hutton		471
Trillium Nivale, Stomata of. F. M. Andrews.....		249

V.

Viola Cucullata, A New Leaf Spot of. H. W. Anderson.....		187
Violet, The Primrose-leaved, in White County. Louis F. Heimlich....		213




2



Academy of Science

-75534

W. S. Hoar
9. 1930



AMNH LIBRARY



100170263